



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:


- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

Ceg

Z6.3





Ceg

Z6.3







II
CATALOGVS
PLANTARVM
HORTI ACADEMICI ET AGRI
GOTTINGENSIS
CONSCRIPTVS

A.
JOHANN GOTTFRIED ZINN
M. D. MED. ET BOT. P. P. O. ACAD. REG.
SC. BEROL. SODALI ET SOC. REG.
SC. GOTTING. MEMBRO EXTRAOR-
DINARIO.



GOTTINGAE
SVMTIBVS VIDVAE ABR. VANDENHOECK
1757.

CATALOG
PLANTARVM

HORTI ACADEMICI ET AGRI

GOTTINGENSIS

COMPENDIOSA

JOHANNI GOTTFRIDU NIN

MD. D. MED. ET BOT. P. B. C. ACADE. REG.

SC. BEROL. SOCIET. ET SOC. REG.

SC. GOTTING. ACADE. REG. AGRIC.

1794

IN OFFICINA

GOTTINGAE

PRINTED AT THE UNIVERSITY PRESS

1794

VIRIS ILLVSTRISSIMIS
REGIAE MAIESTATIS
BRITANNICAE
IN
REGIONIBVS GERMANICIS
STATVS MINISTRIS
ET
CONSILIARIIS INTIMIS
DOMINIS SVIS CLEMENTISSIMIS

DEMISSO ANIMO

D. D. D.

A V C T O R.

SECRETARY GENERAL

REPUBLIC OF THE PHILIPPINES

EXECUTIVE ORDER

Ag

763

STATE OF CALIFORNIA

COMMISSIONER OF THE LAND OFFICE

DOMINION OF THE PHILIPPINES

RECEIVED

D. D. D.

ACTS

VIRI ILLVSTRISSIMI

DOMINI CLEMENTISSIMI

Summa, qua Academiam Georgiam
Augustam fovetis, VESTRA indul-
gentia, VIRI ILLVSTRISSIMI, ani-
mum & fiduciam mihi addit, vt hunc
Horti Academici, cuius cura clementissi-
me mihi demandata est, plantarum catalo-
gum, VESTRO nomini inscribam. Muni-
ficentia VESTRA factum est, vt sumti-
bus non spernendis fabrica caldarium
emendata, & ad alendas servandasque
plantas rariores aptior facta, sed & ipse
hortus vario modo ornatus, & intra paucis-
simos

simos annos tot plantis ex omni terrarum
ambitu adlatis ita instructus fuerit, vt ca-
talogus aliquot eorum cephala auctor
nunc prodire possit. Tantam tamque
insignem benevolentiam, qua semper pre-
cibus meis in commodum horti factis in-
dulxistis, demisse veneror, & publico hoc
monumento gratissimam mentem profi-
teor. Vt vero porro benignitate VE-
STRA res nostras adiuuare, hortumque
Academicum ornatorem reddere curae-
que VESTRAE commendatum habere
velis, summa, qua par est, veneratione
obsecro. Ita Deus in reipublicae, in A-
cademiae, &, quae pars eius non exigua
est, in horti commodum saluos & inco-
lumes VOS seruet, & ad ultimos fene-
stratis terminos VOS, & ILLV-
STRISSIMAS VESTRAS FAMI-
LIAS perpetua felicitate ornet. Gottin-
gae d. 14. Apr. MDCCCLVII.

PRAE-



PRAEFATIO.

Dum catalogum plantarum horti & agri Gottin-
gensis edendi consilium sumsi, nam nulla prae-
mittere visum est, quae rationem operis exponant.
Munificentia Illustrissimorum Viroxum Regi Magnae
Britanniae & Electoris Brunsvico-Luneburgensis &
Consiliis intimis, factum est, ut non solum sumtibus
non spernendis Horti Academici vaporarium melius
instructum, & ad alendas servandasque plantas ra-
tiores aptius redditum, sustentaculisque, in quae
ollae disponantur, horto non exiguum ornamentum
& commodum paratum fuerit, sed eadem quoque de-
bemus, si catalogus intra triennii annorum spatium
aliquot centenis plantarum auctior prodire possit.
Auditores autem libellum optabant, qui, dum plan-

P R A E F A T I O.

tai recentior decerpas demonstramus, memoriam
 adiuuare, illisque regere possit, ut in studio Bota-
 nico proprio Marte ulterius progredi queant. Omnem
 ergo operam dedi, ut, commisso illorum inservitum,
 viamque facilem & tutam, qua ad uberiores plan-
 tarum cognitionem pervenire possint, illos docerem,
 animo in id quoque intento, ut libellum, quo & aliis
 in locis & diversis a nostra Florae regionibus uti
 possent, citis traderem. Methodum elegi, quae,
 quantum ipsa conditio Studii Botanici sinit, ad na-
 turalem proxime accedat, & hanc ob rationem ma-
 xime ex parte illam a Perillustri HALLERO in Enu-
 meratione stirpium Helveticarum adhibitam serva-
 vi, ita tamen, ut classes quasdam naturales priori-
 bus adderem, variisque in locis plantas atium
 in ordinem disponerem. Classis cuiuslibet defi-
 nitionem ubique addidi, Juacinctam, quae sa-
 men scopo nostro sufficiat. Generibus va-
 riis, aut propter minutiam partium obscurioribus,
 aut quorum limites confluere videbantur, praemissi
 characteres breves & essentielles, experientia edoctus,
 nimiam difficultatem in investigandis generibus
 tardium saepe tyronibus creare posse, ut a suavis-
 simo studio facile deterteri se patiantur. In plurima

INDEX GENERVM.

A.

<i>Abies</i>	3	<i>Aloides</i>	113
<i>Abrotanum</i>	397	<i>Alópecurus</i>	60
<i>Abstinium</i>	397	<i>Alisanthemum</i>	252
<i>Abutilon</i>	157	<i>Alíne</i>	185
<i>Acalypha</i>	52	<i>Althaea</i>	152
<i>Acanthus</i>	285	<i>Alysson</i>	322
<i>Acer</i>	205	<i>Amaranthus</i>	32
<i>Achillea</i>	411	<i>Amarylís</i>	95
<i>Aconitum</i>	120	<i>Ambrosia</i>	13
<i>Acórus</i>	82	<i>Amethystina</i>	318
<i>Actaea</i>	117	<i>Ammi</i>	271
<i>Adiantum</i>	285	<i>Anforpha</i>	371
<i>Adonis</i>	128	<i>Amýgdalus</i>	145
<i>Adoxa</i>	184	<i>Anacampteres</i>	170
<i>Aethusa</i>	217	<i>Anacyclus</i>	460
<i>Aegilops</i>	72	<i>Anagallis</i>	252
<i>Aegopodium</i>	214	<i>Ananas</i>	91
<i>Aeschynomene</i>	347	<i>Anastatica</i>	321
<i>Agave</i>	100	<i>Archusa</i>	259
<i>Ageratum</i>	406	<i>Andromeda</i>	202
<i>Agrimonia</i>	198	<i>Androsace</i>	254
<i>Agrostema</i>	192	<i>Andryala</i>	434
<i>Agrostis</i>	60	<i>Anemone</i>	132
<i>Aira</i>	63	<i>Anethum</i>	213
<i>Ajuga</i>	290	<i>Angelica</i>	221
<i>Aizoon</i>	166	<i>Anisum</i>	219
<i>Alaternus</i>	16	<i>Anonis</i>	359
<i>Aleca</i>	152	<i>Anthemis</i>	412
<i>Alchimilla</i>	139	<i>Anthericum</i>	105
<i>Alectorolophus</i>	282	<i>Anthospermum</i>	234
<i>Alisma</i>	102	<i>Anthoxanthum</i>	79
<i>Alkekengi</i>	264	<i>Anthyllis</i>	343
<i>Allium</i>	107	<i>Antirrhinum</i>	282
<i>Alnus</i>	3	<i>Aphanes</i>	159
<i>Aloë</i>	96	<i>FF</i>	<i>Apies</i>

INDEX GENERVM

<i>Apias</i>	340		
<i>Apium</i>	213	<i>Baccharis</i>	403
<i>Apocynum</i>	270	<i>Ballote</i>	306
<i>Aquifolium</i>	238	<i>Barba Caprae</i>	119
<i>Aquilegia</i>	120	<i>Basella</i>	232
<i>Arabis</i>	335	<i>Belladonna</i>	265
<i>Arachis</i>	361	<i>Bellis</i>	417
<i>Arbutus</i>	201	<i>Berberis</i>	206
<i>Argemone</i>	386	<i>Bermudiana</i>	93
<i>Arctotis</i>	416	<i>Beta</i>	25
<i>Argemone</i>	115	<i>Betonica</i>	304
<i>Aristolochia</i>	53	<i>Betula</i>	8
<i>Armeniaca</i>	144	<i>Bidens</i>	406
<i>Arnica</i>	425	<i>Bignonia</i>	181
<i>Artedia</i>	225	<i>Biscutella</i>	324
<i>Artemisia</i>	397	<i>Biserrula</i>	340
<i>Arum</i>	83	<i>Bistorta</i>	40
<i>Aruncus</i>	119	<i>Blitum</i>	35
<i>Arundo</i>	70	<i>Boerhaavia</i>	379
<i>Afarum</i>	53	<i>Borrago</i>	260
<i>Asclepias</i>	269	<i>Bofea</i>	20
<i>Asparagus</i>	104	<i>Brassica</i>	333
<i>Asperugo</i>	260	<i>Briza</i>	65
<i>Asperula</i>	229	<i>Bromelia</i>	91
<i>Asphodelus</i>	105	<i>Bromus</i>	68
<i>Aster</i>	418	<i>Browallia</i>	181
<i>Asterocephalus</i>	381	<i>Brunella</i>	300
<i>Astragalus</i>	340	<i>Bryonia</i>	248
<i>Astrantia</i>	222	<i>Buglossum</i>	259
<i>Athamantia</i>	219	<i>Bugula</i>	290
<i>Atactylis</i>	390	<i>Bupias</i>	320
<i>Atriplex</i>	26	<i>Bupthalmum</i>	410
<i>Atropa</i>	265	<i>Bupleurum</i>	219
<i>Avena</i>	68	<i>Butomus</i>	113
<i>Aurantium</i>	146	<i>Buxus</i>	16
<i>Axyris</i>	35		
<i>Asadarach</i>	123	<i>Cacalia</i>	407
		<i>Cassus</i>	161
			<i>Cala-</i>

INDEX GENERVM.

<i>Adamintha</i>	312	<i>Cergaurea</i>	398
<i>Calceolus</i>	85	<i>Cerastium</i>	187
<i>Calceitrapa</i>	394	<i>Cerasus</i>	144
<i>Calendula</i>	415	<i>Ceratocarpos</i>	36
<i>Calla</i>	84	<i>Cerastonia</i>	438
<i>Callitriche</i>	55	<i>Ceratophyllum</i>	58
<i>Caltha</i>	127	<i>Cercis</i>	372
<i>Campanula</i>	242	<i>Cereus</i>	168
<i>Catsphorata</i>	36	<i>Cerithe</i>	256
<i>Cannabis</i>	80	<i>Cetuispina</i>	126
<i>Canna</i>	439	<i>Chaerophyllum</i>	227
<i>Capparis</i>	147	<i>Chamaeclema</i>	441
<i>Capraria</i>	280	<i>Chamaedrys</i>	290
<i>Captifolium</i>	210	<i>Chamaelea</i>	50
<i>Capicum</i>	263	<i>Chamaemelum</i>	412
<i>Cardaminum</i>	204	<i>Chara</i>	54
<i>Catdamine</i>	334	<i>Cheranthus</i>	336
<i>Cardiaca</i>	300	<i>Chelidonium</i>	117
<i>Cardiospermum</i>	205	<i>Chelone</i>	282
<i>Carduus</i>	387	<i>Chenopodium</i>	27
<i>Carex</i>	74	<i>Chenopodiummorus</i>	35
<i>Cassia</i>	391	<i>Chondrilla</i>	431
<i>Carpesium</i>	399	<i>Christophoriana</i>	117
<i>Carpinus</i>	9	<i>Chrysanthemum</i>	413
<i>Carthamus</i>	390	<i>Chrysocome</i>	404
<i>Carum</i>	214	<i>Chrysosplenium</i>	19
<i>Coryophyllata</i>	134	<i>Cicer</i>	358
<i>Cassia</i>	372	<i>Cichorium</i>	417
<i>Cassida</i>	292	<i>Cicuta</i>	217
<i>Cassine</i>	210	<i>Circea</i>	229
<i>Castanea</i>	10	<i>Cirsium</i>	387
<i>Catananche</i>	435	<i>Cistus</i>	148
<i>Citaria</i>	307	<i>Citrus</i>	146
<i>Caucalis</i>	224	<i>Clematis</i>	333
<i>Celastrus</i>	208	<i>Cleome</i>	339
<i>Celosia</i>	34	<i>Clifortia</i>	52
<i>Cekis</i>	20	<i>Clinopodium</i>	312
<i>Cenchrus</i>	73	<i>Clusia</i>	50

INDEX / GENERVM.

<i>Glypeola</i>	322	<i>Groton</i>	331
<i>Gneorum</i>	30	<i>Crucianella</i>	331
<i>Gnicus</i>	390	<i>Gucubalus</i>	388
<i>Gochlearia</i>	324	<i>Cucumis</i>	347
<i>Coffea</i>	354	<i>Cucurbita</i>	346
<i>Coix</i>	73	<i>Cuminoides</i>	377
<i>Colchicum</i>	94	<i>Cunila</i>	304
<i>Colocynthis</i>	347	<i>Cupressus</i>	35
<i>Colutea</i>	350	<i>Cuscuta</i>	337
<i>Gomarum</i>	335	<i>Cyanus</i>	394
<i>Commelina</i>	90	<i>Cyclamen</i>	353
<i>Conium</i>	317	<i>Cydonia</i>	343
<i>Convolvulus</i>	103	<i>Cynapium</i>	317
<i>Convolvulus</i>	338	<i>Cynara</i>	385
<i>Conyza</i>	404	<i>Cynoglossum</i>	358
<i>Corallodendron</i>	361	<i>Cynolurus</i>	70
<i>Corchorus</i>	347	<i>Cyperoides</i>	74
<i>Cpreopsis</i>	408	<i>Cyperus</i>	78
<i>Coriandrum</i>	316	<i>Cypripedium</i>	353
<i>Corispermum</i>	36	<i>Cyrtus</i>	365
<i>Cypnus</i>	334	D.	
<i>Corona imperialis</i>	101	<i>Damasonium</i>	103
<i>Coronilla</i>	349	<i>Daphne</i>	19
<i>Coronopus</i>	335	<i>Datura</i>	240
<i>Corrigiola</i>	45	<i>Daucus</i>	225
<i>Corylus</i>	126	<i>Delphinium</i>	125
<i>Cotinus</i>	308	<i>Dens Canis</i>	102
<i>Cotula</i>	399	<i>Dentaria</i>	331
<i>Cotyledon</i>	169	<i>Dictamnus</i>	316
<i>Crambe</i>	330	<i>Dianthus</i>	195
<i>Crassula</i>	167	<i>Diervilla</i>	212
<i>Crataegus</i>	243	<i>Digitalis</i>	281
<i>Crepis</i>	428	<i>Dipsacus</i>	383
<i>Crinum</i>	311	<i>Dodartia</i>	280
<i>Crichnum</i>	320	<i>Dolichos</i>	363
<i>Crocus</i>	94	<i>Doronicum</i>	415
<i>Crotalaria</i>	361	<i>Dorycnium</i>	344

Draba

INDEX GENERVM :

<i>Draba</i>	322	<i>Festuca</i>	68
<i>Dracocephalum</i>	309	<i>Ficaria</i>	138
<i>Drosera</i>	182	<i>Ficoides</i>	166
<i>Ebolus</i>	209	<i>Ficoides</i>	163
<i>Echinops</i>	384	<i>Ficus</i>	21
<i>Echium</i>	255	<i>Filago</i>	400
<i>Elaterium</i>	246	<i>Filipendula</i>	119
<i>Elatine</i>	282	<i>Foeniculum</i>	213
<i>Eupetrum</i>	15	<i>Foeniculum gracium</i>	364
<i>Ephedra</i>	2	<i>Fragaria</i>	138
<i>Epimerum</i>	90	<i>Fragula</i>	16
<i>Epilobium</i>	199	<i>Fraxinella</i>	710
<i>Epipedium</i>	273	<i>Fraxinus</i>	441
<i>Epipactis</i>	85	<i>Fritillaria</i>	191
<i>Erica</i>	202	<i>Fumaria</i>	374
<i>Erigeron</i>	403	<i>G.</i>	
<i>Eriacephalus</i>	416	<i>Galanthus</i>	94
<i>Eriophorum</i>	79	<i>Gale</i>	12
<i>Ertea</i>	331	<i>Galega</i>	350
<i>Eruca</i>	310	<i>Galeopsis</i>	302
<i>Ervum</i>	358	<i>Galium</i>	232
<i>Eryngium</i>	228	<i>Gazidella</i>	125
<i>Erythraea</i>	328	<i>Genista</i>	360
<i>Erythronium</i>	361	<i>Gentiana</i>	248
<i>Esculus</i>	302	<i>Geranium</i>	171
<i>Evonymus</i>	204	<i>Geum</i>	171
<i>Eupatorium</i>	208	<i>Gladiolus</i>	91
<i>Euphorbia</i>	405	<i>Glaucium</i>	116
<i>Euphrasia</i>	46	<i>Glaux</i>	252
	289	<i>Glechoma</i>	442
		<i>Gleditsia</i>	374
		<i>Glycine</i>	340
		<i>Glycyrrhiza</i>	351
		<i>Gnaphalium</i>	400
		<i>Gnaphalodes</i>	171
		<i>Gomphrena</i>	134
		<i>Gollypium</i>	158
		<i>Gramen</i>	18
		<i>Granadilla</i>	148

INDEX GENERAL

<i>Gratiola</i>	383	<i>Hypsetis</i>	438
<i>Gutierrezia</i>	144	<i>Hypocoum</i> em.	376
<i>Gypsophila</i>	393	<i>Hypericum</i>	442
H.		<i>Hypochaeris</i>	434
<i>Haemanthus</i>	91	<i>Hypopitys</i>	124
<i>Harmala</i>	183	<i>Hyssopus</i>	302
<i>Hedera</i>	23	<i>Hysterophorus</i>	14
<i>Hedysarum</i>	348	I.	
<i>Helenia</i>	417	<i>Jalappa</i>	342
<i>Heliopsis</i>	408	<i>Jasione</i>	244
<i>Helianthemum</i>	440	<i>Jasminum</i>	274
<i>Heliotropium</i>	257	<i>Jatropha</i>	71
<i>Helleborine</i>	86	<i>Iberis</i>	325
<i>Helleborus</i>	125	<i>Illecebrum</i>	44
<i>Helminthotheca</i>	430	<i>Impatiens</i>	273
<i>Helxine</i>	42	<i>Imperatoria</i>	322
<i>Hemerocallis</i>	100	<i>Indigofera</i>	350
<i>Hepatica</i>	192	<i>Intybus</i>	429
<i>Hieracium</i>	223	<i>Inula</i>	412
<i>Hermannia</i>	198	<i>Ipomaea</i>	239
<i>Herniaria</i>	43	<i>Iris</i>	92
<i>Hesperis</i>	337	<i>Isatis</i>	320
<i>Heschera</i>	198	<i>Isopyrum</i>	125
<i>Hibiscus</i>	151	<i>Iva</i>	14
<i>Hieracium</i>	426	<i>Juglans</i>	9
<i>Hippocastanum</i>	204	<i>Jussiaea</i>	171
<i>Hippocrepis</i>	346	<i>Juncus</i>	89
<i>Hippophae</i>	13	<i>Juniperus</i>	5
<i>Hordeum</i>	71	<i>Justicia</i>	285
<i>Homium</i>	311	<i>Lia</i>	93
<i>Homium</i>	296	K.	
<i>Hottonia</i>	253	<i>Kali</i>	37
<i>Hottulus</i>	23	<i>Kalmia</i>	257
<i>Hypocinchus</i>	102	<i>Kleinia</i>	205
<i>Hydrocharis</i>	113	<i>Knautia</i>	323
<i>Hydrocotyle</i>	213	<i>Kanwel</i>	18
<i>Hypocyanus</i>	267		

INDEX GENERVM.

L.			
<i>Laetia</i>	431	<i>Linum</i>	180
<i>Lagoecia</i>	377	<i>Liquidambar</i>	11
<i>Lagurus</i>	61	<i>Liriodendrum</i>	127
<i>Lamium</i>	302	<i>Lithospermum</i>	257
<i>Lampsona</i>	437	<i>Lobelia</i>	244
<i>Lantana</i>	377	<i>Lolium</i>	67
<i>Lapathum</i>	38	<i>Lonicera</i>	210
<i>Lappa</i>	386	<i>Lotus</i>	344
<i>Lapsana</i>	437	<i>Lunaria</i>	323
<i>Larix</i>	3	<i>Lupinus</i>	362
<i>Lascrpitium</i>	228	<i>Lupulus</i>	23
<i>Lathraea</i>	277	<i>Luteola</i>	123
<i>Lathyrus</i>	352	<i>Lychnidea</i>	253
<i>Lavatera</i>	154	<i>Lychnis</i>	192
<i>Lavendula</i>	315	<i>Lycium</i>	254
<i>Laurocerasus</i>	144	<i>Lycopsis</i>	260
<i>Laurus</i>	21	<i>Lycopus</i>	317
<i>Lens</i>	358	<i>Lysimachia</i>	259
<i>Leontiscus</i>	11	<i>Lythrum</i>	197
<i>Leontodon</i>	425		
<i>Leonurus</i>	309		
<i>Lepidium</i>	328		
<i>Lerchia</i>	30		
<i>Leuconium</i>	95		
<i>Libanotis</i>	226		
<i>Ligusticum</i>	218		
<i>Ligustrum</i>	275		
<i>Lilac</i>	274		
<i>Liliasphodelus</i>	100		
<i>Lilium</i>	101		
<i>Limnopenice</i>	55		
<i>Limon</i>	146		
<i>Liponium</i>	376		
<i>Limofella</i>	276		
<i>Linagrostis</i>	79		
<i>Linaria</i>	282		
<i>Linosyris</i>	18		

M.	
<i>Maierana</i>	316
<i>Malva</i>	154
<i>Malus</i>	145
<i>Mariscus</i>	791
<i>Marrubium</i>	306
<i>Matricaria</i>	413
<i>Maurocentia</i>	2101
<i>Mays</i>	73
<i>Medicago</i>	365
<i>Melampyrum</i>	282
<i>Melia</i>	182
<i>Melanthus</i>	271
<i>Melica</i>	64
<i>Melilotus</i>	368
<i>Melissa</i>	312
<i>Melitis</i>	312
<i>Melo</i>	247

INDEX GENERVMJ

Melocactus	161	Nardus	99
Melothria	248	Nasturtium	326
Menispermum	24	Nepeta	307
Menyanthes	253	Nerium	268
Mentha	316	Nicotiana	269
Mercurialis	52	Nigella	124
Mesembryanthemum	163	Nissolia	398
Messpilus	142	Nyctanthes	274
Meum	219	Nymphaea	114
Micropus	154		
Milium	62	O.	
Mimosa	373	Obeliscotheba	409
Mimulus	280	Ocimum	299
Mirabilis	241	Odontites	289
Mitella	198	Oenanthe	215
Moldavica	309	Oenothera	198
Mollugo	44	Olea	275
Mofucella	310	Omphalodes	259
Momordica	246	Onagra	198
Monarda	299	Onobrychis	348
Monotropa	184	Ononis	359
Moritia	276	Onopordon	385
Morus	26	Ophrys	85
Moschatellina	184	Opulus	209
Musa	84	Opuntia	161
Muscari	102	Oreoselinum	221
Myagrum	321	Origanum	315
Myelotis	258	Ornithogalum	106
Myosurus	128	Ornithopodium	346
Myrica	11	Ornithopus	346
Myriophyllum	57	Orobanche	276
Myrrhis	227	Orobus	352
Myrsine	202	Osteospermum	415
Myrtus	142	Osbonna	421
		Oxalis	173
N.		Oxycozum	200
Napaea	157	Oxys	173
Narcissus	95		

P.

INDEX: GENERVM

<i>P.</i>		<i>Phleum</i>	60
<i>Padas</i>	144	<i>Phlemis</i>	301
<i>Paeonia</i>	116	<i>Phlox</i>	253
<i>Palustris</i>	116	<i>Rhylica</i>	50
<i>Palma</i>	83. 84	<i>Rhyllis</i>	212
<i>Panicastrella</i>	73	<i>Rhyfalis</i>	264
<i>Panicum</i>	61	<i>Phyteuma</i>	244
<i>Papaver</i>	114	<i>Phytolacca</i>	439
<i>Parietaria</i>	30	<i>Picris</i>	429
<i>Paris</i>	184	<i>Pimpinella</i>	215
<i>Parkinsonia</i>	373	<i>Pinus</i>	3
<i>Parnassia</i>	124	<i>Pistacia</i>	11
<i>Parthenium</i>	14	<i>Pisum</i>	351
<i>Pastiflora</i>	248	<i>Plantaginella</i>	276
<i>Pastinaca</i>	214	<i>Plantago</i>	236
<i>Pavia</i>	204	<i>Platanus</i>	10
<i>Pedicularis</i>	287	<i>Plumbago</i>	231
<i>Paganum</i>	183	<i>Poa</i>	64
<i>Peireskia</i>	161	<i>Podegraria</i>	214
<i>Pelecinus</i>	240	<i>Polemonium</i>	242
<i>Pentapetes</i>	398	<i>Polyanthes</i>	103
<i>Penasphyllum</i>	335	<i>Polygala</i>	375
<i>Pentapteris</i>	57	<i>Polygonatum</i>	105
<i>Peplis</i>	397	<i>Polygonum</i>	43
<i>Percepier</i>	339	<i>Populago</i>	127
<i>Periploca</i>	269	<i>Populus</i>	7
<i>Persea</i>	145	<i>Portulaca</i>	167
<i>Perficaria</i>	41	<i>Potamogeton</i>	56
<i>Pervinca</i>	268	<i>Potentilla</i>	335
<i>Petafites</i>	404	<i>Poterium</i>	235
<i>Petiveria</i>	418	<i>Praefium</i>	311
<i>Peneedandum</i>	220	<i>Praeanthes</i>	431
<i>Phaca</i>	342	<i>Primula</i>	253
<i>Phalaris</i>	99	<i>Prunus</i>	144
<i>Phar naceum</i>	45	<i>Psidium</i>	144
<i>Phaseolus</i>	362	<i>Rfomalea</i>	343
<i>Phellandrium</i>	216	<i>Rfyllium</i>	236
<i>Philadelphus</i>	247	<i>Preleae similis</i>	207

INDEX. GENERVM.

<i>Pulegium</i>	317	<i>Ruellia</i>	281
<i>Pulmonaria</i>	256	<i>Rumex</i>	36
<i>Pulsatilla</i>	132	<i>Ruscus</i>	24
<i>Punica</i>	144	<i>Ruta</i>	183
<i>Pyrethrum</i>	414		S.
<i>Pyrola</i>	203	<i>Sabina</i>	3
<i>Pyrus</i>	143	<i>Sagittaria</i>	112
	Q.	<i>Salicaria</i>	197
<i>Quercus</i>	239	<i>Salicornia</i>	541
<i>Quercus</i>	9	<i>Salix</i>	61
<i>Quinquefolium</i>	135	<i>Salisolia</i>	371
	R.	<i>Salvia</i>	294
<i>Radiola</i>	181	<i>Sambucus</i>	209
<i>Ranunculus</i>	128	<i>Samolus</i>	245
<i>Raphanus</i>	333	<i>Sanguisorba</i>	235
<i>Rapum</i>	333	<i>Sanicula</i>	205
<i>Rapunculus</i>	244	<i>Santolina</i>	399
<i>Rapuntium</i>	244	<i>Saponaria</i>	193
<i>Rasteda</i>	123	<i>Satureia</i>	313
<i>Rhabarbarum</i>	37	<i>Saxifraga</i>	87
<i>Rhagadiolus</i>	436	<i>Saururus</i>	83
<i>Rhamnus</i>	16	<i>Saxifraga</i>	171
<i>Rhaponticum</i>	391	<i>Scabiosa</i>	380
<i>Rheum</i>	37	<i>Scandix</i>	227
<i>Rhizanthus</i>	288	<i>Schoenus</i>	79
<i>Rhodiola</i>	169	<i>Scirpus</i>	78
<i>Rhus</i>	208	<i>Sclarea</i>	395
<i>Ribes</i>	141	<i>Scolymus</i>	438
<i>Ricinus</i>	51	<i>Scorpioides</i>	347
<i>Rivinia</i>	439	<i>Scorpiurus</i>	347
<i>Robinia</i>	350	<i>Scorzonera</i>	432
<i>Rorilla</i>	182	<i>Scrophularia</i>	285
<i>Rosa</i>	140	<i>Scutellaria</i>	292
<i>Rosmarinus</i>	299	<i>Secale</i>	71
<i>Rubra</i>	219	<i>Securidaca</i>	349
<i>Rubia</i>	234	<i>Sedum</i>	169
<i>Rubus</i>	139	<i>Selinum</i>	221
<i>Rudbeckia</i>	409	<i>Sempervivum</i>	169
			Senecio

INDEX GENERVM.

<i>Senecio</i>	411	<i>Stellaria</i>	53
<i>Senecio</i>	372	<i>Stipa</i>	70
<i>Serpillum</i>	313	<i>Stoechas</i>	315
<i>Serratula</i>	386	<i>Stramonium</i>	340
<i>Serapias</i>	85	<i>Stratiotes</i>	113
<i>Seseli</i>	215	<i>Succisa</i>	380
<i>Siberardia</i>	229	<i>Symphytum</i>	256
<i>Sicyos</i>	248	<i>Syringa</i>	274
<i>Sida</i>	304		T.
<i>Sideritis</i>	257	<i>Tagetes</i>	416
<i>Siegesbeckia</i>	409	<i>Tamarix</i>	203
<i>Silene</i>	189	<i>Tanacetum</i>	396
<i>Silva</i>	222	<i>Taraxacum</i>	426
<i>Silybum</i>	389	<i>Tarchoanthus</i>	14
<i>Sinapis</i>	332	<i>Taxus</i>	2
<i>Sinapisfrum</i>	339	<i>Telephium</i>	206
<i>Sison</i>	218	<i>Terebinthus</i>	11
<i>Sisymbrium</i>	328	<i>Tetragonia</i>	167
<i>Sisyrinchium</i>	93	<i>Tetralix</i>	202
<i>Sium</i>	217	<i>Teucrium</i>	296
<i>Smilax</i>	24	<i>Thalictrum</i>	112
<i>Smyrnium</i>	214	<i>Thelygonum</i>	53
<i>Solanum</i>	261	<i>Thesium</i>	18
<i>Solidago</i>	420	<i>Thlaspi</i>	326
<i>Sonchus</i>	430	<i>Thlaspidium</i>	324
<i>Sophora</i>	372	<i>Thuja</i>	9
<i>Sorbus</i>	142	<i>Thymelaea</i>	19
<i>Sparganium</i>	82	<i>Thymus</i>	315
<i>Spartium</i>	359	<i>Thyffelinum</i>	221
<i>Spermatocoe</i>	234	<i>Tilia</i>	147
<i>Sphondylium</i>	221	<i>Tinus</i>	209
<i>Spinachia</i>	26	<i>Tithymalus</i>	46
<i>Spiraea</i>	217	<i>Tormentilla</i>	135
<i>Squamaria</i>	277	<i>Toxicodendron</i>	208
<i>Stachys</i>	303	<i>Trachelium</i>	245
<i>Stapelia</i>	270	<i>Tradescantia</i>	90
<i>Staphylaea</i>	206	<i>Tragopogon</i>	433
<i>Statice</i>	376	<i>Trogoelinum</i>	215
		<i>Trapa</i>	

INDEX GENERVM.

<i>Trapa</i>	207	<i>Veronica</i>	279
<i>Tribulus</i>	182	<i>Viburnum</i>	209
<i>Tribulus aquaticus</i>	207	<i>Vicia</i>	355
<i>Triclis</i>	44	<i>Viola</i>	268
<i>Tridentalis</i>	292	<i>Viola</i>	272
<i>Trifolium</i>	369	<i>Virga aurea</i>	426
<i>Triglochin</i>	81	<i>Viscago</i>	188
<i>Trigonella</i>	364	<i>Viscum</i>	16
<i>Trinitas</i>	132	<i>Vitex</i>	277
<i>Triticum</i>	66	<i>Vitis</i>	25
<i>Trollius</i>	125	<i>Violaria</i>	219
<i>Tropaeolum</i>	204	<i>Vlmus</i>	20
<i>Tuberosa</i>	103	<i>Vnifolium</i>	104
<i>Tulipa</i>	102	<i>Vrena</i>	157
<i>Tulipifera</i>	127	<i>Vrtica</i>	92
<i>Tuna</i>	161	<i>Vua vrfi</i>	204
<i>Tunica</i>	195	<i>Vulneraria</i>	343
<i>Turritis</i>	335		
<i>Tussilago</i>	404	X.	
<i>Typha</i>	82	<i>Xanthium</i>	13
<i>Typhoides</i>	60	<i>Xeranthemum</i>	406
		<i>Xiphium</i>	92
V.			
<i>Vaccinium</i>	200	Y.	
<i>Valantia</i>	231	<i>Yucca</i>	20
<i>Valeriana</i>	377		
<i>Valerianella</i>	378	Z.	
<i>Vella</i>	321	<i>Zadantha</i>	436
<i>Vetatrum</i>	440	<i>Zarichellia</i>	57
<i>Verbascum</i>	265	<i>Zea</i>	73
<i>Verbena</i>	318	<i>Ziziphora</i>	299
<i>Verbescina</i>	407	<i>Ziziphus</i>	16
		<i>Zygophyllum</i>	183



P R A E F A T I O.

generibus ejusmodi characteres essentialia addere animus erat, nisi moles libelli nimis incrementum obtinisset. Inter nomina specifica & synonyma plantarum tam ea selegi, quae descriptionem fere brevem, proximasque discriminis notas exprimentem, contineant, quam illa, quae in officinis aut pinacibus v. g. Bauhinianis vel Tournefortianis recepta esse videbam. Descriptiones vero uberiores specierum & observationes parcius adspersi, illasque nonnisi breves paucioribus speciebus, quae aut plane novae, aut nonnisi obscure notae minisque definitae videbantur, addidi. Animus autem est, proxime variarum plantarum, quarum historia nondum satis extricata est, aut discriminis notae minus plénæ datae esse videntur, uberiores descriptionem & fusiores Synonymiam proprio libello exhibere; uti supplemento novos cives, de quibus, etsi in horto iam adsint, quum nondum tamen floruerint, certi quidquam vix satis tuto determinari posse videtur, & plantulas illas ad cryptogamiam pertinentes, spatio hic exclusas, quamprimum feret occasio, subiungam. Paucissimas species prioris Enumerationis omisi, easque tantum, quae in regionibus Germaniae ab agro Gottingensi remotioribus

PRAEFATIO.

bus crescant, nec inter cives Florae nostrae referri possunt. Nullas, si quae forte hoc anno in horto nostro desiderantur, nota peculiari signavi, cum copia plantarum in omnibus hortis res sit mutationi & vicissitudini summe obnoxia, & satis multae, quae in priori enumeratione perisse dicuntur, hortum nostrum iterum ornent. Loca natalia forte non satis plena me dedisse nonnullis videbor. Sed mole libelli impeditus, in ipsis demonstrationibus plura imprimis circa plantas indigenas fusius addi posse credideram. Iconem addidi unicam plantae, quae tam propter novitatem quam ob singularem fabricam peculiari attentione digna videbatur.

Göttingae. d. 20. Apr. MDCCLVII.



APETA.



APETALAE.

Plantae petalis nullis vel obscurissimis, semine, flore, & veris staminibus donatae.

CLASSIS. I.

AMENTACEAE.

Flores mares & flores feminas a se invicem remotos, aut in eadem, aut in distinctis plantis, ex receptaculo, sive axi communi stirpiformi gerunt, squamis inter flores interpositis.

ORDO. I.

AMENTACEAE CONFERAEE.

Flos masc. Squamae circa axin communem in conum congestae, quae ad basin internam aetheras fovent, filamentis aut nullis aut coactis.

Flos femin. Squamae similes, ut in mares, quae ad basin internam semina fovent; in non-

A

nullis

APETALAE

nalis fructus baccam aemulatur. Omnibus præterea indoles resinosa, & angusta sempervirentia folia.

EPHEDRA. LINN. 1007.

Planta anomala, & media inter Equifeta & Coniferas. Diphyta. *Fl. Masc.* Ad squamas coni stamina septem; *Fem.* Semina duo calyce baccato tecta.

H. Ephedra pedunculis oppositis: amentis geminis. LINN. *Spec. 1. distachya.* p. 1040. *Hort. Cliff.* 465. *ROY. Prodr. Fl. Lugd. Bat.* p. 87. HALLER *Enum. Helv.* p. 145. Ephedra minima minor. I. R. H. HALL. *Enum. Gott.* p. 9. Vua marina Monspelienium. LOBEL. *icon.* p. 796. In Valleja. Gallia. Narbon. Hispania. Fr.

TAXUS LINN. 1006. HALL.

Helv. 146.

Diphyta. *Flas. masc.* Ad squamas coni antheræ peltatae. *Flas. fem.* Calyx in baccam maturatus semen vacuū fovet.

Taxus. omnium anchorum. LINN. *Cliff.* 464. ROY. *Lugd. bat.* 87. HALL. *Helv.* 146. *Enum. Gott.* p. 9. C. BAVH. *Pin.* 505. Taxus foliis approximatis. *baccata.* LINN. *Sp. 1.* p. 1040. *Taxbaum. Ebenbaum.* In sylvis circa Pless abunde. Fr.

JUNIPERUS. LINN. 1005. HALL. 146.

Diphyta. *Masc.* Ad squamas coni tres antheræ connatae. *Fem.* Baccæ trisperma, tribus tubis.

Juni-

AMENTACEAE CONIFERAE. 7

Juniperus foliis patentibus. LINN. *Cliff.* 465. ROY. *Lugd. bat.* 89. HALL. *Helv.* p. 146. *Enum. Gort.* p. 9. JUNIPERUS vulgaris & fruticosa. B. arborea. C. BAVH. *Pin.* 488. *Wachholder-Steude.* *Krautw. Deeren-Steude.* Göttingae rarius in declivibus montis *Heimberg* ad sinistram viae *Rohringens*. Fruticosa tamen in aliis regionibus Germaniae. v. g. circa *Jenam* & *Noribergam* copiose; arborea rarius reperitur. Fr.

H. *Juniperus* foliis inferne adnatis; oppositionibus concatenatis. LINN. *Cliff.* 464. ROY. *Lugd. bat.* p. 90. BOEHMER. *Fl. Lips.* p. 283. HALL. *Enum. Helv.* p. 9. *Juniperus Sabina* LINN. *Spec.* 6. BLACKW. *T.* 244. *Mas.* *Sabina* sterilis. I. B. *Femina.* *Sabina baccifera.* I. B. *Hist. plant.* I. b. 288. *Sebenbaum.* *Segelbaum.* In Helvetia. Italia. Sibiria. Fr.

H. *Juniperus* foliis inferioribus ternis, superioribus binis decurrentibus subulatis patulis acutis. *Bermudiana.* LINN. *Sp.* 5. ROY. *Lugdb.* 90. *Juniperus Bermudiana.* HERM. *Lugdb.* 345. *T.* 347. In America. Fr.

PINUS. LINN. 956. Abies. Larix.
TOWN.

Monophyta. *Masc.* Ad basin interiorum squamarum coni duo loculi antheriferi. *Fem.* Ad basin squamarum duae nuculae alatae.

Pinus. *Abies.* LINN. *Spec.* 10. *Abies* foliis solitariis apice acuminatis. LINN. *Cliff.* 449.

APETALAE. PINEA

HALL. *Helv.* 147. *Enum. Gott.* p. 9. Picea major prima s. Abies rubra BAVH. *Pin.* 493. Picea varior. auctor. Rothe Tanne. Coni penduli longiores, squamae coni fem. apice subrotundo. Folia solitaria. Vbiq̃ue in sylva Hercynica & reliquis borealibus Europae, Fr.

Pinus. *Picea*. LINN. *Spec.* 8. Abies foliis solitariis apice emarginatis. LINN. *Cliff.* 449. HALL. *Enum. Gott.* p. 9. Abies foliorum apice bifido. HALL. *Helv.* p. 148. BLACKW. *T.* 203. Abies Taxi folio, fructu sursum spectante. TOURN. 585. *Weistanne*. Folia inferne albida. In montosis Germaniae, Circa Noribergam, Fr.

Pinus. *Larix*. LINN. *Spec.* 7. Abies foliis fasciculatis obtusis. LINN. *Cliff.* 450. Larix folio deciduo. HALL. *Enum. Helv.* p. 149. Larix. C. BAVH. & reliquorum auctorum. *Lerchenbaum*. In alpinis Helvetiae, Sibiriae. In sylvis Hercynicis fata laete provenit. Fr.

Pinus. *Sylvestris*. LINN. *Spec.* 1. Pinus foliis geminis, tono pendulo, nucleis strigosis. HALL. *Enum. Helv.* 149. Pinus sylvestris C. BAVH. *Pin.* 491. *Fiechte. Kibnbaum*. Folia bina. Coni femini squamae apice quadrangulo. Vbiq̃ue in arenosis Germaniae. Fr.

H. Pinus foliis ternis. *Taeda*. LINN. *Sp.* 3. Pinus foliis longissimis ex vna theca ternis COLD. *Novch.* 230. In Virginiae, Canadae paludosa. Fr.

THVJA.

AMENTACEAE CONIFERAE.

THUJA. LINN. 975.

Monophyta. *Masc.* Ad basin squamarum amen-
ti ovati stamina quatuor. *Fl. fem.* Squamae biflo-
rae, styli subulati, semina oblonga alata.

H. Thuja strobilis laevibus : squamis obtusis.
occidentalis. LINN. *Spec.* 1. BLACKW. T. 210.
Thuja Theophrasti. CL. BAVH. *Pin.* 485. Arbor
vitae CLVS. *Hist.* 1. p. 36. *Lebensbaum.* In Po-
lonia. Canada. Sibiria. Fr.

H. Thuja strobilis squarrosis : squamis acu-
minatis reflexis. *orientalis.* LINN. *Spec.* 2. HALL.
Enum. Gott. p. 107. In China. Fr.

CYPRESSVS. LINN. 976.

Monophyta. Thuiae maxime affinis. *Fl. masc.*
Ad singulas squamas peltatas antherae quatuor.
Fl. fem. Ad basin squamarum puncta numerosa
truncata loco styliorum. Nux angulata.

H. Cupressus foliis imbricatis erectis. LINN
Cliff. 449. ROY. *Lugd. bat.* 88. Cupressus *sem-*
pervirens. LINN. *Spec.* 1. p. 1002. BLACKW.
T. 127. Cupressus meta in fastigium convolu-
ta, quae femina Plinii. I. R. H. HALL. *Enum.*
Gott. p. 7. 587. Cupressus BAVH. *Pin.* 488. &
officin. *Cypresse.* Sponte in Creta. Fr.

ORDO. II.

AMENTACEAE JVLIFERAE.

Aliae monophytae, aliae diphytae. *Fl. masc.*
Amentum, *Julus* dictum, pendulum constans

6. APETALAE

ex squamis circa axem communem congestis, quae ad basin flores, staminum filamentis distinctis, fovent. *Fl. sem. & fructus* mire variat. Omnes, fruticolae.

SALIX. LINN. 926.

Diphyta, Fl. mae. Ad singulas squamas staminum duo aut tria, *Fl. sem.* Julius ex capsulis bivalvibus.

Salix foliis ferratis glabris, acuminatis, appendiculatis. LINN. *Spec.* 349. *ROEHMER. Lips.* n. 697. p. 286. *Salix* Persicae folio auriculato. HALL. *Enum. Gott.* p. 8. *Salix* folio amygdalino utrinque aurito C. BAVH. *Pin.* 473. *Salix amygdalina* LINN. *Sp.* 6. p. 1016. Ad rivos.

Salix foliis elliptico lanceolatis, utrinque glabris, ferratis, appendiculatis. HALL. *Enum. Gott.* p. 8. *Salix* folio auriculato, splendente, flexilis. RAR. *Salix triandra*. LINN. *Sp.* 2. p. 1016. Passim circa pagos.

Salix lento vimine & squama florali flavis, foliis elliptico lanceolatis, subtus sericeis. HALL. *Gott.* p. 8. *Salix vitellina*. LINN. *Spec.* 9. BLACKW. *T.* 327. α . *Salix* sativa lutea folio crenato. C. BAVH. β . *Salix* vulgaris, alba, alborefcens. C. B. & *Enum. Gott.* l. c. Vimen flexillimum. Ubique feritur.

Salix foliis utrinque levibus, inferne glaucis, squamis & capsulis torbentosis. HALL. *Helv.* n. 9. p. 153. *Salix Helix*. LINN. *Spec.* 11. In pratis.

Salix

AMENTACEAE JVLIFERAE.

Salix foliis ovatis integerrimis, subtus tomentosis nigris. ROY. *Lugdb.* p. 84. *Salix Armaria*. LINN. 21. p. 1019. *Salix* foliis utrinque candicantibus & lanuginosis. C. B. *Pin.* 474. In paludosis arenosis, circa Berolinam.

Salix foliis ovatis sericeis, iulis crassissimis. HALL. *Gott.* p. 9. *Salix Caprea*. LINN. *Spec.* 26. p. 1020. *Salix latifolia* rotunda. C. BAVH. *Pin.* 474. HALLERVS ad hanc speciem multas alias salices vt varietates refert. Vbiq; in dumetis & sylvis.

Salix angustis & longissimis foliis, subtus albicantibus. HALL. *Gott.* p. 9. C. BAVH. *Hist.* I. P. II. p. 212. *Salix fragilis*. C. BAVH. *Prodr.* p. 159. Circa *Htperhausen*.

POPVLVS. LINN. 996.

Diphyta. *Fl. masc.* Ad singulas squamas stamin. octo. *Fl. fem.* vt in *Salice*.

Populus foliis subrotundis, dentato-angulatis, utrinque levibus. LINN. *Hort. Cliff.* 460. HALL. *Helv.* 156. *Populus tremula*. LINN. *Spec.* 2. C. BAVH. *Pin.* 429. *Espen. Zitterpappeln.* In dumetis & sylvis.

Populus foliis deltoidibus acuminatis serratis. LINN. *Cliff.* 460. HALL. *Helv.* 156. *Enon. Gott.* p. 9. *Populus nigra*. LINN. *Spec.* 3. BLACKW. T. 248. C. BAVH. 429. *Rhein-Weide. Schwarzer Pappelbaum.* Ad aquas.

Populus foliis subrotundis, dentato-angulatis, subtus tomentosis. LINN. *Cliff.* 400. HALL. *Helv.* n. 1. p. 158. *Populus alba.* LINN. *Spec.* 1. *Rappet. Weide.* *Weisser Pappelbaum.* *An der Unstrut.* In Rhem infulis. Circa Beralium.

ALNVS. HALLER. 157. *Betulae species* LINN.

Monophyta. *Fl. masc.* Ad singulas squamas tres flores tetraspermones. *Fl. fem.* Semina inter squamas apice crassiores.

Alnus rotundifolia, glutinosa viridis. c. BAYH. *Pin.* 428. HALL. *Enum. Gott.* p. 10. *Alnus auctorum:* *Betula alnus.* LINN. *Spec.* 5. *Ertm.* In paludosis & ad rivos. Fr.

BETVLA. HALLER. 158. LINN. 933.

Monophyta *Alno* Smilis. Fr. Inter squamas coni semina alata.

Betula foliis acuminatis serratis. *Cliff.* 447. HALL. *Helv.* 158. *Betula* BLACKW. T. 240. BAYH. *Pin.* 447. & auctor. *Betula alba.* LINN. *Spec.* 1. p. 982. *Birch.* Ubique in dumis sylvisque. Fr.

Betula foliis orbiculatis crenatis. nana LINN. *Fl. Lapp.* 266. T. 5. f. 4. *Amoen. Acad.* 1. p. 4. T. 1. HALL. *Helv.* 158. *Enum. Gottlag.* p. 10. *Alder. Birche.* In paludosis circa montem Bruchert. v. g. *Auf der Heinrichs-Höhe* Fr.

AMENTACEAE JVLIFERAE. 29

CARPINVS. LINN. 952.

Monophyta. *Masc.* Julius ex squamis polystemonibus. *Fem.* Inter squamas foliaceas nuxes monospermae.

Carpinus squamis strobilorum plantis. Britan. LINN. *Spec.* 1. p. 998. *Carpinus* HALL *Helv.* 158. & *austriacum. Charno.* Gall. *Maynab.* *eben. Steinh.* In sylvis. Fr.

QUERCVS. LINN. 949.

Monophyta. *Masc.* Julius ex Floribus distitis. *Fem.* Aliquot scuta plana loco tubae. *Fr.* Nux oblonga in calyce haemispherico, scabro, persistente.

Quercus foliis anhuus oblongis superne latioribus; sinibus acutioribus; angulis obtusis Cliff. 448. HALL. *Helv.* 159. *Quercus Robur.* LINN. *Spec.* 12. p. 996. *Quercus vulg.* BLACKW. T. 487. & *auctor. Eiche.* Ubique in sylvis. Fr.

JUGLANS. LINN. 950.

Monophyta. *Masc.* Ad squamas singulas stamina plurima. *Fem.* Stigmata maxima clavata, lacera. *Fr.* Sub cortice carnoso nux, in quo nucleus quadrilobus.

H. *Juglans foliolis ovalibus glabris subserratis, subaequalibus. Regia.* LINN. *Sp.* 1. p. 997. *Nux Juglans. f. regia vulgaris.* BAVH. *Pin.* 417. *Nux iuglans.* BLACKW. T. 247. *varior. auctor. Welsche Nüsse.* Colitur in hortis exoticis. Fr.

CORYLVUS, LINN. 953.

Monophyta. *Masc.* Ad squamas singulas stamina octo ad decem. *Fem.* Fructus s. nuxes conglobatae singulae in calyce lacero.

CORYLVUS stipulis ovatis obtusis. *Avelana*. LINN. *Spec.* 1. p. 998. HALL. *Helv.* 159. *Corylus sylvestris*. BLACKW. T. 293. BAVH. *Pin.* 418. & *officin.* *Häselstande, Haselnüsse.* In sylvis frequens. Fr.

FAGUS, LINN. 951. Castanea. HALL.

TOVERN.

Monoph. *Masc.* Flores campanulati quinquefidi axi communi affixi. *Fem.* Fr. Nuxes duae in capsula quadrivalvi echinata.

H. *Fagus Castanea*. LINN. *Spec.* 1. *Fagus* foliis lanceolatis acuminato serratis subtus nudis LINN. *Cliff.* 447. *Castanea* & *sylvestris* β sativa. p. BAVH. *Castanea Helv.* 169. BLACKW. T. 330. & *officinarum*. *Châtaigner* Gall. *Castanien*. Flores mares in cylindrum dispositi. Variis in locis colitur, non indigena. Fr.

FAGUS foliis ovatis obsolete serratis *Cliff.* 447. *Fagus officin.* & *omnium fere auctorum*. *Hære*. Gall. *Büchen*. *Rotb-Büchen*. Flores mares in globum digesti. Vbiq. in sylvis. Fr.

PLATANVS, LINN. 954.

Monoph. *Mares* in apertum globosum digesti. *Fem.* Fructus plures in globum dispositi.

Platanus foliis palmatis. LINN. *Cliff.* 447.
Platanus orientalis. LINN. *Spec.* 1. p. 199.
Platanus. BAVH. *Pin. & auctor.* Colitur in hor-
 tis principum. v. g. in horto *Wilhelmsbal* pro-
 pe *Cassel.* Fr.

LIQUIDAMBAR. LINN. 955.

Modoph. *Mares* polystermones in amentum,
 conicum, laxum. *Feminae* ad basim spirae masculae,
 in globum digestae.

H. *Liquidambar* foliis palmato angulatis.
Synaciflua. LINN. *Spec.* 1. p. 999. *Liquidam-*
bar. *Enum.* Gott. p. 12. LINN. *Mat. med.* 432.
 C. BAVH. *Pin.* 503. *offic. & Auctor.* Nota ar-
 bor ob resinam officinalem *Storacis liquidae.* In
 Virginia. Fr.

PISTACIA. LINN. 981. *Terebinthus.*
Lentiscus.

Diphyta. *Mares* quinquefidi, pentastemones
 in amento laxo. *Fem.* Amentum nullum. Styli
 tres. Fr. Nux ovata intra drupam.

H. *Pistacia* foliis abrupte pinnatis, folio-
 lis lanceolatis. *Lentiscus.* LINN. *Spec.* 5. *Mat. med.*
 453. *Lentiscus vulgaris.* BAVH. *Pin.* 399. & *offi-*
cingrum. *Mastixbaum.* In calidiori Europa. In
 Palaestina. Fr.

MYRICA. LINN. 981. *Gale.*

Diphyta. *Musf.* Ad singulas squamas stamina
 quatuor. *Fem.* Styli duo. Fr. bacca unilocu-
 laris.

H. *Myrica* foliis lanceolatis, fructu scoto.
 LINN. *Mat. med.* 451. HALL. *Enum. Ger.* p.
 12. *Myrica* Gale. LINN. *Spec.* 1. *Myrica* ba-
 bandica offic. In ducatu Cellensi. Fr.

H. *Myrica* foliis oblongis opposite sinuatis.
Quercifolia. LINN. *Spec.* 4. *Laurus Africana*
 minor, folio. *Quercus*. COMME. H. 2. p. 111.
 T. 81. In Aethiopia. Fr.

HIPPOPHAE. LINN. 989.

Diphyta. *Mist.* In calyce bipartito, laminae
 quatuor. *Pem.* In calyce tubuloso, bifido, styli
 simplex. Fr. bacca unilocularis.

H. *Hippophae* foliis lanceolatis. *Rhamnaoides*.
 LINN. *Sp.* 1. HALL. *Helv.* 161. *Hippophae*.
Cliff. 454. ROYEN. 207. In littoribus arenosis
 fluviorum. Fr.

CLASS. II.

NVCAMENTACEAE.

Flores masculini & feminini distincti aut in eadem
 planta, aut in eodem disco. Mares multi, tubu-
 losi, monopetali, squamis interpositis, recepta-
 culo communi affixi, antheris in singulis con-
 ventibus. Flores femine aut ad basem marium,
 aut in eodem disco cum ovaribus. Fructus in
 aliis nux, in aliis semina ex placenta communi
 plura, squamis interpositis.

NYCAMENTACEAE.

139

XANTHIUM. LINN. 937.

Flores *mares* in capitulum collecti. *Fem.* Capsulae infra-masculos congestae floscae, aculeatae, apice bifido, biloculares.

Xanthium caule inermi. *Strumarium*. LINN. *Spec.* 1. p. 987. *Mat. Med.* 424. HALL. *Helv.* 151. Xanthium. BLACKW. *T.* 444. Xanthium. Dioscoridis L. *Lappa minor* C. BAVH. *Pin.* 198. *Spitzkletten*. Circa *Salz der Heiden*. & *Allendorf*. *Berolini* in ipsa vrbe prope sterquilina. ANN.

H. Xanthium spinis ternatis. *spinosum*. LINN. *Sp.* 1. Xanthium Lusitanicum spinosum. HERM. *Parad.* *T.* 246. HALL. *Enum. Gott.* p. 11. In Lusitania. ANN.

AMBROSIA. LINN. 938.

Flores *mares* congesti, calyce communi monophyllo. *Fem.* ad basin marium, tuba bifida. *Fr.* Nax ovata, aculeata, monosperma.

H. Ambrosia foliis multifidis, racemis solitariis pilosis. *maritima*. LINN. *Sp.* 4. p. 988. Ambrosia *maritima*. BAVH. *Pin.* 138. Ad ripas maris mediterranei. A.

H. Ambrosia foliis trilobis serratis. *trifida*. LINN. *Sp.* 1. Ambrosia *Canadensis altissima*, hirsuta Platan folio. HALL. *Enum. Gott.* p. 11. vbl characterem emendatum huius plantae dedit. In Virginia. Canada. A.

IVA. LINN. 940. Tarchonanthus.

VAILL.

Flores *mares* & *fem.* in discum congesti, calyce communi. *Mares* in disco, tubulosi, staminibus quinque connatis. *Feminae* in ambitu pauciores, ex apice capsulae superne crassioris condensatae, in qua semen exiguum latet; styli duo, capillares, vagina quinquedentata inferius. *Andr.*

H. Iva foliis lanceolato ovatis, caule herbaceo. *annua*. LINN. *Sp.* 11 p. 988. *Hort. Vp.* 285. Tarchonanthos foliis cordatis serratis trinerviis. ROYEN. 538. In America meridionali. A.

H. Iva foliis lanceolatis, caule fruticoso. *frutescens*. LINN. *Spec. 2. Gen. nov.* 1106. Parthenium foliis lanceolatis serratis. *Cliff.* 443. ROYEN. 26. In Virginia, Peru. Fr.

PARTHENIVM. LINN. 930. Hysterophorus. VAILL.

Flores *feminint* & *androgyni* calyce communi excepti. *Androg.* plures in disco tubulosi. *Fem.* pauciores in radio lingulati, stylo bifido. Semina turbinato cordata.

H. Parthenium foliis composito-multifidis. *Hysterophorus*. LINN. *Sp.* 1. Partheniastrum Americanum Ambrosiae folio. NISSOL. *Act. Bot. Berol.* 1711. p. 423. T. 13. f. 2. In Jamaica. A.

MICROPVS. LINN. 892. Gnaphalodes.

TOVRN.

Flores *androgyni* & *feminini* congesti ad alas foliorum. *Androg.* steriles tubulosi, plures in disco. *Fem.* in radio nudi, quorum germina latent intra singulas calycis squamas galeatas, dorso echinatas. Omnes partes plantae disruptae fila trahunt.

H. *Micropus supinus*. LINN. *Spec.* 1. p. 1. 927. ROY. 145. *Gnaphalodes Lusitanica* TOVRN. I. R. H. 439. HALL. *Enum. Gott.* p. 12. Ad ripas maris mediterranei. A.

CLASS. III.

Sexus varius. Calyx quadri- aut quinquepartitus, plerumque interne coloratus. Corolla nulla aut petala minima. Stamina tot, quot segmenta calycis, ex calyce orta, paucioribus exceptis, ubi stamina bis tot. Tubae duae vel tres. Fructus plerumque bacca, tot seminibus foeta, quot stigmata adfunt.

ORDO I.

VEPRECVLAE LINN.

EMPETRVM LINN. 977.

Diphyt. *Masc.* Calyx trifidus. Stamina tria. *Fem.* Calyx similis, Styli novem, bacca totidem seminibus foeta.

Empe-

Empetrum procumbens nigrum. LINN. *Sp.* 2. p. 1022. HALL. *Helv.* 162. *Erica baccifera*. CLVSII & *aliorum*. In monte *Bructero*. Fr.

VISCUM. LINN. 979. HALLER.
Helv. 162.

Monoph. HALLER. *Diphyta* LINN. *Masc.* *Stamina* sessilia quatuor *Fem.* Calyx quadrifolius, bacca monosperma, glutine plenissima. Flores ad ortum foliorum conglomerati. Planta parasitica.

Viscum album. LINN. *Sp.* 1. HALL. *Helv.* 162. *Viscum baccis albis*. C. BAVH. *Pin.* 423. *Viscum* BLACKW. T. 184, & *officin.* *Mistel*. In variis arboribus, in primis in *Pyro*. Fr.

BUXUS. LINN. 934. HALLER. 163.

Monoph. Flores utriusque sexus misti in gemmis plantae. *Masc.* Calyx quadrifolius. *Fem.* Calyx similis. Capsula trirostris, trilocularis. Folia sempervirentia.

H. *Buxus sempervirens*. LINN. *Spec.* 1. HALLER. 163. BLACKW. T. 196. & *offic.* *Buxbaum*. Sponte in calidiori Europa, in horta culta & tonsa. Fr.

RHAMNUS. LINN. 235. *Frangula*. *Paliurus*. *Alaternus*. *Ziziphus*. *Cervi spina*.

TOURN.

Calyx quadri- & quinque-partitus. Ad singulas divisiones segmentorum squamulae introrsum conniventes, sub quibus stamina.

Rhamnus

Rhamnus spinosus, foliis acuminatis, serratis.
HALL. *Helv.* 163. *Enum. Gott.* p. 13. *Rhamnus*
catharticus. LINN. *Sp.* 1. *Rhamnus catharticus*
BAVH. *Pin.* 478. BLACKW. *T.* 135. Spina cer-
vina *Offic. Greutzbeer. Wegdorn.* Planta diphyta.
Calyx quadrisfidus. Stigma quadrisfidum. In ad-
fensu M. *Alengst.* circa *Plesf.* & alibi in sepi-
bus. Fr.

H. *Rhamnus inermis*, foliis glabris, crenatis.
HALL. *Comment. Soc. Sc. Gott.* p. 349. *T.* 16.
En. Gott. p. 13. *Rhamnus alpinus*. LINN. *Sp.* 2.
In horto tabernae adest femina, cuius calyx qua-
drisfidus, stigma tri- & quadrisfidum. Sponte in
alpidibus Helveticis. Fr.

Rhamnus inermis, floribus monogynis herma-
phroditis, foliis integerrimis. *Frangula*. LINN.
Spec. 3. ROY. 224. *Frangula* ora folii integra.
HALL. *Enum. Helv.* 164. *Frangula*. *Alnus nigra*.
BLACKW. *T.* 152. *Offic. Faulbaum.* Calyx quin-
quesfidus. Stigma emarginatum. Bacca disperma.
In humidis v. g. circa *Landotsbausen.* Ad sepes
super *Weende.* Fr.

H. *Rhamnus inermis* floribus dioicis, stigma-
te triplici. *Alaternus*. LINN. *Spec.* 4. *Physica*
elator. C. BAVH. *Pin.* 427. *Alaternus* CLVS. &
varior. auctor. Calyx quinquesfidus. Squamae ca-
lycis deficiunt. In calidiori Europa. v. g. Hispa-
nia. Fr.

H. *Rhamnus aculeis geminatis rectis*, floribus
digynis, foliis ovato-oblongis. *Zizyphus*. LINN.
Sp. 9. *Juiba sylvestris*. BAVH. *Pin.* 446. *Zi-*
zyphus.

zyphus. DOD. *Pempt.* 807. *Tubae Offic. Rotbe Brust-Beeren.* Calyx quinquefidus. Baccae nucleis bilocularis. In Europa Australi, Africa & Syria. Fr.

H. *Rhamnus aculeis geminatis, inferiore reflexo, floribus trigynis. Paliurus.* LINN. *Sp.* 5. *Paliurus.* TOURN. 387 & Auctor. Nucleus trilocularis, calyx quinquefidus. In Europa australi. Fr.

THESIUM. LINN. 258. HALLER. 183.

Linofyris. Linophyllum.

Calyx quinquepartitus. Tuba stigmate globosa. Semen in fundo calycis.

Thesium panicula foliata, foliis linearibus. Linophyllum. LINN. *Sp.* 1. *Thesium Cliff.* 41. HALL. *Helv.* 183. In montosis siccis Germaniae, v. g. M. *Bratfero, Weisner.* Fr.

Quae sequuntur duo genera, stamina bis tot, quot segmenta calycis habent.

ORD. II.

SCLERANTHUS. LINN. 497. KNAWE. T.

Calyx semiquinquefidus, segmentis acutis oris albis. In fundo calycis capsula disperma.

Scleranthus calycibus fructus patulis. annuus. LINN. *Sp.* 1. p. 406. KNAWE. *annuum maius, flore minori.* HALL. *En. Gert.* p. 32. *Polygonum minus polycarpon.* TABERN. p. 834. Ubique in agris. A. Scl-

Scleranthus calycibus fructus, clausis. perennis.
 LINN. Sp. 2. Knewel inqnum, flore maiore,
 perenne. HALL. Helv. 186. Polygonum Polo-
 nicum, cocciferum. 1. BAVH. III. p. 378. Cel-
 bris olim planta propter gallam, radici adhae-
 rentem, enatam ex sic dicto Progall-insecto, quae
 ad rem tinctoriam & Confectionem *Alkermes* in-
 tu fuit. In arenosis *Cellenfis*. P.

DAPHNE. LINN. 436. Thymelaea. HALL. 188.
 TOURN. 366.

Calyx petalo similis, caducus, tubulosus, lim-
 bo quadrifido. Bacca unilocularis.

Daphne floribus sessilibus infra folia elliptico-
 lanceolata. LINN. Cliss. 147. Mat. med. 179.
 Daphne Mezereum. LINN. Spec. 1. 2. 356.
 Thymelaea spica foliis levibus, mollibus, lanceo-
 latis subiecta. Enum. Helv. 166. Enum. Gott. p.
 166. Laureola folio deciduo flore purpureo of-
 ficinis laureola femina. BAVH. Pin. 462. Kell-
 bals. Vbique in sylvis primo vere. Fr.

Daphne Laureola. LINN. Sp. 5. Thymelaea
 foliis duris levibusque, floribus intermissis HALL.
 Enum. Gott. p. 166. Laureola semper virans, flo-
 re viridi, quibusdam Laureola mas. 0. BAVH. Pin.
 462. BLACKW. T. 62. In Helvetiae montosis
 circa Heidelberg. Fr.

CHRYSOPLIVM. LINN. 483.

Calyx quadri- & quinquepartitus, coloratus,
 patulus. Stamina ex receptaculo orta. Capsu-
 la unilocularis, birostris, calyce cincta.

1. *Chrysofenitum* foliis alternis. LINN. *Spec.* 1. HALL. *Helv.* 189. Chryf. foliis pedunculis oblongis insidentibus. I. R. H. 146. In vdis opacis. Flores racemosi. A.

ORD. III.

Prioribus omnino adsimiles. Styli autem duo ex fructu sicco monospermo.

VLMVS. LINN. 281.

Stamina calyce colorato longiora. Fructus latus. Flosculi ex eadem gemma plures.

1. *Vlmus* fructu membranaceo. LINN. *Cliff.* 83. HALL. *Helv.* 167. *Vlmus campestris*. LINN. *Sp.* 1. p. 225. *Vlmus campestris* & Theophrasti. BAVH. *Pin.* 246. *Vlmenbaum*. In sylvis & ad pagos. Fr.

BOSEA. LINN. 280.

Calycis segmenta lanceolata. Bacca globosa. Flores racemosi.

H. Bofea. *Yervamora*. *Spec. pl.* 1. p. 225. ROY. *Lugdb.* 223. Bofea arbor. HALL. *Gott. p.* 14. Frutex peregrinus horto Bosiano Yervamora dictus. WALTH. *Hort.* 24. T. 10. In insulis Canariis. Fr.

CELTIS. LINN. 1012. LVDW. 782.

Calycis segmenta patula. Flores *androg.* & *masc.* Stamina brevia. Styli pubescentes. Fr. rotundus nucis foetus. Flores in alis foliorum solitarii.

H. *Celtis* foliis ovato-lanceolatis: *australis*. LINN. Sp. 1. *Celtis* fructu nigricante. HALL. Gott. p. 14. *Lotus* arbor. LÖB. ic. 186. In Europa australi. Fr.

ORD. IV.

Plantae variae arboreae, prioribus multum adsimiles, vario tamen modo anomalae.

MORVS. LINN. 936.

Pl. Monophyta. Flores fere amentacei. *Mat.* Stamina calyce longiora quatuor. *Fem.* Bacculae multae pulposae, baccam compositam faciunt.

H. *Morus* foliis oblique cordatis laevibus. *alba*. LINN. Spec. 1. *Morus* fructu albo. BAVH. Pin. 459. *Morus alba*. HALL. Gott. p. 14. *Weißer Maulbeerbaum*. Celebris propter Bombycem, quae foliis illius pascitur. Sponte in China, nunc ubique fere in Europa colitur, coeli nostri satis patiens. Fr.

H. *Morus* foliis cordatis scabris. *nigra*. LINN. Sp. 2. *Mat. med.* 422. *Morus nigra*. BLACKW. T. 126. BAVH. Pin. 459. *Schwartzer Maulbeerbaum*. Sponte in Italia. Nobis in hortis colitur. Fr.

LAVRVS. LINN. 452. LVDW. 133.

Stamina novem, quorum tria inferiora ad basia glandulifera. Ex tuberculis filamenta circa germen. Tuba simplex. Sub drupa nux. Sexus & corolla variat. Flores racemosi. Fr.

H. *Laurus* foliis lanceolatis venosis perennantibus, floribus quadrifidis diopis. *nobilis*. LINN. Sp.

4. *Mat. med.* 194. *Laurus vulgaris*. C.B. *Pin.* 460.
HALL. Gott. 15. *BLACKW. T.* 175. *Lorbeerbaum*.
 In Italia Graecia. Fr.

H. *Laurus* foliis lanceolatis perennantibus veno-
 sis planis, ramulis tuberculatis cicatricibus, floribus
 racemosis. *Indica*. LINN. *Sp.* 5. *Laurus Indica*. PLUK.
Bot. Zeyl. ubi vide characterem. In Virginia. Fr.

FICUS. LINN. 1032.

In interna superficie calycis communis carnosus
 osculi mares & feminae. *Mas.* Calyx tripartitus,
 stamina tria. *Fem.* Calyx quinquepartitus. Tuba
 bifida.

H. *Ficus* foliis palmatis. *Cavica*. LINN. *Spec.* 1.
 1059. *Mat. med.* 478. *Ficus communis*. BAYH.
Pin. 457. *BLACKW. T.* 125. *Feigenbaum*. In Eu-
 ropa australi. Asia. Fr.

H. *Ficus* foliis ovatis integerrimis obtusis, caule
 inferne radicato. *Benghalensis*. LINN. *Sp.* 4. *Ficus*
Bengalensis. &c. *H. Gott.* p. 14. In India. Fr.

CLASSIS IV.

SARMENTACEAE. LINN.

Omnes isostemonae, stamina quatuor ad sex, tu-
 ba una ad tres. Fructus ficcus aut bacca. Caules
 plerumque volubiles aut sarmentosi. Flores ra-
 cemosi.

CANNABIS. LINN. 988.

Diphyta. *Mas.* calyx quinquepartitus. *Fem.* Ca-
 lyx monophyllus altero latere trias. Stylis duo.
 Nux bivalvis.

H.

H. *Cannabis foliis digitatis.* LINN. *Mat. med.*
 457. *sativa.* LINN. *Sp.* 1. *Cannabis mas & femi-*
na. Auct. HALL. *Gott.* 14. BLACKW. *T.* 223. a. b.
 Hanf. Sponte in India: nobis in agris colitur. A.

HVMULVS. LINN. 989. *Lupulus.*
 HALL. 166.

Pl. *Diphyca. Mares pentastemonones racemosi.*
Flores fem. in amentis. Ad squamas maiores syn-
gulas styli duo. Semen vnicum. Caulis volubilis.

Humulus. LINN. *Sp.* 1. p. 1028. *Mat. med.*
 458. *Lupulus.* HALL. *Helv. & offic. Hopfen.* Ad
 sepes. P.

VITIS. LINN. 250. HALL. 166.

Petala quinque emarcida; minima. Tuba obtu-
sa. Bacca unilocularis. Flores racemosi. Caulis ca-
pucolis scandens.

H. *Vitis foliis palmato angulatis.* LINN. *Mat. med.*
 97. *vinifera.* LINN. *Sp.* 1. HALL. *Helv.* 166.
 BLACKW. *T.* 153. *Weinstock.* Sponte in Europa
 calidiori. ad sepes. Fr.

HEDERA. LINN. 249. HALL. 165.

Petala quinque oblonga. Stylus simplex. Bacca
pentasperma. Flosculi umbellati.

Hedera foliis ovatis & lobatis. *Helix.* LINN. *Sp.*
 1. *Mat. med.* 98. HALL. *Helv.* 164. BLACKW. 188.
 a. sterilis, β. arborea. BAVH. *Pin.* 305. *Ephru.*
Wintergrün. Cirrhis arboribus & muris adhaeret, &

scandit parasitica, Prior g. vbiq; ad arbores. β .
in calidioribus floret, sed & circa *Ples.* Fr.

H. *Hedera* foliis quinq; ovatis serratis, *quinquefolia*, LINN. *Sp.* 2. *Vitis* quinq; folia Canadensis, CORNYT. *Canad.* T. 100. HALL. *Gott.* 170. In Canada. Fr.

Menispermum, LINN. 413.

Calyx sexfolius. Petala sex. Styli tres, Baccae tres monospermae. Caulis volubilis.

H. *Menispermum* foliis peltatis subrotundis angulatis, *Canadense*. LINN. *Sp.* 1. *Menispermum* Canadense scandens, umbilicato folio. TOWN. *Ad.* Par. 1705. p. 311. In Virginia. Canada. P.

Smilax, LINN. 992.

Diphyta. Calyx sexfolius in utroque sexu. *Fem.* Styli tres. Bacca trilocularis. Planta capreolis scandens.

H. *Smilax* caule aculeato angulato, foliis dentato aculeatis cordatis, *aspera*. LINN. *Sp.* 1. *Smilax* aspera fructu rubente. BAVH. *Pin.* 296. HALL. *Gott.* 78. In Hispania. Italia. Fr.

H. *Smilax* caule aculeato angulato, foliis inermibus sagittatis, *excelsa*. LINN. *Sp.* 2. *Smilax* orientalis, aculeatis fargmentis, foliis non spinosis. TOWN. *Cor.* 45. HALL. *Gott.* 79. In Oriente. Syria. Fr.

Ruscus, LINN. 1005.

Pl. Diphyta. Calyx sexfolius, & nectarium ovatum, monophyllum, in utroque sexu. *Mar.* Stamina tria.

SARMENTACEAE.

24

tria. Fem. Seylus vnicus, Bacca globosa: Gaulia geniculatus.

H. *Ruscus foliis supra floriferis nudis, aculeatus.* LINN. *Sp.* 1. *Mat. med.* 768. *Bruscus* vel *Ruscus.* BLACKW. *T.* 155. *Ruscus myrtifolius aculeatus.* I. R. H. 79. HALL, *Gott.* 78. In Anglia. Gallia. Italia. Fr.

H. *Ruscus foliis subtus floriferis sub foliolo. Hypoglossum* LINN. *Sp.* 3. *Laurus Alexandrina* fructu pedunculo infidente. BAVH. *Pm.* 304. *Hypoglossum Bilingua.* BLACKW. *T.* 128. In Italia. Hispania. Fr.

H. *Ruscus racemo terminali hermaphrodico, racemosus.* LINN. *Sp.* 5. *Laurus Alexandrina* ramosa, fructu e summitate caulium prodeunte. HERMANN, *Lugd. bat.* 379. *T.* 681. P.

CLASSIS. V.

OLERACEAE. LINN.

Calyx quadri- aut quinquepartitus, segmentis concavis. *Petala* nulla. *Sexus* variat. *Stamina* tot, quot partes calycis, ex thalamo orta, in plerisque elastica. *Styli* duo vel tres, in paucioribus quatuor. *Ovarium* calyce plerumque persistente compressum, siccum, monospermum. *Pleraeque* herbae. *Flores* in plurimis in arctissimos glomeres, ad alas foliorum, aut in racemos dispositos, congesti.

BETA. LINN. 274.

Calyx quinquefolius. *Capsula* semine reniformi. *Flores* in alis foliorum.

H. *Beta vulgaris*. LINN. *Sp.* 1. *Mat. med.* 113. BLACKW. *T.* 235. α . *Beta sylvestris maritima*. C.B. 118. β . varietates ex colore caulis & radice. *Man-gold. Rorhe Rüben*. In Angliæ & Belgii littoribus marinis. A.

SPINACHIA. LINN. 986.

Diphyta. *Mas.* Calyx quinquefidus. *Fem.* Calyx quadrifidus, laciniis duabus minimis. Siliqua quatuor. Flores in alis.

H. *Spinachia oleracea*. LINN. *Sp.* 1. *Mat. med.* 556. BLACKW. *T.* 49. α . femine aculeato. β . semine non aculeato. Num varietates? *Spinach*. A.

H. *Spinachila foliis angulosis*, caule ramosissimo. HALL. *Gott. p.* 22. Seminis capsula foliolo terminata. A.

ATRIPLEX. LINN. 1011. HALL. 173.

Flores in eadem planta androgyni, Omares & feminae. *Andrag.* Calyx quinquefolius. Stamina glauca. Ovarium compressum. *Masc.* Calyx & staminum similia. *Fem.* Calyx maior bivalvis, cuius valvulae compressae, magnae, semen androgyni, simile includunt.

H. *Atriplex caule erecto herbaceo*, foliis triangularibus. *hortensis*. LINN. *Sp.* 4. *Mat. med.* 473. HALL. *Gott. p.* 18. α . *Atriplex hortensis alba*. β . *Atriplex hortensis rubra*. BAVH. *Pin.* 119. BLACKW. *T.* 99. *Melte*. Sponte in Tataria. A.

Atriplex

Atriplex foliis incanis, serratis, capsulis quadrangulicibus, dentatis. HALL. *Gott.* 19. *Atriplex sylvestris*, fructu roseo, compresso. C. B. *Prodr.* 58. In Thuringia. A.

Atriplex foliis triangularis, rariter dentatis. HALL. *Gott.* 1. c. *Atriplex hastata.* LINN. *Sp.* 6. *Atriplex sylvestris* folio hastato, f. deltoide. BAYH. *hisor.* 192. Circa loca salia *Harste*, & alibi in vicina. A.

Atriplex foliis imis sagittatis, superioribus lingulatis, valvis seminis dentatis. HALL. *Gott.* 19. *Atriplex patula.* LINN. *Sp.* 7. *Atriplex angustifolia* folio oblongo. BAYH. *Pin.* 219. In hortis sponte, ad fossas & vias. A.

Atriplex fruticosa, foliis ovato-triangularibus, incanis. HALL. *Gott.* 1. c. *Atriplex Halimus.* LINN. *Sp.* 1. *Halimus latifolius* f. fruticosus, BAYH. *Pin.* 120. Ad sepes maritimas Hispaniae, Lusitaniae & Virginiae. Fr.

Atriplex fruticosa, foliis lingulatis incanis. HALL. *Gott.* p. 19. *Atriplex Portulacoides.* LINN. *Sp.* 2. *Portulaca marina.* DOD. *Pempt.* 771. Calyx feminae quinquefolius. Ad littora marina Sueciae, & ad mare mediterraneum. Fr.

CHENOPODIUM. LINN. 273. HALL. 175.

Flores omnes androgyni, androgyni prioris generis simillimi. Stylus bipartitus. Semen orbiculatum.

Chenopodium foliis sinuatis, subhirsutis. HALL. *Gott.* 20. *Chenopodium foliis oblongis sinuatis, racemis nudis multifidis.* Boiss. LINN. *Sp.*

8. *Mat. med.* 109. Botrys. *BLANCH.* 314. In Europae australis arenosis. A.

H. *Chenopodium* foliis subhirsutis, oblongis dentatis, incanis. *HALL. Gott.* 20. *Chenopodium* foliis lanceolatis dentatis, racemis foliaceis simplicibus. *Ambrosioides* LINN. *Sp.* 9. Botrys ambrosioides Mexicana. *BAVH. Pin.* 138. In Lusitania, Mexico. A.

H. *Chenopodium* foliis sinuatis, subtus incanis. *HALL. l. c.* *Chenopodium glaucum*. LINN. *Sp.* 12. *Chenopodium angustifolium laciniatum minus*. *TOURN. I. R. H.* 506. Ad Europae fimeta & aquas salvas. A.

Chenopodium foliis levibus, difformiter dentatis, calycibus rubris. *HALL. l. c.* 4. *Chenopodium rubrum*. LINN. *Sp.* 3. *Chenopodium n.* 3. *HALL. Helv.* 174. *Atriplex sylvestris latifolia*. *BAVH. Pin.* 119. & *offic.* *Pes anserinus*. *TABERN. ic.* 427. *Gänse-Fuß. wilde Mette*. Inter segetes & locis ruderalis. A.

Chenopodium foliis levibus, viridibus, septangulis. *HALL. Gott.* 20. *Chenopodium hybridum*. LINN. *Sp.* 7. *Chenopodium stramonii folio*. *VAILL. Par.* 36. *T. 7. f. 3.* *Neun-Spitzen. Sautodt.* In locis cultis. A.

Chenopodium foliis subtus farinosis, rhomboideis, dentatis. *HALL. l. c. Helv.* 175. *Chenopodium viride*. LINN. *Sp.* 6. *Chenopodium Opuli folio* *VAILL. T. 7. f. 1. a.* *Chenopodium* foliis inferioribus ovatis, acutis, summis lineari-lanceolatis.

NALL.

HALL. l. c. *Chenopodium folio sinuato, candicante*,
I. R. H. Locis similibus. A.

Chenopodium foliis triangulari-sagittatis integerrimis. Bonus Henricus. LINN. *Spec. 1. Mat. med.* 106. HALL. *Helv.* 174. *Lapathum vinctuosum.* BAVH. *Pin.* 115. *Bonus Henricus offic. Guter Heinrich.* Ad pagos & ruderosa. A.

Chenopodium ex alis racemosum, foliis glabris, rhomboideis, serratis. HALL. *Gott. p.* 21. *Chenopodium Atriplicis sativae facie, spicis surrectis racemosis.* RVFF. *Fl. Jen. Hall. p.* 338. Ad lacum falsum, aliisque locis Saxoniae. A.

Chenopodium foetidum, foliis farinosis, glaucis, obtuse rhomboideis. HALL. l. c. *Helv.* 175. n. 6. *Chenopodium. Vulvaria.* LINN. *Spec. 13. Mat. med.* 108. *Atriplex olida.* BLACKW. *T.* 100. *Atriplex foetida.* C. B. *Pin.* 116. *Vulvaria* TABERN. ic. 428. *Stinckende Melde.* In horto sponte, in cultis & ruderosis. A.

Chenopodium foliis ovato-lanceolatis integerrimis. HALL. l. c. *Helv.* n. 7. p. 176. *Chenopodium polyspermum.* LINN. *Sp.* 14. *Chenopodium Betae folio.* I. R. H. 506. In agris & hortis sponte. A.

H. *Chenopodium foliis lineari-lanceolatis, ramis recurvis, dichotome ramosis, setis terminatis.* HALL. *Comm. Soc. Sc. Gott.* 1752. p. 347. *T.* 15. f. 3. *Chenopodium aristatum.* LINN. *Sp.* 21. In Sibiria. A.

H. *Cheno-*

h. *Chenopodium foliis tubulatis* superne planis, subtus convexis. LINN. *Gliff.* 86. HALL. *Gott.* p. 28. *Chen. maritimum*. LINN. *Sp.* 16. Kali minus album, semine splendente. BAVH. *Pin.* 289. Ad marina littora Europae. A.

h. *Chenopodium foliis linearibus*, hirsutis, ascendentibus. HALL. *A. c. Chenopodium. Scoparia*. LINN. *Spec.* 15. *Chenopodium Lini folio* I. R. H. *Belvedere* hortulanorum. In Graecia. China. Japonia. A.

LERCHIA. HALL. *Comm. Soc. Sc. Gott.* 1754. p. 223.

Priori generi valde simile genus. Sed tubae tres ad quinque. Fructus oblongus, sursum gracilior, in cuius fundo semen lenticulara. Flores in summitatibus inter folia.

h. *Lerchia foliis obtusis*. HALL. *Gott.* 21. *Comm. Gott. T. I. p. 223*. *Chenopodium frutescens*. LINN. *Sp.* 17. In Persia. Ad mare Britannicum. Fr.

h. *Lerchia foliis filiformibus* jacutis. HALL. *Gott.* 22. *Comm. Gott. l. 6. p. 224*. *Chenop. altissimum*. LINN. *Sp.* 18. Ad *Astracum*. Fr.

PARIETARIA. LINN. 1020. HALL. 177.

In eadem planta flores androgyni, mares & feminae. *Androg.* Calyx profunde quadrifidus, stamina elastica. Tuba plumosa. *Mas.* Stamina maiora. *Fem.* Calyx & fructus, vt in androgyno. Glomeruli florum densissimi.

Parie-

Parietaria foliis ellipticis. HALL. *Gott.* p. 17. *Parietaria officinalis.* LINN. *Sp.* 1. p. 1052. *Mat. med.* 472. *Parietaria* BLACKW. *T.* 156. *Parietaria officinarum & Dioscoridis.* C. B. *Pin.* 121. *Gloss. Kraut.* Circa Noriberm & alibi in rudementis. A.

Parietaria foliis ovatis. HALL. *Gott.* l. c. *Parietaria Lusitanica.* LINN. *Spec.* 2. *Parietaria ocimifolio, l. minor.* HALL. *Fl. Jen. Rupp.* 340. In muris Germaniae inferioris. Olim *Göttingae* in area horti. Nunc evanuit.

11. VRTICA. LINN. 935. HALL. p. 177.

Monophyta praeter unicum speciem. *Mat.* *Parietariae mari similis.* *Fem.* Calyx bivalvis; tuba plumosa, semen ovatum. Flores in racemis.

12. *Vrtica foliis oppositis, amentis fructiferis globosis. pilulifera.* LINN. *Sp.* 1. p. 983. *Mat. med.* 420. *Vrtica pilulifera, foliis cordatis, circumferratis.* HALL. *Gott.* p. 291. *Vrtica Romana.* BLACKW. *T.* 321. *Vrtica pilulifera, foliis ovatis integris.* *Gott.* l. c. Num distinctae species? In Europa australi. A.

Vrtica foliis ovatis, amentis cylindraceis. HALL. *Helv.* 178. *Vrtica vrens.* LINN. *Sp.* 3. *Vrtica vrens minor.* BAYH. *Pin.* 232. *Heiter-Nessel.* Vbi-que ad sepes. A.

Vrtica foliis oblongo-cordatis. LINN. *Mat. med.* 419. HALL. *Helv.* 177. *Vrtica dioica.* LINN. *Spec.* 4. *Vrtica vrens maxima.* BAYH. *Pin.* 232. BLACKW. *T.* 12. *Brennnessel.* Vbi-que in rudementis. A.

H. *Vertica* foliis oppositis, tripartitis, incis. *tan-nobina*. LINN. *Sp.* 5. *Vertica* foliis profunde laciniatis, semine lini. AMMANN. *Ruth.* 249. T. 25. HALL. *Gott.* 27. In Sibiria. P.

H. *Vertica* foliis alternis orbiculato vtrinque acutis, subtus tomentosis. *nivea*. LINN. *Sp.* 12. *Vertica* racemifera maxima Sinarum, foliis subtus lanuginosis. HALL. *Gott.* p. 27. In China. P.

AMARANTHVS. LINN. 941.

Glomeres florum in racemos dispositi. Pl. Monophyta. *Mas.* Calyx persistens plerumque coloratus tri- aut quinquefolius, foliis lanceolatis. Stamina tria aut quinque. *Fem.* Calyx, ut in mare. Stylis tres. Capsula ovata, monosperma.

H. *Amaranthus* capitulis subrotundis, lateralibus, foliis lanceolatis acuminatis. LINN. *Cliff.* 444. HALL. *Gott.* p. 23. *Amaranthus tricolor*. LINN. *Sp.* 1. BLACKW. T. 319. In India. A.

H. *Amaranthus* erectus, foliis emarginatis. HALL. T. 6. *Amaranthus hybridus*. LINN. *Sp.* 4. *Blitum* pulchrum, magnum, rectum, rubrum. F. B. *Hist.* 2. p. 966. In Virginia. A.

H. *Amaranthus* foliis ellipticis, racemis in aliis sessilibus. HALL. *Gott.* p. 24. *Amaranthus* floribus conglomeratis axillaribus, foliis lanceolatis obtusis. *Gracchanus*. LINN. *Sp.* 5. Racemi virides. In Virginia. A.

H. *Amaranthus* procumbens foliis ovatis emarginatis. LINN. *Cliff.* 444. HALL. *Helv.* 176. *Amaranthus*

ranthus floribus glomeratis lateralibus, foliis ovatis retusis, caule diffuso. *Blitum*. LINN. *Spec.* 6. Glomeres virides. In Europa temperiori. A.

H. *Amaranthus racemis triandris cylindricis, interruptis, axillaribus, terminalibusque, foliis rhombeis obtusis, petiolo longioribus.* An *Amaranthus spicis triandris, glomeratis sessilibus, axillaribus, terminalibusque, foliis rhombeis obtusis?* LINN. *Cent. I. Plant. n. 95.* Videtur, etsi in nostro neque spicae aristatae, neque folia latiora quam longa sint. A.

H. *Amaranthus racemis cylindricis, pendulis, longissimis.* LINN. *Cliff.* 443. HALL. *Gott.* 23. *Amaranthus caudatus.* LINN. *Spec.* 7. *Amaranthus maximus.* C. BAVH. *Pin.* 120. α . viridis. β . maculatus. γ . ruber. &c. In Peru. Persia. Zeylona. A.

H. *Amaranthus spicis cylindricis lateralibus conjunctis* LINN. *Cliff.* HALL. *Gott.* 24. *Amaranthus hybridus.* LINN. *Spec.* 8. *Amaranthus sylvestris maximus novae Angliae, spicis viridibus.* RAI *Hist.* 101. In Virginia. A.

H. *Amaranthus racemis cylindricis erectis caule inermi.* *Hypochondriacus.* LINN. *Sp.* 9. *Amaranthus sylvestris maximus novae Angliae, spicis purpureis.* TOURN. I. R. H. 238. *Amaranthus latifolius, spicis purpureis tenuibus.* HALL. *Gott.* 24. In Virginia. A.

H. *Amaranthus racemis cylindricis erectis, axillis spinosis.* *spinosus.* LINN. *Sp.* 11. ROYEN. 419. *Amaranthus indicus spinosus, spica herbacea.* HERM. *Lugd. bat.* 31. T. 33. In India. A.

CELOSIA. HALL. Gott. 25.

LINN. 255.

Androgyna. Calyx Amarantho similis quinquefolius cum squamis ad basin accessoriis tribus. Stamina connata in anulum. Stylus simplex. Fructus unilocularis.

H. Celosia foliis lanceolatis, pedunculis angulatis, spica ovato-oblonga. *argentea*. LINN. *Spec.* 1. Amaranthus spica albescente habitiori. MART. *Cent.* 7. *T.* 7. Flores magni, iuniores rubri, adultiores argentei. In America. A.

H. Celosia foliis lanceolato-ovatis. LINN. *Cliff.* 43. HALL. *Gott.* 25. Celosia *cristata*. LINN. *Sp.* 2. Amaranthus α . spicatus. β . cristatus. Flores rubri & flavi. In India. A.

H. Celosia floribus rigidis retroveris. HALL. *Gott.* 25. Achyranthes caule erecto, calycibus reflexis spicae adpressis. *aspera*. LINN. *Sp.* 1. Amaranthus Siculus spicatus. BOCC. *Sic.* 16. *T.* 9. Flores maturi contra caulem reflexi. Circa germen valvulae quinque plumosae. In Sicilia. Ceylona. Jamaica. Fr.

GOMPHRENA. LINN. 279.

Calyx Celosiae similis. In medio rubus cylindraceus ore quinquesido, totidem staminibus clauso. Styli duo. Capsula unilocularis.

H. Gomphrena caule erecto, foliis ovato-lanceolatis, capitulis solitariis, pedunculis diphyllis. *globosa*. LINN. *Sp.* 1. HALL. *Gott.* 348. α . capitulis rubris, β . capitulis argenteis. Amaranthus globosus vulgo. In India. A.

BLITVM. LINN. 14. Chenopodiumorus T.

Habitus sere Chenopodii. Calyx trifolius. Stamen unicum. Calyces conglomerati succulenti maturi baccam referunt.

H. Blitum capitellis spicatis lateralibus. *capitatum. LINN. Sp. 1.* Chenopodiumorus foliis rari-ter serratis. HALL. Gott. p. 25. Huc quoque pertinet Blitum Helv. 176. *Erdbeeren-Spinat.* In Rhæ-tia & Tyrolensi Comitatu. Anglia. A.

H. Blitum capitellis sparsis lateralibus. *virgatum. LINN. Sp. 1.* Chenopodiumorus foliis argute-
dentatis. HALL. Gott. l. c. In Tataria. A.

AXYRIS. HALL. Comment. Soc. Sc. Gott. T. II. p. 345. LINN. Gen. 429.

Monophyta. *Maris* dense spicati terminales, ob-
scuri singuli, stamine unico. *Fem.* alares. Calyx ex-
terior bi- aut trifolius, interior monophyllus, quadrifi-
dus, lacer, plumosus, tubae duae, semen com-
pressum.

H. Axyris herbacea amens masculis corymbosis
pedunculatis. HALL. Gott. p. 26. & *Gemm. Gott.*
1752. T. II. p. 345. Tab. XV. f. 2. An Axyris
caule erecto spicis simplicibus. *Amaranthoides LINN.*
Spec. 2? Folia ovato-lanceolata punctata. A.

H. Axyris foliis ovatis, caule erecto, spicis con-
glomeratis. *hybrida. LINN. Spec. 3?* Caulis ra-
mosus, lanatus in primis in summitate ramulorum.
Folia ovata, inferne incana, villosa, superne pun-
ctata; spica marium pedunculata, conglomerata,
glomerulis densissimis. A.

CAMPHORATA. HALL. 183. Camphorosma.

LINN. nov. gen. 1042.

H. Camphorata foliis lanceolatis, hirsutis, ciliatis, glomerulis florum laxè spicatis. Camphorosma. *Montpeliaca*. LINN. *Spec.* 1. *Mat. med.* 43. An Camphorata hirsuta. C. BAVH. *Pin.* 486. Glomeruli florum ad alas foliorum; cuilibet flosculo præter foliolum a latere positum maius, calyx quadrifolius, foliis duobus maioribus, glumarum instar carinatis, duobus minoribus, planis, stamina quatuor calyce minora, styli duo calice multo longiora, capsula monosperma, semen lenticulare. Vide quam diversa sit fabrica a caractere, Linnaeano. In Gallia Narbonensi. Tataria. P.

CORISPERMUM. LINN. 12.

Flores solitarii alares. Calyx bifolius. Stamen unum, duo, tria. Tubae duae. Semen compressum, marginatum.

H. Corispermum floribus lateralibus. *hyssopifolium*. LINN. *Spec.* 1. Corispermum floribus alternis. *Cliff.* 3. ROY. 205. Corispermum hyssopifolium. JUSSIEV. *Ag. Par.* 1712. p. 244. HALL. *Gott.* 26. In Tataria. Montpelii. A.

CERATOCARPVS. LINN. 921.

Monoph. Flores alares. *Mar.* Calyx bipartitus. stamen unicum. *Fem.* Calyx bifolius. Semen compressum bicornè.

H. Ceratocarpus arenarius. LINN. *Spec.* 1. p. 969. BUXBAVM. *Att. Petropol.* 1. p. 241. Ceratocarpum GERBERI. In arenosis Tatariae. A.

OLERACEAE.

37

SALSOLA. LINN. 275. Kali.

TOURN. 128.

Calyx quinquefolius. Sepalus bipartitus. Capsula globosa. Semen cochleatum. Flores in alas congesti.

H. Salsola (herbacea) foliis pungentibus. LINN.

Cliff. 86. HALL. Gott. 28. Salsola Kali. LINN.

Sp. 1. Kali spinosum cochleatum. BAVH. Pin. 289.

In littoribus marinis. A. n. 2. zylis

CLASSIS VI.

VAGINALES.

Caulis geniculatus, vaginis per intervalla cinctus. Calyx quinque aut sexpartitus, plerumque coloratus. Stamina quinque ad octo. Tubae duae vel tres. Fructus calyce comprehensus, monospermus, semen triquetrum aut ovato compressum. Flores in plerisque spicati, aut racemosi terminales in paucioribus ad alas foliorum congesti.

RHEUM. LINN. 478. Rhabarbarum.

TOURN. 118.

Calyx sexpartitus. Stamina novem. Tubae tres plumosae. Semen triquetrum, margine membranaceum. Flores paniculati.

H. Rheum foliis glabris cordiformibus. HALL.

Gott. 15. Rheum Rhabarbarum. LINN. Sp. 1.

Mat. med. 198. Rhabarbarum. ALPIN. Exot. 5.

137. BLACKW. 262. & Offic. in Thracia. Tartaria.

H. *Rheum* foliis acutis crispis subulnatis. HALL.
Gott. 15. Rheum Rhabarbarum. LINN. Spec. 2. Mat.
197. Rhabarbarum verum Offic. HELMII pro-
pe murum & in Sibiria. P.

LAPATHUM. HALL. 168. Rumex.
LINN. 407.

Calyx sexfolius. Stamina sex. Tubus tres plum-
 bae. Semen triquetrum. Sexus variat.

Lapathum acetosum textu distinctum, foliorum
 sagittatorum laminae longis & brevibus. HALL.

Helv. 168. Rumex floribus dioicis, foliis lanceo-
lato-basis. Acetosella. LINN. Sp. 22. Acetosa

avenensis lanceolata. BAVH. Pin. 114. In agris la-
pidosis rupibus muris. P.

Lapathum acetosum textu distinctum, foliorum
 sagittatorum laminae brevibus. HALL.

Rumex floribus dioicis, foliis oblongis sagittatis.
Acetosa. LINN. Sp. 21. Acetosa pratensis. BAVH.

Pin. 114. BLAC. 250. Acetosa montana ma-
xima. BAVH. Pin. 811. 4. Acetosa montana ari-

lato rotundo folio. BAVH. Pin. 811. 4. Acetosa
montana in montibus altioribus. P.

H. *Lapathum* floribus superioribus masculis, in-
 ferioribus femininis, foliis cordatis. *Lapathum ob-*

rusifolium, spica confertissima. HALL. Helv. 6. 179.
Rumex alpinus. LINN. Sp. 23. Lapathum rotundi-

folium. CAUS. Hist. 206. In alpinis. P.

Lapathum (Rumex) floribus monoicis, Glyceribus
fructus vacinatis. LINN. Cliff. 139. ROY. 230.

Rumex.

Rumex spinosus LINN. *Spec.* 10. *Beta cretica*,
 Enine aculeato. BAYH. *Pin.* 118. *Pradr.* 57. Mares
 superiores longe spicati, feminae flores alares, con-
 gesti. In Grece. P.

H. *Lapathum* (*Rumex*) valvulis laevibus, caule ar-
 boreo, foliis subcordatis. *Linnaeus* LINN. *Spec.* 15.
Lapathum acetosum calycibus triphyllis. HALL. *Gott.*
 16. *Acetosa rubra* fœns, subrotundo folio. BLACKW.
T. 252. f. 3. In Canariis. Fr.

H. *Lapathum acetosum*, valvis exterioribus
 amplissimis streptentibus. HALL. *l. c.* *Rumex vesica-*
rius LINN. *Sp.* 16. In Africa. A.

H. *Lapathum acetosum* pedunculis incrassatis
 bicornibus. HALL. *l. c.* *Rumex* fructibus dentatis
 calyce reflexis, caput bovinum referentibus. LINN.
Clas. 139. bucephalophoros. *Sp.* 13. *Acetosa ocymi*
 folio, bucephalophoros. COL. *Ecpbr.* T. 150. In Italia
 & Germania meridionali. A.

Lapathum foliis oblongis, paniculis aphyllis. HALL.
Gott. 17. a. *Lapathum* folio acuto, crispo. BAYH.
Pin. 115. b. *Lapathum aquaticum* folio cubitali.
Rumex LINN. *Sp.* 14. C. B. *Pin.* 116. γ. *Lapathum*
 acutum f. *Oxylapathum*. *ITKVF.* *Hist.* 2. p. 983.
Rumex. LINN. *Sp.* 19. HALLERVS omnes hasce plan-
 tas tanquam varietates ad vnam speciem refert. Ha-
 bitat α. β. in paludosis. γ. in siccioribus. P.

H. *Lapathum*, (*Rumex*) valvulis integerrimis
 unica granifera, foliis cordato-lanceolatis. *Sangui-*
neum. LINN. *Sp.* 2. *Lapathum* folio acuto rubente.
 C. BAYH. *Pin.* 115. In Virginia. P.

Lapathum foliis acutis verticillatis. HALL. *Gott.*
 17. *Rumex* ad alas foliorum conglomeratis; foliis
 lanceolatis integerrimis. BOER. *Phil. Lyp.* 576. *Hy-*
drolapathum minus. LOBEL. *Icon.* p. 286. In
 foliis. P.

Lapathum foliis obtusis; ora crispa, calycis val-
vis interioribus serratis. HALL. *Gott.* 18. *Rumex*
obtusifolius. LINN. *Spec.* II. *Lapathum vulgare* sp-
 lio obtuso. I. BAVH. *Ad vias*. P.

H. *Lapathum arvense subhirsutum foliis retuso,*
caule longius brachiato; capsula seminis crenata.
 HALL. *Gott.* 18. In TILLII *hort. Pij.* p. 93. *Tup.*
 17. folium pinguis. Caulis ramosissimus ad angu-
 los rectos brachiatus. A.

Lapathum perenne foliis longissimis; calycibus
ciliatis. HALL. *l. c.* *Lapathum anthoxanthum*. L. B. II.
 987. In vdis circa Gottingam. In Saxonia superio-
 ri. Circa Berolinum.

H. *Lapathum annuum foliis lingulatis, verticillis*
densissimis, ciliis calycis longissimis. HALL. *l. c.*
Rumex aegyptiacus. LINN. *Sp.* 6. In Aegypto. A.

BISTORTA. HALL. 178.

Calyx quinquepartitus. Stamina octo. Tubae
 duae. Semen triquetrum. Radix contorta. Flores
 dense spicati.

Bistorta foliis ovato-oblongis acuminatis. LINN.
Cliff. 150. 207. 217. *Bistorta foliis imis appendicu-*
latis, semine triquetro. HALL. *Helv.* 178. *Polygonum*
Bistorta. LINN. *Sp.* 2. *Bistorta radice a. magis,* 3.
 minus

VAGINALES.

minus inserta. C. B. *Pin.* 192. *Bistorta* BLACKW. *T.* 254. & *offic. Krehwarzel, Schlangewurzel.* In pratis. ydis. P.

Bistorta foliis lanceolatis. LINN. *Cliff.* 150. *Bistorta* foliis ad oram nervosis, imis ovalibus, superioribus linearibus. HALL. *Helv.* 2. 179. *Polygonum viviparum.* LINN. *Sp.* 3. *Bistorta alpina* α . media & minor. In Alpibus. P.

PERSICARIA. HALL. 179. LVDW. 778.

Flores densius spicati. Stamina quinque ad octo. Tuba aut bifida aut duplex. Semen ovato compressum.

Persicaria foliis serratis, spica densa, cylindrica. HALL. *Helv.* 1. 180. *Persicaria* florum staminibus quinis corollam superantibus, stylo bifido. LINN. *Cliff.* 41. *Polygonum amphibium.* LINN. *Spec.* 7. *Potamogeton* falcis folio. BAVH. *Pin.* 193. In foliis folia laevia, & caulis procumbens, in siccioribus locis caulis erectus & folia hirsuta. Glandulae ad basin segmentorum.

Persicaria vaginis florum ciliaribus, spicis interruptis densioribus. HALL. *Helv.* 180. *Gott.* 28. *Persicaria* floribus hexandris digynis. LINN. *Fl. Suec.* 319. *Polygonum Persicaria.* LINN. *Sp.* 111. *Persicaria* mitis. BLACKW. *T.* 118. maculosa, & non maculosa. BAVH. *Pin.* 101. α . *Persicaria* foliis subtus tomentosis. *Helv.* 181. β . *Persicaria* angustifolia. BAVH. *Prodr.* 43. HALL. *Helv.* n. 11. *Flöb-Wein. Pfirsingkraut.* In hortis & ad vias. A.

7. *Perficaria* senis, spicis longis strigosis, vaginis non effatis. HALL. *Helv.* 181. GOTT. 29. *Perficaria* florum staminibus senis, stylo bifido. LINN. *Cliff.*

45. *Polygonum* *Hydropiper*. LINN. *Spec.* 10. *Perficaria* vrens. f. *Hydropiper*. BATH. *Pm.* 101. BLACKW., T. 119. *Wasserpfeffer*. In foliis. A.

8. *Perficaria* florum staminibus sex pluribusque, stylo duplici. LINN. *Cliff.* 41. *Perficaria* *Orientalis* Nicotianae folio, calyce florum purpureo. TOURN. *Cor.* 38. COMMEL. *rar.* 43. T. 43. HALL. *Gott.*

29. *Polygonum orientale*. LINN. *Spec.* 13. Glandulae octo ad basin segmentorum. In Oriente. A.

FAGOPYRVM. HALL. 172. LVDW. 794. HELXINE LINN.

Stamina octo, tubae tres, glandulae octo ad basin segmentorum. Flores laxè spicati.

9. *Fagopyrum* (*Helxine*) caule diffuso, foliis lanceolatis. LINN. *Hort. Vpf.* 96. *Polygonum* *divaricatum*. LINN. *Spec.* 18. *Perficaria* montana, angustifolia, racemosa. AMMAN. *Ruth.* 240. p. 167.

HALL. *Gott.* 29. Glandulae obscurae. In Sibiria. P.

10. *Fagopyrum* caule erecto, fructu ex calyce eminentè laevi. HALL. *Gott.* 29. *Helv.* 172. *Polygonum* *Fagopyrum*. LINN. *Sp.* 23. *Mat. med.* 188.

63. *Frumentum* *Sarracenicum*. MATH. 394. *Heydakorn*. *Barbancize*. Sponte in Asia. Nobis collatum. A.

11. *Fagopyrum* caule erecto, fructu ex calyce eminentè serrato. HALL. l. c. *Polygonum* *Tataricum*. LINN. *Spec.* 22. In Sibiria. Finnia. A.

Fago-

VAGINALES.

43

Pagopyrum scandens, semine calyce inclusis.
HALL. *Helv.* 173. *Helxine* caule volubili. LINN.
Lapp. 154. ROYEN. 217. *Polygonum Convolvula-*
ce. LINN. *Spec.* 104. *Pagopyrum vulgare scandens*.
L. ROM. 518. *Vogelzunge wild Heydskorn*. Vbi-
que in agris, & horti. A. 181. 182.

POLYGONVM. HALL. 181.

Stamina octo. Tubae tres. Glandulae nullae.
Flores solitarii albes.

Polygonum caule procumbente herbaceo, foliis
lanceolatis integerrimis. LINN. *Mat. med.* 188.
Polygonum aviculare. LINN. *Spec.* 15. *Polygo-*
num vulgare α . *angustifolium* β . *latifolium*. BAVH.
Pin. 281. *Polygonum* HALL. *Helv.* 182. BLACKW.
T. 319. *Wegwieser*. *Weggras*. In cultis & rudera-
tis. A.

Polygonum foliis lanceolatis, caule stipulis
obtectis fruticoso. *maritimum*. LINN. *Sp.* 8. *Po-*
lygonum perenne, vaginis amplissimis. HALL. *Gott.*
30. *Polygonum maritimum latifolium*. BAVH. *Pin.*
289. In Europa australi. LINNAEVS inter species
pentandras refert. P.

HERNIARIA. LINN. 274.

Calyx patens, interne coloratus. Stamina quin-
que sterilia, quinque abortiva. Tuba bifida. Glo-
meruli florum alares.

Herniaria calycibus bractea nudis. LINN. *Mat.*
med. 104. HALL. *Helv.* 181. *Herniaria glabra*.
LINN. *Spec.* 1. BLACKW. *T. 320.* *Polygonum tri-*
nus.

mix. f. Millegrana maior BAVH. *Pin.* 281. *Bruch-*
krant. In arenosis, siccis, v. g. circa *Northeim-*
Herzberg. A.

H. *Herniaria squamis nigris, flores superantibus.*
LINN. *Clus.* 4. H. ROY. 279. HALL. *Gott.* 31.

Illecebrum Paronychia. LINN. *Sp.* 4. *Paronychia*
hispanica. CLVS. *Hist.* 2. 183. In Gallia Narbo-
nensi & Hispania. A.

ILLECEBRVM. LINN. 256.

Calyx coloratus, cartilagineus. Stamina quinque.
Tuba simplex. Flores densa, verticillati.

Illecebrum floribus verticillatis nudis, caulis
procumbentibus. verticillatum. LINN. *Spec.* 1081.

Illecebrum. HALL. *Gott.* 31. *Illecebrum spuria.* R. & P.
99. In arenosis *Cellensis* & circa *Berolinam.* A.

Quae sequuntur tria genera ob habitum hic re-
linquenda, etsi petalis, numero staminum, aut capsu-
la polysperma differant.

MOLLYGO. LINN. 39.

Calyx interne coloratus. Stamina tria. Tubae
tres. Capsula polysperma.

H. *Mollugo* foliis quaternis obovatis, panicula
dichotoma. LINN. *Spec.* 2. *Triclis tristemon* foliis
coniugatis. HALL. *Gott.* 25. Vid. descriptionem &
iconem in *Comment. Soc. Sc. Gott.* T. V. Petala
quinque calyce minora. Folia opposita, quae ad ra-
morum ortum quaterna videntur. In Gallia Narbo-
nensi. A.

H. Mollugo foliis verticillatis tunciformibus acutis, caule subdiviso procumbente, pedunculis unifloris. *verticillata*. LINN. *Sp. 4. H. Vps.* 24. Vid. icon floris in *Comm. Gott. T. V. In Africa. Virginia. A.*

PHARNACEVM. LINN. 341.

Omnia fere prioris. Differt tantum staminibus quinque & floribus umbellatis.

H. Pharnaceum glabrum, pedicellis folia aequantibus. *Cerviana*. LINN. *Sp. 1. Phil. Bot.* 458. Triclis pentastemon, floribus umbellatis. HALL. *Comm. Soc. Sc. Gott. T. II. p. 346. T. XV. f. 2. Rostochii. In Sibiria. A.*

CORRIGIOLA. LINN. 340.

Calycis folia & petala & stamina quinque. Tuba trifida. Capsula monosperma. Folia alterna.

Corrigiola caule fasciculis florum terminato. LINN. *Cliff. 112. Spec. p. 271. HALL. H. 391. Polygonifolia. DILL. Nov. Gen. 95. Tab. III. In arenosis, v. g. Cellensibus.*

CLASSIS. VII.

TRICOCCAE.

Stamina in multis plura, quam segmenta calycis, in aliis segmentorum numero aequalia. Petala in aliis adsunt, in aliis desunt. Fructus tricoccus, trilocularis, tot styli quor loculi fructus. Sextus variat. Omnes acres, suspectae.

EUPHOR.

EUPHORBIA. LINN. 536. Tithymalus.

HALL. 189. TOURN. 18.

Ex marginē calycis monophylli petala quatuor aut quinque, crassa, plana. Fructus ex flore pendulus, petiolatus. Styli tres bifidi. Vbique lacte turgēt.

H. Euphorbia aculeata nuda subquingularis, aculeis geminatis. *Canariensis*. LINN. Sp. 2. Euphorbium tetragonum & pentagonum spinosum Canariinum. BOERH. p. 258. HALL. Gott. 32. Euphorbium. BLACKW. T. 340. f. 1. In Africa calidiori. Fr.

H. Euphorbia aculeata nuda multangularis, spinis solitariis subulatis. *cereiformis*. LINN. Sp. 5. Euphorbium cerei effigie, caulibus gracilioribus. BOERH. Lugdb. 1. p. 258. In Aethiopia. Fr.

H. Euphorbia aculeata nuda multangularis, aculeis geminatis. *officinatum*. LINN. Sp. 6. Euphorbium polygonum spinosum, cerei effigie ISNARD. Act. Par. 1720. T. 10. BLACKW. T. 340. f. 2. In Africa calidiori. Fr.

H. Euphorbia inermis imbricata tuberculis, foliolo lineari instructis. *caput Medusae*. LINN. Sp. 8.

α. Euphorbium erectum, aphyllum, ramis rotundis, tuberculis tetragonis BURM. T. 7. f. 2. H. Gott. 32.

β. Tithymalus aizoides africanus, simplici squamoso caule. COMMEL. Prael. 58. T. 8. In Aethiopia. Fr.

H. Euphorbia inermis nuda fruticosa, filiformis volubilis: cicatricibus oppositis. *viminalis*. LINN. Sp. 9. Felsel Taval. ALP. Aegypt. 190. T. 190. In Africae maritimis. Fr. H.

g. *Euphorbia inermis* seminuda fruticosa, filiformis, flaccida, foliis alternis. *Mauritanica*. LINN. Sp. 10. *Tithymalus aphyllor Mauritanicae*. DILL. Etb. T. 289. In *Africae* maritimis. Fr.

h. *Euphorbia inermis* subnuda fruticosa filiformis erecta, ramis parvis determinate confertis. *Ternatilis*. LINN. Sp. 31. *Tithymalus indicus frutescens*. COMMEL. H. Amst. L. p. 27. T. 14. In *India*. Fr.

i. *Euphorbia umbella* quadrifida, dichotoma, foliis oppositis integerrimis. *Lathyris*. LINN. Sp. 31. *Tithymalus*. HALL. Helv. n. 1. 189. *Lathyris* f. *Cataputia minor*. 1. B. III, p. 380. In *Helvetia*. *Gallia*. *Italia*. P.

Euphorbia (Tithymalus) foliis subhirsutis, ad caulem ellipticis, cymbis florum perfoliatis, petalis lunatis acutis. HALL. Helv. 190. *Euphorbia sylvatica*. LINN. Sp. 55. *Tithymalus sylvaticus lunato flore*. BAVH. Pin. 290. In sylvis. P.

k. *Euphorbia involucellis* perfoliatis emarginatis, petalis obtusis foliis integerrimis, hirsutis, caule frutescente. *Euphorbia Characias*. LINN. Sp. 56. *Tithymalus characias* L. CLUS. Hist. 2. 88. In *Gallia*. *Hispania*. Fr.

Euphorbia (Tithymalus) altissimus, levis, foliis sub foliis ternis, cordatis, fructu verrucoso. HALL. Helv. 191. n. 6. Gott. 34. *Euphorbia palustris*. LINN. Sp. 51. *Tithymalus palustris fruticosus*. BAVH. Pin. 292. P. In pratis paludosis *Thuringiae* *Helvetiae*

Euphor-

1 Euphorbia (*Tithymalus*) foliis glabris, serratis, acuminatis, involucris bifoliis, cordatis, fructu toto verrucoso. *Helv.* 191. *E. platyphyllus*. LINN. *Sp.* 47. *Tithymalus arvensis latifolius Germanicus*. B. *Pin.* 191. Vbique in agris, A.

H. Euphorbia (*Tithymalus*) foliis ovato-lanceolatis, subhirsutis, umbellae involucro quinquefolio. *HALL. Gott.* 34. Num Varietas?

H. Euphorbia (*Tithymalus*) foliis ovatis, spina terminatis, subhirsutis, petalis longe bicornibus. *HALL. Gott.* l. cit. A.

H. Euphorbia (*Tithymalus*) foliis ellipticis, spinula terminatis, petalis lunatis, serratis. *HALL. Gott.* 35. *E. Pitbyusa*. LINN. *Sp.* 35. *Tithymalus foliis brevibus aculeatis*. BAVH. *Pin.* 292. In arenosis Belgii, Hispaniae. P.

H. Euphorbia (*Tithymalus*) foliis linearibus spinula terminatis, radiis umbellae plurimis reflexis, petalis obtusis. *HALL. Gott. l. c. Helv.* 192. n. 10. *E. paralias*. LINN. 37. *Tithymalus maritimus*. BAVH. *Pin.* 291. In maritimis, P.

Euphorbia (*Tithymalus*) foliis linearibus, obtusis, petalis emarginatis. *Helv. l. c.* *Tithymalus amygdaloides angustifolius*. I. R. H. TABERN. p. 591. *MVPP.* p. 273.

1 Euphorbia (*Tithymalus*) foliis linearibus, obtusis, confertis, petalis lunatis. *HALL. Helv.* 193. n. 11. *E. cyparissias*. LINN. *Sp.* 49. Rarius circa *Gottin-gam* v. g. circa *Grossschneen*.

Euphorbia (Tithymalus) foliis linearibus obtusis, petalis argute bicornibus. *Helv.* 193. n. 12. *Esula exigua*. LINN. *Sp.* 29. C. BAVH. *Pin.* 291. Vbi-que in hortis. A.

Euphorbia (Tithymalus) foliis subrotundis serratis, radiis umbellae quinis latissimis. *Helv.* 194. n. 14. *E. Helioscopia*. LINN. *Sp.* 40. *Tithymalus helioscopius*. BAVH. *Pin.* 291. In cultis & hortis. A.

Euphorbia (Tithymalus) foliis ovatis, involu-
cris bifoliis cordatis, obtusissimis. *Helv.* 193. n. 13. *E. Peplus*. LINN. *Spec.* 27. *Peplus* f. *Esula rotunda*. BAVH. *Pin.* 292. A.

H. *Euphorbia* foliis subrotundis obtusis, floribus solitariis in foliorum alis. CVETT. *Stamp.* 2. p. 240. *Euph. Peplis*. LINN. *Sp.* 23. *Peplis maritima fœto obtuso*. BAVH. *Pin.* 293. *Peplis* CLVS. *Hist.* 2. p. 187. Petala argute bicornia, fructus in dorso valvarum verrucosus. In *Narbonae* maritima. A.

H. *Euphorbia* dichotoma, foliis serratis oblongis, floribus axillaribus solitariis, ramis patulis. *maculata*. LINN. 21. Synonymum FLVCK. 172. T. 65. f. 8. ad sequentem potius refererim. Folia inferiora maculata, superiora pura, petala obtusa, fructus glaber, laevis. In *America* septentrionali. A.

H. *Euphorbia* dichotoma, foliis crenulatis oblongis, floribus solitariis axillaribus, caulis procumbentibus. *Chamaesyce*. LINN. *Sp.* 22. *Chamaesyce*. BAVH. *Pin.* 293. CLVS. *Hist.* 2. p. 187. Habitus prioris, sed tota planta omni ex parte minor, & hirsuta, & procumbens. Folia semper maculata
D prioris

prioris spec. similia, petala integra, fructus hirsutus.
In *Sibiria* & *Europa* australi. A.

CLVTIA. LINN. 1009.

Pl. Diphyta. In utroque sexu calyx, quinquefolius, quinque petala, & nectaria intra petala posita. *Mas.* Stamina quinque ex columella media orta, patentia. *Fem.* Styli tres, bifidi. Capsula sexsulca.

H. *Clutia* foliis ovatis integerrimis, floribus lateralibus. *pulchella*. LINN. *Sp.* 2. *Clutia* foliis petiolatis. *Cliff.* 500. HALL. *Gott.* 36. In *Aethiopia*. Fr.

CNEORVM. LINN. 47.

Calyx tridentatus. Petala tria, totidemque stamina. Stylus trifidus.

H. *Cneorum tricocon*. LINN. *Sp.* 1. ROY. 819. *Chamalaea tricoctos*. BAVH. *Pin.* 462. HALL. *Gott.* 36. In *Hispania*, *Gallia Narbonensi*. Fr.

PHYLICA. LINN. 226.

Plurimi flores in discum collecti. Calyx proprius trifolius. Petalum limbo quinquefido; squamae stamina munientes. Stylus simplex.

H. *Phylica* foliis ovato-linearibus. ROY. 199. HALL. *Gott.* 36. Ph. *ericoides*. LINN. *Sp.* 1. *Alaternoides Africana ericae foliis, floribus albicantibus & muscosis*. COMMEL. *Hort. Amst.* II. p. 1. T. 1. In *Aethiopia*. Fr.

CROTON.

CROTON. LINN. 960.

n. Croton foliis ovatis tomentosis integris serratis. argenteum. LINN. Sp. 4. Cliff. 444. Flores in corymbo, superius masculo, inferius feminino. Calyx in femina multo maior ab exteriori latere tripartitus in folia oblonga tomentosa, quibus singulis ad basin squama glandulifera adhaeret, ab interiori fissus in duas facinias minimas. Petala nulla vidi. Fr. multo minor, calyx similis, filamenta staminum hirsuta ex receptaculo hirsuto, vix coalita. Cons. vix. ad III. WERLHOFF; ubi fassus hanc descripsit.

JATROPHA. LINN. 961.

Monoph. Mar. Petalum hypocrateriforme limbo quinquefido. Stamina decem. Fem. Corolla pentapetala. Styli dichotomi.

n. Jatropha foliis cordatis angulatis. LINN. Sp. 4. Mar. med. 439. HALL. Gott. 36. In America calidiori. Fr.

RICINUS. LINN. 962.

Monoph. Mares inferius positi. Calyx quinquepartitus. Filamenta staminum inferne ramosae coalita. Fem. Calyx tripartitus. Styli bipartiti,

n. Ricinus foliis peltatis serratis: petioli glanduliferis. LINN. Cliff. 450. Mar. med. 438. HALL. Gott. 37. Ric. communis. LINN. Sp. 1. Ricinus vulgaris. BAVH. Pin. 439.

Var. a. Ricinus amer. Fructu glabro.

β. Ricinus caule rutilante. Sponte in Africa, Asia, America & Europa calidiori. P. nobis annua.

ACALYPHA. LINN. 959.

Monoph. Mares plurimi in sulum fere conferti, calyce minimo quadrifolio; stamina vt in Ricino. Feminae plures involucri foliaceo magno crenato exceptae. Styli ramosi.

H. Acalypha involucri femineis cordatis incilis, foliis ovato lanceolatis petiolo longioribus. *Virginica*. LINN. Sp. 1. Mercurialis triccocos hermaphroditeq. ad foliorum iuncturas & foliis cristatis infusca simul & fructum ferens. BURM. Zeylan. 248. T. 99. fig. 4. Sponte in Ceylona. Virginia, A.

CLIFFORTIA. LINN. 1004.

Diph. Mas. Calyx trifolius. Stamina plurima. Fem. Calyx similis germini insistent. Fructus bilocularis.

H. Cliffortia foliis subcordatis dentatis. *ilicifolia*. LINN. Sp. 1. Hort. Cliff. 463. T. 36. Arbutula *Afrasolio acuto ilicis caulem amplexo rigido*. DILL. Elsb. 36. T. 31. f. 38. In Aethiopia. Fr.

MERCVRIALIS. LINN. 998.

HALL. 194.

Diph. Mares in inulos congesti, calyce trifolio, staminibus duodecim circiter. Feminae solitariae. Styli duo reflexi. Capsula bilocularis.

Mercurialis caule simplicissimo, foliis scabris perennis. LINN. Sp. 1. HALL. Helv. 194. 2. Gott. 37. Mercurialis montana a. testiculata β. spicata. SAVH. Pin. 122. In sylvis primo vere. P.

Mer-

TRICOCCAE.

33

Mercurialis caule brachiato, foliis glabris: *anaya*. 1
LINN. *Spec.* 2. *Mat. med.* 464. HALL. *Helv.* 194.
1. *Mercurialis* BLACKW. T. 168. & *offic. Binglei*,
krout. In ruderalis, umbrosis. A.

Quae sequuntur tria genera, partibus quibusdam
anomala, ob habitum tamen hic relinquenda.

THELYGONVM. LINN. 947.

H. Monoph. *Mat.* Calyx bifolius stamina duo,
decim. *Fem.* Fructus globosus unilocularis, stylus
simplex.

H. *Thelygonum Cynocrambe*. LINN *Sp.* I. HALL.
Got. 38. *Cynocrambe Dioscoridis*. BAVH. *Pih.* 122.
Prodr. 59. In *Italia, Monspelii*. P.

ASARVM. LINN. 522. HALL. 195.

Calyx monophyllus, semitrifidus. Stamina duo,
decim, stylus stigmate sexfido. Fructus sexlocularis
sub calyce.

Asarum foliis reniformibus obtusis, *Europaeum*,
LINN. *Sp.* 1. *Mat. med.* 219. *Asarum*. HALL. *Helv.*
195. BLACKW. T. 383. & *offic. Haselwurz*. In sylvis, P.

ARISTOLOCHIA. LINN. 911. HALL. 196.

Petalum ex tubo ventricoso ligulatum, sub flo-
re capsula sexlocularis; stylo brevi, sub stigmate sex-
partito stellato, adnascuntur sessiles antherae sex
quadriloculares.

H. *Aristolochia* foliis cordatis, caule erecto, flori-
bus axillaribus confertis. LINN. *Sp.* 13. *Aristolochia*,
Clematitis. BLACKW. 255. *vulgaris* CLVS. *Hist.* 2.
LXXI. Sponte in *Alsatia, Helvetia, Gallia*. P.

11. *Aristolochia* caule infirmo ramoso, foliis cordatis crenulatis petiolatis, floribus solitariis LINN. *Spec.* 10. *Aristolochia Pissolochia dista.* BAVH. *Pin.* 307. In *Hispania: Gallia Narbonensi.* P.

CLASSIS VIII.

INVNDATAE LINN.

Oculis herbaceus, geniculatus. Folia verticillata, aut opposita. Flores aut in alis foliorum, aut in spicam digesti. Stamina unicum ad octo. Semina saepe nuda, totidemque semper tubae. Omnes in paludosis aut stagnis, perfecta florentia sub aqua latent.

SALICORNIA. LINN. 10.

Habitus aphyllus. Flores in conum congesti. Calyx truncatus. Stamen unicum. Stylus apice bifido. Semen unicum.

11. *Salicornia* articulis, apice crassioribus obrutis. *Europaea.* LINN. *Sp.* 1. *Mat. med.* 8. *a. annua.* *B. sempervirens.* *Salicornia* HALL. *Gott.* 5. *Kali genti- culatum* BAVH. *Pin.* 289. *Soda offic.* Ad aquas sal- sas & in litoribus marinis.

CHARA. LINN. 1066.

Folia verticillata in quorum alis anthera sessilis. Semen unicum.

Chara catulis laevibus: *vulgaris.* LINN. *Spec.* 2. p. 1156. ROY. 533. *Chara vulgaris foetida.* VAILL. HALL. *Gott.* 38. *Hell.* 198. n. 1. In aquis pigris.

LIMNO.

INVNDATAE.

55

LIMNOPEUCE. HALL. 197. Hippuris

LINN. II.

Folia verticillata, in quorum alio semen unicum & stamen unicum.

Linnopeuce. HALL. Gott. 38. Helv. 197. Hippuris vulgaris. LINN. Sp. 1. ROY. 205. In rivis & stagnis, P.

ZANNICHELLIA. LINN. 920.

HALL. 198.

Folia coniugata, in eadem ala folii, flos unus stamine unico, & femina feminibus quatuor nudis, stigmate lato patente, calyce communi exceptis.

Zannichellia tubae ligula integra. HALL. Helv. 198. Gott. 1. c. Zannichellia palustris. LINN. Sp. 1. Vid. iconem MICHEL. gen. 71. T. 34. f. 1. 2. In vivo ad Rasmühle. A.

STELLARIA. HALL. 198. Callitriche.

LINN. 13.

Folia coniugata. Flores alares, mares stamine unico, feminae fructu biloculari. Vtrique sexui petala duo lunata.

Stellaria foliis omnibus angustis apice resecto. HALL. Helv. 199. n. 1.

β. Stellaria foliis ad caulem angustis, in summitate subrotundis. HALL. 1. c. n. 2.

γ. Stellaria foliis omnibus subrotundis. HALL. n. 3. Callitriche palustris LINN. Sp. 1. omnes, ut varietates, iungit. In paludibus, A.

POTAMOGETON, LINN. 16p.

HALL. 199.

Flores spicati ex alis foliorum. Petala quatuor, totidemque stamina antheris gemellis, & semina nuda quatuor.

1. Potamogeton foliis oblongo-ovatis petiolatis, natans. LINN. Sp. 1. HALL. Helv. l. c. n. 1. Potamogeton rotundifolium. BAVH. Pin. 193. In fluviis & stagnis.

Potamogeton foliis ovatis longissime petiolatis. Helv. 199. Potamogeton foliis lanceolatis planis in petiolos desinentibus. lucens. LINN. Sp. 4. Fontinalis lucens maior. BAVH. Hist. III. p. 769. In stagnis. v. g. Lichtenmeer auf dem Heimberg.

2. Potamogeton foliis cordatis amplexicaulibus. perforiatum. LINN. Sp. 2. HALL. Helv. 199. n. 3. Potamogeton rotundifolium alterum LOESEL. Pruff. ic. 65. In lacubus. v. g. lacu Seeburgensi ad Roerenshausen.

3. Potamogeton foliorum oris plicatis, molliter spinosis. Helv. 202. n. 5. Potamogeton crispum. LINN. Sp. 5. Fontinalis crispa. 1. B. Hist. III. p. 778. Potamogeton foliis crispis f. Lactuca ranarum. BAVH. Pin. 193. In rivis v. g. am Schützenhof.

4. Potamogeton foliis lanceolatis oppositis acuminatis. setaceum. LINN. Sp. 2. HALL. Helv. 200. n. 6. Fontinalis media lucens. 1. B. III. p. 777. In aquis lente fluentibus.

Potamogeton foliis planis longissimis, spica densa & crassa. HALL. l. c. n. 8. Potamogeton racemosum

angn-

angustifolium. G. B. *Prodr.* p. 101. n. 2. In rivo
am Schützenbause & alibi.

Potamogeton foliis linear-lanceolatis alternis
sessilibus, stipula latioribus. *gramineum*. LINN. *Sp.*
10. Potamogeton HALL. *Helv.* n. 7. Potamogeton
gramineum latifolium, LÖESEL. *icon.* 66. p. 206.
In fluviis lente fluentibus vt in *Leina*.

Potamogeton foliis gramineis vndulatis spicis exi-
guis. HALL. *t. c.* n. 9. *P. compressum*. LINN. *Sp.* 7.
Potamogeton caule compresso, folio graminis canti-
ni. RAI. *Hist.* 189. In fossis paludosis.

Potamogeton pedunculis florigeris longissimis,
glomerulis florum remotis. HALL. *l. c.* n. 11. Po-
tamogeton *marinum*. LINN. *Sp.* 11. Potamogeton
pusillum. BOCCON. *Sicil.* 42. T. 20. f. 5. Circa
Göttingam.

MYRIOPHYLLVM. LINN. 945. Pentapteris.

HALL. 201.

Monophyta. *Mas.* Calyx quadrifolius, stamina
octo. *Pem.* Capsula tetrasperma, tubis quatuor.
Interdum & androgyni adfunt. Folia verticillata.

Myriophyllum (*Pentapteris*) scapo florifero nu-
do. HALL. *Gott.* 40. Myriophyllum *spicatum*
LINN. *Sp.* 1. Myrioph. *aquaticum pennatum spir-*
catum. G. BAVH. *Prodr.* 73. T. 73. Flores masculi
li verticillati quaterni, staminibus, calyce explicato,
longe pendulis, feminini calyce trifolio mihi ob-
servati sunt. In lacu *Seeburgensi*.

Myriophyllum (*Pentapteris*) floribus in alis sessilibus. *Helv.* 202. n. 2. *M. verticillatum*. LINN. Sp. 2. Millefolium aquaticum flosculis ad foliorum nodos. *BAVL. Pin.* 141. In fossis v. g. am Schützenhaufs.

CERATOPHYLLUM. LINN. 244.

HALL. 202.

Pl. monophyta. *Mas.* Calyx multipartitus, stamina plurima. *Fem.* Calyx similis, capsula monosperma, acuminata. Folia verticillata, flores alares.

Ceratophyllum foliis asperis, bis bifurcatis. *HALL. l. c.* *C. demersum*. LINN. Sp. 1. Dichotophyllum. *DILL. nov. Gen. p.* 100. T. 3. In fossis.

CLASSIS. IX.

GRAMINEAE.

In omnibus amplissima huius classis plantis caulis, qui *culmus* dicitur, est simplex, articulatus, foliis vaginantibus cinctus; folia longa sunt, angusta, integerrima, sine acuto: flores in summitate caulis aut in spicam aut in paniculam dispositi, simplices, aut in capitulum, quod *lobusa* dicitur, collecti: calyx & folliculus glumofus, ut plurimum bivalvis; stamina in plurimis tria, filamentis capillariibus, antheris sagittatis; ex apice germinis stylus duplex aut simplex: stigmata plumosa. Semen vnicum folliculo comprehensum.

ORDO.

ORDO. I.

GRAMINA.

Calyx biglumis, rarius vnglumis, & triglumis.
 Folliculus bivalvis, (rarius univalvis,) valvula interiori planiori, exteriori concava, saepe aristata.
 Stamina tria, (excepto *Anthoxantho*, cui duo,) styli duo, semen unicum.

I. Locusta uniflora.

ANTHOXANTHVM. LINN. 40.

Calyx & folliculus bivalvis, hic aristatus, nectarium diphyllum. Stamina duo.

Anthoxanthum spica ovaro-oblonga, flosculis, subpeduncularis aristis longioribus, *odoratum*. LINN. *Spec. 1. Gr. pratense spica flavescente*, BAVH. *Pin. 3. In ficcis.*

NARDVS. LINN. 65.

Calyx nullus, Folliculus bivalvis. Stylus simplex.
 ex LINN.

Nardus spica setacea recta. *stricta*. LINN. *Sp. 1. Gramen sparteum suncifolium*. BAVH. *Pin. 5. In sylva bercynica. v. g. am dñe Oderbrücke.*

PHALARIS. LINN. 69.

Calyx biglumis; folliculus bivalvis; panicula spiciformis.

Phalaris, panicula subovata, glumis carinatis.
Canariensis. LINN. *Sp. 1. α. Phalaris maior semine albo aut nigro. β. Pin. 28. β. Gramen spicatum perenne.*

GRAMINA.

perenne, semine miliaceo, fibrosa radice. HALL. Gott. 51. In Europa australi. Canariis.

Phalaris panicula oblonga ventricosa. arundinacea. LINN. Sp. 3. Gramen *arundinaceum spicatum.* B. Pin. 6. Th. 94. Gr. *aquat. paniculatum phalaridis semine.* SCHEVCHZ. HALL. Gott. p. 46. In stagnis & fossis.

Phalaris panicula cylindrica. phleoides. LINN. Sp. 11. Gramen *typhoides asperum primum.* C. B. Pin. 4. HALL. Gott. 50. In versuris agrorum.

PHLEVM. LINN. 71. Typhoidis spec.
HALL.

Calyx biglumis, apice bicuspidate, glumis superne latis aristatis. Folliculus bivalvis. Spicatum.

Phleum spica cylindrica longissima. pratense. LINN. Sp. 1. Gramen *typhoides maximum, spica longissima.* C. B. Pin. 4. In pratis & agris.

β. Gramen *typhoides asperum alterum.* C. B. Th. p. 32. Caulis fere bulbosus. Num varietas?

ALOPECURVS. LINN. 72. HALL.
Gott. 42.

Calyx biglumis. Folliculus univalvis, ex basi arista. Spicatus.

Alopecurus culmo spicato erecto. pratensis. LINN. Sp. 1. Gramen *spicatum, spica cylindracea longiore tenuissima.* SCHEVCHZ. 69. In pratis v. g. auf dem Rosdorffer Weg.

Alope-

GRAMINA



Alopecurus culmo spicato infracto, geniculatus. LINN. Sp. 2. *Alopecurus culmo infracto aristis glau-
cis longioribus.* ROY. 34. *Gramen aquaticum gen-
culatum spicatum.* C. B. Pin. 3. In fossis natadleri.

PANICVM. LINN. 70. HALLERI Panici

et Mili species.

Calyx triglumis, glumis teretibus minimis, post ala-
ram posita. Folliculus bivalvis interdum aristatus.
Flores spicati aut paniculati.

H. *Panicum spica composita, aristis flosculis he-
vioribus.* LINN. Hort. Vps. 19. *Panicum italicum.*
LINN. Spec. 4. *Panicum italicum f. panicula ma-
ior.* B. Pin. 27. In Indiis. A.

*Panicum spica tereti, involuclis bifloris fascicu-
lato-pilosis, glaucum.* LINN. Sp. 2. Var. β. *Gram-
panicum f. panicum sylvestre, spica simplici.*
SCHEVCHZ. 46. In agris, exeunte aestate.

Panicum spicis alternis remotis, laxis, compositis.
LINN. Cliff. 27. *Pan. Crus Galli.* LINN. Sp. 5.
Gr. *panicum, spica divisa.* C. B. Pin. 8. Ad fossas.

Panicum spica composita, aristis spica longioribus. ROYEN. 55. *Gynosurus panicum.* LINN. Sp. 8. *Gram. panicum spica aspera.* B. Pin. 8. In agris cultis.

Panicum spicis aggregatis, basi interiori nodosis, flosculis geminis muticis, vaginis foliorum punctatis. *sanguinale.* LINN. Sp. 8. *Gram. distylon satis latiore.* C. B. Tb. 119. HALL. Gott. 146. In segetibus auf dem Hagen.

11. *Panicum panicula laxa flaccida, foliorum vaginis pubescentibus. Milicium* LINN. *Sp.* 16. *Adat.* med. 34. *Milium semine lato & albo.* 2. *Poa* aff. In India. A nostris colitur.

MILIUM, LINN. 73.

Calyx & folliculus bivalvis, muticus, valvulis ovalibus. Stigmata penicilliformia. *Paniculatum.*

20. *Milium floribus dispersis. effusum* LINN. *Sp.* 1. *Gramen sylvaticum, panicula multica sparsa.* C. B. *Poa* 8. In Europae umbratis nemoribus.

AGROSTIS, LINN. 74.

Locusta acuminata. Calyx & folliculus bivalvis, aristatus. *Paniculata.*

Agrostis petalo exteriori exsertente aristam rectam stridam longilamam. Spica densa. LINN. *Spec.* 1. *Gram. segetum panicula sparsa.* C. B. (*Bromus*.) *Agrostis. Holo.* 25. Inter segetes.

Agrostis panicula inferiori verticillata laxa, superior contracta. LINN. *Lapp.* 46. *Agrostis radia.* LINN. *Sp.* 4. *Gram. serotinum* 270. *panicula contracta pyramidalis. (Arundo.)* HALL. *Gest.* 46. *By. Stollberg.*

Agrostis panicula tenuissima. LINN. *Lapp.* 43. *Agrostis capillaris.* LINN. *Sp.* 8. *Gram. mont. panicula spadicem delicatiora.* B. *Prodr.* 12. (*Poa*.) HALL. *Det.* 46. In pratis & locis siccis.

LAGURVS. LINN. 36.

Calyx biglumis, glumis arista villosa. Folliculi valvula maior arista triplici, media e dorso longissima. Spicatus.

H. Lagurus spica ovata. LINN. Sp. 1. Gram. alopecuroides spica rotundiore. BAVH. Pin. 4. (Typhoides) HALL. Gott. 56. In Europa Australi. A.

II. Locusta multiflora.

AIRA. LINN. 75.

Calycis biflori glumae ovato-acutae. Folliculus calyci similis. Fl. paniculati.

Aira panicula spicata, floribus muticis pedunculo longioribus; glumis acuminatis inaequalibus, cristatis. LINN. Sp. 2. Gram. spica cristata subirsutum. C. B. Pin. 2. Prodr. 8. (Poa) HALL. Gott. 46. In siccoribus v. g. Auf dem Heymberg.

Aira foliis planis, panicula patente. floribus muticis laevibus calyce longioribus. aquatica. LINN. Sp. 6. Gramen aquaticum miliaceum. SCHEVCHZ. 228. HALL. Helv. 218. In aquis. v. g. ad viam versus Geismar.

Aira foliis planis, panicula patente, petalis basi villosis aristatisque; arista recta brevi. cespitosa. LINN. Sp. 8. Gramen segetum panicula arundinea. C. B. Pin. 3. In agris. B.

Aira foliis setaceis, panicula angustata, flosculis basi pilosis aristatis; arista tortili longiore. montana. LINN. Sp. 14. Gram. avenaceum capillaceum, minoribus

noribus glumis. C. B. *Pin.* 10. (*Avena*.) HALL. *Helv.* 228. In sylvis super Hardeggen. B.

Aira flore, masculino aristato, femineo mucoso. LINN. *Cliff.* 27. *Holcus lanatus*. LINN. *Sp.* 4. Gram. *pratense paniculatum molle*. C. B. *Prodr.* 5. In vdis.

MELICA, LINN. 76.

Calyx & folliculus bivalvis, muticus. Corpusculum inter flosculos. Paniculata.

Melica petalis imberbibus, panicula nutante simplici. *nutans*. LINN. *Sp.* 2. Gram. *montanum avenaceum, locustis rubris*. C. B. *Prodr.* 10. In rupibus. Vm *Pless*, & in *M. Hengst*.

POA, LINN. 77.

Locusta ovato-oblonga mutica, disticha. Calyx & folliculus vt in Aira. *Fl. Paniculata*.

Poa flosculis confertis vno versa dispositis LINN. *Cliff.* 28. *Dactylis glomeratus*. LINN. *Sp.* 2. Gr. *spicatum, folio aspero*. C. B. *Prodr.* 9. HALL. *Helv.* 231. Ad vias.

Poa panicula diffusa, spiculis sexfloris linearibus. *aquatica*. LINN. *Sp.* 1. Gram. *palustre paniculatum altissimum*. C. B. *Pin.* 2. Im *Deuskerhäuser-Tisch*.

Poa panicula diffusa, spiculis quinquefloris basi pubescentibus, culmo erecto tereti. *trivialis*. LINN. *Sp.* 3. Gram. *pratense paniculatum medium*. C. B. *Pin.* Vbique in pascuis.

Poa panicula diffusa, spiculis quadrifloris pubescentibus, culmo erecto tereti. *angustifolia*. LINN. *Sp.*

Sp. 4. Gr. *pratense paniculatum maius angustiori folio*. C. B. Pin. 2. Prodr. 6. In versuris agrorum.

Poa panicula diffusa, spiculis trifloris glabris, culmo erecto, rereci. *pratensis*. LINN. Spec. 5. Gram. *pratense paniculatum maius* C. B. Pin. 2. In pratis.

Poa panicula diffusa angulis rectis, spiculis obtusis, culmo obliquo compresso. *annua*. LINN. Sp. 6. Gram. *pratense paniculatum minus*. C. B. Pin. 2. Ad vias.

Poa panicula coarctata secunda, culmo obliquo compresso. *compressa*. LINN. Sp. 15. Gram. *murorum radice repente*. C. B. Prodr. 2.

Poa panicula attenuata, spiculis subbifloris mucronatis scabris, culmo incurvo. *nemoralis*. LINN. Sp. 16. α. Gr. *panicul. angustifolium alp. locustis varioribus & angustioribus non aristatis*. SCH. 164. β. Gr. *cirrboſa & villoſa ſpongia ad ſingula genicula donatum*. ROCCON. Mus. 2. p. 70. T. 59. Ad radices vmbroſas montium.

BRIZA. LINN. 78.

Locusta cordata disticha. Calycis valvulae cordatae. Paniculata.

Briza spiculis ovatis, flosculis septem. *media*. LINN. Sp. 2. Gram. *tremulum maius*. C. B. Pin. 2. In siccioribus & pratis.

h. Briza spiculis cordatis, flosculis septemdecima. *maxima*. LINN. Sp. 3. Gram. *tremulum maximum*. C. B. Pin. 2. In Italia. Monspeltii.

FESTUCA. LINN. 82.

Locusta oblonga, apice acuminato. Folliculus bivalvis aristatus. Culmus alternis incisuris exsculptus, quem media plana facies locustae respicit.

Festuca panicula secunda coarctata aristata, culmo tetragono nudiusculo, foliis sataceis. *ovina*. LINN. *Sp.* 1. *Gt. foliis lanceis brevibus, maius, radice nigra.* C. B. *Prodr.* 34. In locis ficcis.

Festuca panicula nutante inferne ramosa, spicis adscendentibus hispidis, foliis setaceis. *duriuscula*. LINN. *Spec.* 2. *Gr. pratense, panicula duriore laxa, unam partem spectante.* RAII *hist.* 1286. In ficcis.

TRITICVM. LINN. 94.

Locusta saepius triflora, obtusa. Folliculi valvula exterior obtusa cum acumine. Culmus vt in priori.

Triticum radice annua spica glabra aristata. *aestivum*. LINN. *Spec.* 1. Triticum *aestivum*. C. B. *Pin.* 21. *Sommer-Weitze.* Rarius seritur.

Triticum radice annua, spica mutica. LINN. *Cliff.* 24. Triticum *hybernum*. LINN. *Sp.* 2. *Winter-Weitze.* A nostris seritur.

Triticum flosculis aristatis hermaphroditis, intermedio neutro. *Spelta*. LINN. *Sp.* 4. *Zea dicoccos vel Spelta maior.* B. *Pin.* 22. *Spelt.*

Triticum radice repente, foliis viridibus. LINN. *mat. med.* 37. Triticum *repens*. LINN. *Sp.* 6. Gramen *caninum arvense* s. *gramen Dioscoridis*. B. *Pin.* 1. HALL. *Helv.* 2. *Quecken.* In hortis & arvis.

Triti-

Triticum. Gramen latifolium spica triticea compacta, G. B. *Prodr.* 53. HALL. *H.* 207. n. 1.

β. Gramen latifolium, spica triticea divulsa. C. B. *Prodr.* 54. HALL. *l. c.* β. ad rivos.

Triticum caninum. LINN. *Sp.* 7. Gramen *foliaceum*, radice repente, f. *Gram. offic.* longioribus aristis donatum. HALL. *Helv.* 208. Gramen *spica triticea compacta hirsutum aristatum*. HALL. n. 6. Num duae haec plantae modo dictae varietates sunt *graminis canini vulgaris officinarum*?

Triticum culmo indiviso, spiculis alternis subsessilibus. *Bromus pinnatus*. LINN. *Sp.* 10. Gramen *foliaceum corniculatum*. TVERN. *Gram. spica Brizae maius*. C. B. *Prodr.* 53. In sylvis.

β. Gramen *foliaceum corniculatum spicis glabris*. I. R. H. *Festuca dumetorum*. C. B. *Prodr.* 69. In pratis.

LOLIUM. LINN. 90.

Locustae oblongae acuminatae distichae, dorso culmi alternis incisuris obversae & appressae. Calyx uniglumis, folliculus bivalvis. Fl. spicati.

Lolium spica mutica. perenne. LINN. *Sp.* 1. *Gram. foliaceum angustiori folio & spica*. C. B. *Pin.* 9. Ad vias & versuras agrorum.

Lolium spica aristata. temulentum. LINN. *Spec.* 2. *Gram. foliaceum spica longiore f. Lolium Dioscoridis*. C. B. *Pin.* 9. In segetibus vdiis.

Lolium spicis mediis latioribus. ROY. p. 69. Gr. *loliaceum spicis brevioribus & compressis.* MORISON. III. S. VIII. T. 2. f. 2. Ad vias.

BROMVS. LINN. 83.

Locusta ovata. Calyx biglumis, muticus. Folliculus aristam infra apicem emittit. Spiculæ paniculatae.

Bromus panicula patente, spicis ovatis, aristis rectis. *secalinus.* LINN. Spec. 1. Gr. *avenaceum locustis villosis crassioribus.* 1. R. H. Festuca *graminea*, *glumis hirsutis.* C. B. Pin. 9. Inter segetes.

Bromus panicula patula, spiculis oblongis distichis, glumis subulato-aristatis. *sterilis.* LINN. Sp. 5. Bromus herba, DODON. Cereal. p. 141. Ad vias.

Bromus panicula nutante, spiculis ovato-oblongis. *arvensis.* LINN. Spec. 6. Festuca *graminea iuba effusa.* C. B. Pin. 192.

Bromus panicula nutante, spiculis linearibus. *testorum.* LINN. Spec. 7. Gr. *avenaceum, locustis villosis & candicantibus.* 1. R. H. 526. In agris.

Bromus. HALL. Herb. n. 7. p. 229. Gram. Typhoides molle. C. B. Tb. 50. Gr. *spicata spica cylindracea molli & densa.* 1. R. H. Folia lanuginosa. Ad ripas *Leinae.*

AVENA. LINN. 85.

Locusta laxa. Calyx muticus. Ex dorso folliculi arista geniculo reflexa, spiralis. Fl. paniculati.

Avena

GRAMINA.

63

Avena foliis pubescentibus, panicula contracta, flosculo hermaphrodito mutico, masculo arista, uncinata, calyce brevibre. *Holcus lanatus*. LINN. *Sp. 4. Gr. pratense paniculatum molle*. BAVH. *Prodr.* 3. *Gr. lanatum*. DALECH. 425. In pascuis & locis vdis.

Avena calycibus bifloris, panicula laxe spicata pedunculis prioribus fasciculatis. EVETT. *Avena elatior*. LINN. *Sp. 2. Gr. avenaceum, iuba longe splendente*. RAIL *meth.* 179. In lapidosis. v. g. versus *Herperhausen*.

β. *Gr. nodosum avenacea* panicula, radice tuberibus praedita. C. B. *Prodr.* 3.

Avena calycibus dispermis, seminibus laevibus. *fatua*. LINN. *Sp. 5. Mat. med.* 38. *Avena nigra*. β. *alba* C. B. *Pin.* 23. *Haber.* Copiose feritur.

Avena panicula patente, calycibus trifloris, flosculis basi pilosis. *fatua*. LINN. *Sp. 6. Gr. avenaceum, locustis lanugine flavescentibus*. SCH. 239. *Wild Haber. Flug Haber.* Inter segetes.

Avena panicula laxa, calycibus trifloris brevibus, flosculis omnibus aristatis. *flavescens*. LINN. *Sp. 7. Gr. avenac. pratense elatus, panicula flavescente, locustis parvis*. SCH. 223. In sepibus versus *Edigehausen* & circa *Reinhausen*.

Avena calycibus quinquefloris, panicula spicata. *pratensis*. LINN. *Sp. 9. Gr. avenac. mont. spica simplici, aristis recurvis*. SCH. 226.

β. *Gr. avenaceum*, panicula purpureo argentea splendente. SCHEVCH. 226.

ARVENDO. LINN. 87.

Locusta vni-vel multiflora, mutica. Folliculus intus lanuginosus, cum calyce bivalvis. Fl. paniculati.

H. Arundo calycibus trifloris, panicula diffusa. *Denax*. LINN. Sp. 2. Arundo *sativa*. BAVH. Pin. 17. Arundo *Indica Lacōnica versicolor*. Cliff. 26. Arundo *Americana striata*. MUNT. F. 108. Variet. In *Hispania*. Gallo-*Provincia*.

Arundo calycibus quinquefloris, panicula laxa. *phragmites*. LINN. Sp. 3. Arundo *vulgaris* f. *phragmites* *Dioscoridis*. C. B. Pin. 17. *Schilf-Rohr*. In lacubus & fossis.

STIPA. LINN. 84.

Locusta vniflora, laxa, biglumis. Folliculus aristata longa ex apice prodeunte, basi articulata. Fl. paniculati.

H. Stipa aristis lanatis, *pennata*. LINN. Sp. 1. *Gasperteum pennatum*. C. B. Pin. 5. In *Marchia*. *Helvetia*. *Gallia*.

H. Stipa aristis nudis, *iuncea*. LINN. Sp. 2. *Festuca iuncea folio*. C. BAVH. Pin. 9.

CYNOSVRVS. LINN. 81.

Involucrum laterale trifolium, saepe pinnatifidum. Calyx biglumis. Folliculus aristatus.

Cynosurus bracteis pinnatifidis. *cristatus*. LINN. Spec. 1. Gramen *pratense cristatum*. B. Prodr. 8. In pratis.

Cyno-

Cynosurus bracteis integris. corymbosus. LINN. Sp. 5. Gr. glumis variis. C. B. Prodr. 21. Ad rupes praeruptas M. Plesi.

H. *Cynosurus paniculae spiculis sterilibus pendulis ternatis, floribus aristatis. aureus. LINN. Sp. 9. Gramen panicula pendula, aurea. C. B. Pin. 3. BALL. Gott. 49. ubi vide characterem a reliquis spicibus diversissimum.*

HORDEVM. LINN. 93.

Praeter involucrium triflorum, ex sex aristis compositum, calyx nullus. Folliculus aristatus.

H. *Hordeum flosculis omnibus hermaphroditis aristatis, ordinibus duobus erectioribus. vulgare. LINN. Sp. 1. Hordeum polystichon vernum. C. B. Pin. 22. Gerste. Ad cerevisiam coquendam inprimis feritur.*

Hordeum flosculis lateralibus masculis aristatis. murinum. LINN. Sp. 5. Gram. hordeac. minus & vulgare. B. Pin. 8. β. Gr. spicatum secalinum minus. SCH. 17. Vbique ad sepes.

Hordeum flosculis lateralibus imperfectis, mere aristatis. Gramen hordeaceum montanum spica strigosa, brevius aristata. HALL. Helv. 204. Gott. 42. Gr. secalinum maius. In sylvis über Herperhausen.

SECALE. LINN. 92.

Involucri biflori aristae duae. Folliculus bivalvis, aristatus.

H. Secale glutinarum ciliis scabris. *Linnaeus*. LINN. Sp. 1. α . hybernium vel maias. β . vernum vel minus. B. Pin. 23. Korn. Rocken. Vbiq. colitur.

GRAMINIBVS ADFINES.

Culmo, foliis & tota reliqua habita, pleraeq. etiam staminibus, cum prioribus conveniunt, sed in flore diuersitas est. His enim stylis ut plurimum vnicus & semen est vel vnicum, in bivalvi quidem folliculo, sed flores sexu sunt distincti; aut semen vnicum, ad vnicam glutinam; aut semen vnum, plurave capsulis tecta, cum flore apetalis, sexfolijs & staminibus tribus vel sex.

Semen stylo bicorni in folliculo biglumi, flores sexu distincti.

AEGILOPS. HALL. Gott. 52.

In eadem locusta tri- aut quadriflora, flores & hermaphroditi laterales, & masculi intermedi. Calyx biglumis, aristatus. Folliculi bivalvis glutina exterior triplici arista terminata, interior mutica, quae semen hirsutum obvoluit. Petala duo plumosa. Styli duo. Vide quomodo differat character Linnaeus Gen. 1018.

H. Aegilops spica ovata, aristis brevioribus. *ovata*. LINN. Sp. 1. Aegilops DON. Pempt. 73. Cereal. p. 143. Festuca altera capitulis duris. RAVH. Pin. 10. In Europa australi. B.

GENCHRVS. LINN. 1017. Panicastrella

MICH. 31.

Capitula ex locustis bifloris, ternis calyce communi echinato exceptis, in spicam disposita. Flores alter *androgynus*, alter *masculus*. Calyx proprius & folliculus bivalvis, acuminatus.

H. Cenchrus spica oblonga conglomerata. *echinatus*. LINN. Sp. 3. Gram. *locustis tumidioribus echinatis*. SCHEVCHZ. 77. In *Jamaica*. A.

ZEA. LINN. 926. Mays. TOVRN.

Monoph. Vtrique sexui calyx & folliculus bivalvis muticus. *Mares* in locustis bifloris laxè spicatis. *Feminae* foliis tectae; singulis stylus filiformis longissimus. Semina in receptaculo cylindrico ad dimidium demersa.

H. Zea. *Mays*. LINN. Sp. 1. ROY. 73. Frumentum indicum Mays dictum. BAYH. Pin. 25. a. *granis aureis*, β . *granis rubentibus*. In *America*. A.

COIX. LINN. 927.

Monoph. *Mares*. Locustae biflorae laxè spicatae, singulae calyce & folliculo mutico. *Feminae* ad basin priorum spicatae; fabrica *masculis* simili.

H. Coix seminibus ovatis. LINN. Sp. 1. ROYEN. 72. Lacryma Jobi. CLVS. Hist. 2. p. 216. ZANONI p. 136. T. 101. In *Indiis*. A.

H. Coix seminibus angulatis. *dactyloides*. LINN. Sp. 2. Gram. *dactylon maximum Americ.* PLUCK.

T. 190. f. 2. *Sesamo Indico perpetuo*. ZANON. T. 66. In America. P.

CYPEROIDEAE.

II. Stamina tria, & semen vnicum, stylo tricorni ad glumam vnicam.

CAREX. LINN. 928. *Cyperoides*. Carex

HALLER.

Monophyta vt plurimum. Vtriusque sexus flores spicati. Feminis semen triquetrum in nectario apico bidentato.

1. Flores masculi & feminini in eadem spica.

Carex spica paniculata. HALL. *Helv.* n. 12. p. 246. Carex *racem. composita*. LINN. *Nov. Sp. Cent.* I. n. 93. Gram. *cyperoides palustre*, spica longiori laxa. RAIL. *Cyperus longus inodorus sylvaticus*. C. BAVH. *Prodr.* n. 24. Ad oras stagnorum v. g. bey der Rasemühle. Seeburg. Landolfshausen.

Carex spicis alternis, latis, laxius in spicam dispositis. *Helv.* 244. Circa *Odorbrücke* in sylva *Hercynica*.

Carex spicis teretibus acutis, alternis, se contingentibus. HALL. *Helv.* 244. n. 5. Carex *brizoides*. LINN. *Nov. Sp. Cent.* I. n. 91. Gram. *cyperoides elegans*, spica composita asperiore. GOTT. p. 54. Circa *Göttingam* & in sylva *Hercynica*.

Carex spicis obesis subrotundis, in summo caule se alterne contingentibus. HALL. n. 7. p. cix. Carex *leporina*. LINN. *Sp.* 6. Gram. *cyperoides palustre*

luftre maior spica divisa. B. Pin. 6. Prope *Gottin-*
gam versus *Langlern*, in locis vdis.

Carex spica composita ex plurimis spicis tereti-
bus, acutis, squamae latae infidentibus. HALL.
Helv. p. 245. n. 8. *Gottinae* in pratis vdis. v. g.
auf dem Mafch.

Carex spicis brevibus sessilibus, echinatis in fum-
ma caule cohaerentibus. HALL. l. c. n. 9. *Carex*
vulpina. LINN. Sp. 7. Gr. *cyperoid*, *palustre maius*,
spica compacta. BAYH. Pin. 6. Ad sepes vdas.

Carex spicis brevibus, echinatis, sessilibus, di-
stinctis. HALL. 245. n. 10. *Carex muricata*. LINN.
Sp. 8. Gramen *nemorosum*, *spicis parvis asperis*. B.
Pin. 7. In nemoribus humentibus. v. g. *im Eu-*
lenloch.

Carex spicis parvis sessilibus in foliorum alis.
HALL. 243. n. 1. *Car. remota*. LINN. Sp. nov. Cent.
L n. 92. MORIS. III. p. 243. T. 12. f. 17. In
sylva Hercyniae.

2. Flores masculini & feminini in spicis distin-
ctis. *Cyperoides* HALLERI.

Carex (Cyperoides.) spicis ex luteo & spadiceo
variis, feminali suprema super spicam marem elata.
HALL. *Helv.* 236. n. 10. *Carex digitata*. LINN. Sp.
14. Gram. *caryophyllum*, *mont. spica varia*. C. B.
Prodr. 9. In nemoribus.

β. Gr. *caryophyllum*, *nemorosum*, *spica mul-*
tuplici. C. B. *Prodr.* n. 22. Num varietas?

Carex (Cyper.) capsulis subrotundis spicae vn-
dique rigentibus. HALL. *Helv.* n. 12. *Car. flava*.
LINN.

LINN. *Sp.* 12. *Gr. Cyper. aculeatum Germanicum*
f. minus. BAVH. *Pin.* 7. In paludibus.

Carex (Cyper.) spicis feminalibus breviter pedunculatis, & inter se & a mare remotissimis. HALL. *Helv.* 237. n. 13. *Cyperoides spicis parvis longe distantibus* I. R. H. In vdis.

Carex (Cyper.) spicis feminalibus longius pedunculatis, mutantibus, suprema marem erectam & gracilem contingente. HALL. *l. c.* n. 15. *Car. pallens.* LINN. *Sp.* 23. In paludibus.

Carex (Cyper.) spicis feminalibus breviter pedunculatis, rufis, marem obesam contingentibus. HALL. *l. c.* n. 14. *Cyperoides minus spicis densifloribus.* I. R. H. *Bey Robringen & ad Lützelengen.*

Carex spicis masculis pluribus, femineis pedunculatis, capsulis inflatis acuminatis. *vesicaria,* LINN. *Sp.* 29. α . *Cyperoides vesicarium, spicis viridantibus aut subfuscis.* SCHEVCHZ. 470. β . *Cyperoides minus praecox, spicis turgidis, teretibus, flavescens.* MORIS. *Hist.* III. p. 242. γ . *Gram. Cyperoides sylvatum tenuius spicatum.* SCHEVCHZ. 418. In vdis sylvaticis.

Carex (Cyper.) spicis feminalibus longis & asperis, ex longis pedunculis pendulis. HALL. *Helv.* 239. n. 25. *Carex. Pseudo-Cyperus.* LINN. *Spec.* 26. *Gr. cyperoides spica pendula brevior.* C. B. *Pin.* 6. In vdis.

Carex (Cyperoid.) latifolium, spicis feminalibus semimasculis, ex longis pedunculis pendulis. HALL. *Helv.*

Helv. 239. n. 26. Gr. *Cyperoides spica pendula longiore & angustiore.* C. B. in vdis.

Carex (*Cyperoid.*) spicis maribus obefis, femininis imis longe petiolatis, gracilibus. HALL. *Helv.* 31. Gramen *nigro luteum vernum.* I. B. II. p. 494. Carex. LINN. *Spec.* 28. α. In locis gramineis.

Carex (*Cyper.*) foliis latis & asperis, spicis plurimis, fere sessilibus, duabus androgynis. HALL. *Helv.* 29. Gr. *Cyperoid. spica rufa, f. caule triangulo.* BAVH. *Pin.* 6. Carex. LINN. *Spec.* 28. β. In vdis.

Carex. (*Cyperoides*) HALL. *Helv.* n. 32. Gram. *Cyperoid. caryophyllaeum*, elatius, granis tumidiorebus, spicis pedunculis tenuissimis appensis. MORIS, III. p. 243. n. 13. *Im Wald beym Hessen-dreisch.*

Carex. (*Cyperoides*) HALL. n. 33. *Cyperoides palustre*, spicis purpureo spadiceis, tenuibus pedunculis insidentibus. SCH. 467. In vdis.

Carex. (*Cyper.*) HALL. n. 34. *Cyperoides spicis sessilibus erectis & feminibus confertis compositis.* RAII III. p. 418. In paludibus versus sylvam supra *Geismar.*

Carex spicis remotis, femineis subsessilibus, capsulis hirtis. *hirta.* LINN. *Spec.* 13. *Cyperoides polytachyon lanuginosum.* RAI. I. R. H. In durnetis.

3. Flores masc. & fem. in plantis distinctis.

Carex spica simplici dioica. LINN. *Sp.* 1. *Vlua.* HALL. *Helv.* 242. In vdis circa *M. Brusterum.*

CYPE

CYPERVS. LINN. 61.

Floſculi androgyni in ſpica diſticha.

Cyperus minimus, panicula ſparſa flaveſcente.
HALL. *Helv.* 246. *Cyperus flaveſcens*. LINN. *Sp.*
10. In vdis.

SCIRPVVS. LINN. 62.

Spica florum imbricata. Semen villis calyce brevioribus munitum.

Scirpus caule aphylo, ſpica imbricata, terete
HALL. *Helv.* 248. n. 5. *Sc. paluſtris*. LINN. *Sp.* 2.
Scirpus equiſeti capitulo maior. C. B. In pratis
vdis.

Scirpus culmo vnifolio, ſpica bivalvi breviffima.
HALL. *Helv.* 249. *Sc. cespitofus*. LINN. *Sp.* 4. *Scirpus mont. capitulo brevior*. SCH. 363. T. 7. f. 18.
Circa Oderbrücke in ſylva Hercynica.

Scirpus teres panicula ex ſummo culmo ramoſa.
HALL. 2. p. 247. *Sc. paluſtris*. LINN. *Sp.* 7. *Juncus paluſtris maximus*. f. Schoenus. C. B. In ſtagnis.

Scirpus planifolius, panicula ramoſa foliis infidentibus.
HALL. n. 1. l. c. *Sc. ſylvaticus*. LINN. *Spec.* 22. *Gr. cyperoides miliaceum*. B. Pin. 6. *Am Raſenmühlen Waſſer*.

Scirpus planifolius, ſpicis obefis, foliis infidentibus.
HALL. *Gott.* 55. *Scirpus maritimus*. LINN. *Sp.* 20. *Cyperus rotundus, inodorus Germanicus*. B. Pin. 14. In falſis. v. g. *Bey Harſte, Saltz der Helden*.

LINAGROSTIS. HALL. p. 250: Eriophorum.

LINN. 63.

Priori simillima, a quo tantum differt villo seminis spica ipsa longiore.

Linagrostis foliis teretibus, spica subrotunda, villica. HALL. *Helv.* 250. n. 2: Eriophorum *vaginatum*. LINN. *Spec.* 1. Gram. *tomentosum alpinum* & *minus*. B. *Pin.* 5. In sylva *Hercyn.* & in vdis *Cellensibus*.

Linagrostis foliis planis, spicis multiplicibus. HALL. *l. c.* n. 1. Eriophorum *polystachion*. LINN. *Spec.* 2. Gr. *prat. tomentosum, panicula sparsa*. B. *Pin.* 4. In pratis vliginosis.

III. Stamina tria (duo), semen vnicum, stylo (bicorni) tricorni ad glumas sex.

SCHOENVS. LINN. 60.

Calyx locustae multiflorae biglumis. Panicula foliolo excepta. Semen aristis cinctum.

Schoenus flosculis fasciculatis. LINN. *Cliff.* 22. HALL. *Helv.* 251. *albus*. LINN. *Spec.* 9. Schoenus *locustis albicantibus erectis*. Gott. 57. In *Cellensi Ducatu*.

Schoenus spicis gemellis, obesis. HALL. *Gott. l. c.* n. 2. vbi vide characterem a priori satis diversum.

MARISCVS. HALL. *Gott.* 57.

Habitus prioris. Locustae biflorae. Stamina duo, stylus bicornis. Capsula nucleo foeta, aristis non cincta.

Maris-

Mariscus. HALL. *Helv.* p. 251. *Schoenus culmo teteti*, foliis margine dorsoque aculeatis. LINN. *Spec.* 1. *Cyperus longus inodorus Germanicus.* B. *Pin.* 14. *Im Denckerhäuser Teich.*

IV. Stamina sex, capsula tripartita, stylo tricorni ad glumas sex.

JUNCVS. LINN. 396.

Calyx communis biglumis. Capsula triquetra, unilocularis.

Juncus culmo aphylo, mollior, panicula compactiori. HALL. *Helv.* 252. n. 2. *Juncus conglomeratus*. LINN. *Spec.* 2. *Juncus laevis panicula non sparsa.* B. *Pin.* 12. In locis vdis.

Juncus culmo aphylo panicula magis sparsa. HALL. *l. c. n.* 1. *J. effusus.* LINN. *Sp.* 3. *Juncus laevis panicula sparsa maior.* B. *Pin.* 12. In vdis.

Juncus foliis angulatis, culmo dichotomo, floribus solitariis sessilibus, *bufonius*. LINN. *Spec.* 10. HALL. *n.* 8. *Gr. iuncum bufonium.* TABERN. p. 225. In locis inundatis.

Juncus foliis mollioribus, carinatis, panicula multiplici & ramosa. HALL. *Helv.* n. 10. *Juncus parvus cum pericarpis rotundis.* 1. B. *Ad Leinam.*

Juncus foliis duris, carinatis, floribus cartilagineis. HALL. *l. c. n.* 9. *Juncus squarrosus.* LINN. *Sp.* 7. *Gram. iuncum, foliis & spica iunci.* B. *Pin.* 5. In torfaceis circa Oderbrücke.

Juncus foliis articulatis, ramosae paniculae ramis semel ramosis. HALL. 255. n. 11. *Juncus foliis*

foliis nodoso-articularis. LINN. *Sp.* 8. *Gr. jun-
ceum*, folio articulado, aquaticum. β . idem cum
viriculis. γ . idem sylvaticum. BAVH. *Prodr.* 12.
In pratis. v. g. bey Herperhausen.

*Juncus foliis planis, pilosis, panicula in spicas
congesta*. HALL. *l. c.* 256. n. 14. *Juncus campe-
stris*. LINN. *Spec.* 14. *Gram. hirsutum, capitulis
Pyllyi*. C. B. *Pin.* 7. In sylvis.

*Juncus foliis planis, latis, panicula ramosa, spar-
sa*. HALL. n. 14. *Gram. hirsutum latifolium minus*.
C. B.

*Juncus foliis planis latissimis, panicula sparsa,
duplicato-ramosa*. HALL. 15. *Gram. Lazulæ ma-
ximum*. 1. B. II. p. 493. In sylvis circa Oderbrück.

*Juncus foliis planis, panicula umbellata, flori-
bus minimis*. HALL. 258. n. 20. *Gr. hirsutum an-
gustifolium minus*. C. B. *Prodr.* 16. In sylvis.

TRIGLOCHIN. LINN. 409. HALL.

Helv. 218.

*Junco similis staminibus, & calyce; siliculae au-
tem distinctae tres aut sex, totidemque tubae plu-
mosae.*

*Triglochin capsulis trilocularibus sublineari-
bus, paleis*. LINN. *Sp.* 1. *Triglochin fructu tena-
li*. HALL. *Helv.* 258. In pratis vdis.

*Triglochin capsulis sexlocularibus ovatis, mariti-
mis*. LINN. *Spec.* 2. *Gramen spicatum marit. ultio-
rum*. 108. 16. 17. In pratis salis prope Harste.

GRAMINA

ACORUS. LINN. 392. HALL. 259.

Clava cylindrica recta floribus sessilibus. Calyx & stamina ut in priori. Capsula trilocularis, tuba brevissima.

Acorus. *Calamus*. LINN. *Sp.* 1. *Mat. med.* 16. *Acorus vernus* f. *Calamus aromaticus officinarum*. C. B. *Pin.* 34. In fossis. In Ducatu Cellensi. Circa Eimbeccam. Noribergum.

IV. Monophytae, flores in receptaculo communi, staminibus tribus, tuba simplici, calyce triglumi.

TYPHA. LINN. 324.

In supremia parte clavae cylindrica flores mares, usinae feminae semine papposo.

Typha clava crassissima; cylindrica. HALL. *Helv.* 260. *latifolia*. LINN. *Sp.* 1. *Typha palustris maior*. BAVH. *Pin.* 20. In fossis suburbanis & stagna ad Denckerhausen.

SPARGANIUM. LINN. 925.

Flores in capitula collecti, masculini in superiore, femini stylo bicorni in inferiori parte plantae.

Sparganium foliis erectis triquetris. *erectum*. LINN. *Sp.* 1. HALL. *H.* 259. n. 1. α . *Sparganium ramosum*, β . non ramosum. BAVH. *Pin.* 15. In fossis v. g. circa Bovenden.

Sparganium foliis planis natantibus. *natans*. LINN. *Sp.* 2. HALL. n. 2. 260. *Sparganium minimum* C. B. RAIL. *Hist.* 1910. Circa Berolinum.

SAV-

SAURURVS.

Flóres in sulum digesti, ad singulas squamas stamina sex, & bacca tetrasperma, tubis quatuor.

HALL. *Gott.* 9. bacca monosperma, tubis tribus.

LINN. *gen.* 414.

H. *Saururus* foliis cordatis, petiolatis, spicis solitariis recurvis. LINN. *Sp.* 1. *Serpentaria repens*, floribus stamineis spicatis, *bryoniae nigrae* folia ampliore pingui, *Virgin.* REVK. T. 117. f. 4. In *Virginia.* Aro proxime affinis. P.

PETALOIDEAE.

CLASSIS. I.

PALMAE.

Caulis (*Caudex*) simplex. Fructificationes in spadice cum *Spatha*.

ARVM. LINN. 915. HALL. 261.

In *spatha* cucullata spadix clavae formae, superne nudus; interne germina, in baccas polyspermas abundantia; in medio antherae sessiles, interiectae duplici ordini papillarum cirrhosarum.

Arum foliis sagittatis, cucullo recto. HALL. 261.

Arum acaule, foliis hastatis integerrimis, spadice clavato, *maculatum*. LINN. *Sp.* 12. *Arum* vulg. *a. maculatum* β . non *maculatum*. B. *Pin.* 195. BLACKW. T. 228. In sylvis.

H. *Arum* acaule, foliis peltatis ovatis repandis, basi semibifidis, *Colocasia*. LINN. *Sp.* 6. *Colocasia*. CLVS. *Hist.* 2. p. 75. *Arum maximum*

Aegyptiarum. a. caule viridi, p. ex atro purpureo.
Pin. 193. In Hispania, Aegypto, Syria.

CALLA. LINN. 917. Provençalia. PET.

In spatha ovato cordata spadix digitiformis, vndique fere recta antheris sessilibus, & germinibus in baccas abeuntibus.

H. Calla foliis sagittato cordatis, spatha cucullata, spadice superne masculo. *Aethiopica. LINN. Sp. 1. Arum Aethiopicum, flore odorato Moschum redolente. COMMEL. Hort. 1. p. 95. T. 50. Antherae & germina sine certo ordine digesta mihi visa.*

PALMA. PHOENIX. LINN. 1089, LVDW. 908.

In spatha alari spadix ramosus. Corolla tripartita. Diphyta. Mas. Stamina sex. Fem. Bacca unilocularis, stylo simplici. Caudex apice frondosus.

H. Phoenix frondibus pinuatis; foliolis complicatis. *dactylifera. LINN. Sp. 1. p. 1188. Palma bortenfis. f. dactylifera. BLACKW. T. 202. & auctor. In India. P.*

MVSA. LINN. 1010. LVDW. 370.

Caudex apice frondosus. In spatha alari flores multi, corolla dipetala, staminibus sex, fructu carnoso longo, stylo cylindraceo. In flor. superioribus ovaria abortiunt.

H. Musa racemo simplicissimo. LINN. *H. Cliff. 467. Mus. Cliff. 1. T. 1. Spadice nutante. paradisiaca.*

ORCHIDEAE.

85

fiaca, Sp., I., TREW. Dec. | Pl. rar. II. & Commerces
lit. 1739, 40, 41. In India.

CLASSIS. II.

ORCHIDEAE.

Culmus simplex, foliis simplicibus vaginantibus
cinctus. Capsulae trivalves striatae flores sustinent.
Petalae quinque aut quatuor superiora in cucullum
conniventia, sextum inferius cavum, nectariferum,
interdum calcaratum *labelli* aut *nectarii* nomine,
quod gerit machinam duorum staminum, sub
qua stigma concavum. Vid. HALLER. *Comment.*
Soc. Sc. Gott. Tom. V.

CALCEOLVS. HALLER. I. c. *Cypripedium*.

LINN. 906.

Filamenta staminum duo distincta. Petala supe-
riora quatuor, inferius amplissimum, cavum. Radix
ramosa.

Calceolus. HALL. *Helv.* 276. Gott. 65. *Cypripe-*
dium radicibus fibrosis, foliis ovato lanceolatis equ-
limis. LINN. Sp. I. Calceolus *Marianus*. DOD. *Ma-*
rienschubz Circa *Petersen*.

EPIPACTIS. HALLER. *Serapias*. *Ophrys*.

LINN.

Machina staminum est scapus superne bilabiatus,
labium superius inferne duas antheras nudas, labio
inferiori incumbentes, fovet. Calcar nullum. Cap-
sula ovata. Radix, ut plurimum ramosa.

1. *Helleborine*. Labellum cavum, ex ovato
cordatum.

Epipactis (*Helleborine*) floribus obsolete carneis,
raris, labello obtuso. HALL. Enum. Helv. 275. Hel-
leborine palustris nostras. RAIL. bist. 1231. angu-
stifolia pratensis. B. Pin. 87. Serapias. LINN. Sp. 1.
Ad lacum prope Seeburg. Im Landolfshäuser
Sumpf.

Epipactis (*Helleborine*) amplexicaulis multiflora.
Helv. 275. Helleborine latifolia montana. C. B. α.
flore atrorubente β. flore pallido. C. B. Pin. 186.
Serapias. LINN. Spec. 1. α. In sylvis v. g. circa
Pless. Göttinger Wald.

Epipactis (*Helleborine*) latifolia floribus magnis
paucis albisque. Em. Helv. 276. Helleborine f.
Damasceniuth montanum latifolium. C. B. In sylvis
v. g. Göttinger Wald, circa Pless. &c.

β. Helleborine longifolia, floribus magnis, pau-
cis & albis. Helv. 275. Auf dem Brackshühl.

Epipactis (*Helleborine*) longifolia paucis pur-
pureis floribus. Helv. 274. Helleborine mont. an-
gustifolia purpurascens. B. Pin. 187. In sylvis v. g.
circa Pless.

Epipactis (*Helleborine*) bifolia radice ovali, cu-
tculo sparso. Comm. Soc. Sc. Gott. Tom. I. T. XIV.
p. 215. Circa Betersen.

2. *Ophrys*. Labellum longum, bicornis.

Epipactis (*Ophrys*) foliis ovatis. LINN. Cliff. 429.
ROYEN, 14. HALL. Helv. 277. *Ophrys ovata*.

LINN.

LINN. Sp. 5. *Ophrys bifolia*. B. Pin. 87. In pratis & silvis.

Epipactis (Ophrys) aphylla. HALL. Helv. 278. *Ophrys. Nidus avis*. LINN. Sp. 1. *Orchis abortiva fusca*. B. Pin. 86. *Nidus avis*. LOB. ic. 195. I. R. HALL. Omnes plantae partes lignei coloris. In locis vinbrosis nemorum.

ORCHIS. HALLER. *Orchis. Satyrium*.

LINN. 900. 901.

Machina staminea. In duobus loculis styli brevioris membrana clausis antherae duae ex filamentis convolutis factae. Calcat modo adest, modo deest.

Orchis radicibus subrotundis, cucullo purpureo, maculato, labello bifido. HALL. Gott. p. 61. Helv. n. 3. *Orchis* LINN. Sp. 10. d. *Orch. militaris major*. B. Pin. 81. I. M. Hengst.

Orchis radicibus subrotundis, brachiolis labelli brevioribus, spinula ex divisione medii trunculi eminente. HALL. Helv. 262. Gott. l. c. *Orchis* LINN. Sp. 10. a. *Orchis galea* & alis fere cinereis. I. n. II. p. 757. In *Suevia*, rarius supra *Bouenden*.

Orchis radicibus subrotundis, spica brevissima labello punctato, ferrato, mucico. HALL. Gott. p. 61. *Orchis* LINN. 10. γ. *Orchis militaris minor*. I. n. RVPP. III. T. VI. In gramineis ficcis.

n. *Orchis* radicibus subrotundis, bracteis ovario longioribus, albo flore. Gott. 62. *Orchis foetida, Sylvatica, alba, praecox, barba luteola*. RVPP. III. p. 297. Circa *Jenam*. In horto adest.

Orchis radicibus subrotundis, floribus obsoletis, cimicem olentibus, labello retro reflexo. *Enam.* *Helv.* 264. Orchis LINN. *Sp.* 6. Orchis *birci* odore minor. B. *Pin.* 82. In pratis circa *Dorsten* ad viam versus *Osterode*, circa *Hardeggen*, ad lacum *See-burgensem*.

1 Orchis radicibus subrotundis, cucullo sparso, labello ampliter trifido, crenato, emarginato. H. *Helv.* 265. Orch. LINN. *Sp.* 8. δ . Orchis *morio mas.* B. *Pin.* 81. *Satyrium mas.* BLACKW. T. 53. In sylvis.

1 Orchis radicibus subrotundis, cucullo clauso, lineato, labello ampliter trifido, crenato. HALL. *Helv.* 266. Orchis LINN. *Sp.* 8. α . Orchis *morio femina.* B. *Pin.* 82. In pratis siccis & caeduis sylvarum.

1 Orchis radicibus conicis, labello lingulato, longissimo, HALL. *Helv.* 266. Orchis *bifolia.* LINN. *Sp.* 2. Orchis *alba bifolia minor*, *calcare oblongo.* BAVH. *Pin.* 83. SEGV. *Veron.* T. 15. f. 10. In sylvis.

1 Orchis radicibus subrotundis, calcare nullo, labello latiusculo, bisulco, brachiolis brevibus. HALL. *Helv.* 268. *Ophrys insectifera.* LINN. *Sp.* 15. α . Orchis *muscam referens.* B. *Pin.* 83. Rarius in *Göttinger Wald bey der Brück*, circa *Pless*, &c.

1 Orchis radice subrotunda, calcare nullo, labello trifido. HALL. *Gott.* 63. *Helv.* 269. η . 20. *Ophrys* LINN. *Sp.* 11. *Herminium Fl. Lapp.* 317. *Monorchis montana minima*, flore obsoleto, vix conspicuo

MICHEL.

MICHEL. p. 30. T. 26. Monorchis. c. B. In pratis inter Dorsten & Osterode. Ad viam inter Hessendreisbach & Roßbrünn.

Orchis radicibus palmatis, calcare ovario longiore, labello aequaliter trifido, unicolore. HALL. Helv. 271. Orchis conopsea. LINN. 13. Orchis palmata minor, calcaribus oblongis. C. B. Pin. 85. In pratis sylvarum.

Orchis radicibus palmatis, calcare crasso, foliis maculatis; caule fistuloso. HALL. Gott. 63. Helv. n. 27. Orchis latifolia. LINN. Sp. 11. Orchis palmata pratensis a. maculata. β. non maculata. γ. latifolia. &c. C. B. In pratis ubique.

Orchis radicibus palmatis, calcare crasso, labello profunde trifido, serrato, caule solido. HALL. Gott. 64. Orchis maculata. LINN. Sp. 12. Orchis palmata, montana, maculata. C. B. Pin. 86. In sylvis montanis.

Orchis radicibus multis cylindricis, calcare brevissimo. HALL. Helv. 270. Satyrium albidum. LINN. Sp. 4. Pseudorchis alpina, flore herbaceo MICHEL, gen. 32. T. 26. In summo M. Brütten.

CLASSIS. III.

LILIACEAE.

Calyx nullus, cuius loco in variis spatha membranacea, marcescens. Corolla tripetala, aut bis tripetala, aut monopetala limbo bis trifido. Stamina tria aut sex. Stylus simplex, stigmate trifido, rarius styli

styli tres. Fructus trigonus trilocularis, trivalvis. Folia simplicia, integra, nervis parallelis. Caulis aut nullus, aut foliis vaginantibus cinctus, simplex, herbaceus. Radix bulbosa, vel tuberosa, perennis.

I. Fructu sub flore posito,

a. flore tristemone. His fere omnibus folia ensata, & spathae sub flore.

COMMELINA. LINN. 58.

Corolla bis tripetala, tria petala calycem fermentientia. Filamentis tribus corpuscula cruciformia insistant.

Commelina corollis inaequalibus, foliis ovato-lanceolatis acutis, caule procumbente glabro. *communis*. LINN. *Sp.* 1. *Commelina procumbens annua, saponariae folio*. DILL. *Elth.* T. 78. f. 89. In America. A.

Commelina corollis inaequalibus, foliis lanceolatis glabris obtusis, caule repente. *africana*. LINN. *Sp.* 2. *Commelina procumbens flore lutea*. ROYEN. 538. HALL. *Gott.* 66. In Aethiopia.

Commelina corollis aequalibus, foliis ovato lanceolatis subciliatis. *tuberosa*. LINN. *Sp.* 5. *Commelina radice Anacampserotis*. DILL. *Elth.* T. 79. f. 90. In Mexico.

TRADESCANTIA. LINN.

Spatha foliaceae flores continens. Calyx trifolius. Petala tria. Stamina villosa.

H. Tradescantia. *Virginiana*. LINN. *Sp.* 1. *Ephem. merum*. TERN. 193. In Virginia.

BRO-

BROMELIA. LINN. 356. Ananas.

TOURN.

Flores in nostra specie in conum foliis coronatum collecti, spathis intermistis. Calyx trigonus. Petala tria. Bacca umbilicata.

H. Bromelia foliis ciliato spinosis mucronatis, spica compola. *Ananas*. LINN. *Sp.* 1. *Ananas aculeatus*, *fructu pyramidato, carne aenea*. 1. R. H. 653. In *America meridionali*.

HAEMANTHVS. LINN. 394.

Involucrum sexfolium umbelliferum. Corolla profunde sexpartita. Bacca subrotunda.

H. Haemanthus foliis linguiformibus planis, coccineus. LINN. *Sp.* 1. Haemanthus *Africanus* COMM. H. II. p. 127. T. 64. Ad C. bon. Spei.

H. Haemanthus foliis lanceolatis, *puniceus*. LINN. *Sp.* 2. Haemanthus *colchici folijs, perianthia hex- baceo*. DILL. *Eleb.* 167. f. 149. In *Africa*.

GLADIOLVS. LINN. 55.

HALL. 279.

Corolla sexpartita, bilabiata. Stamina adscendentia; stigmata undulata.

H. Gladiolus foliis ensiformibus, floribus distantibus. *communis*. LINN. *Sp.* 1. HALL. *Helv.* 279. Gladiolus *floribus uno versu dispositis* C. B. Pin. 41. In *Marchia & Europa australi*.

IRIS. LINN. 57. HALL. 279. *Xiphium*.
Silyrinchium.

Corolla sexpartita, petala tria reflexa. Stigmata
lata, petaloidea, antheras deprimentia.

1. Petalis reflexis barbatis.

H. Iris corollis barbatis, caule foliis longiore uni-
floro. *Susiana*. LINN. *Sp.* 1. *Iris Chalcidonica latifolia*.
I. B. II, p. 721. *Iris Susiana, flore maximo ex atro-*
bo nigricante. C. B. *Pin.* 31. In *Oriente*.

H. Iris corollis barbatis caule foliis longiore,
multifloro. *Germanica*. LINN. *Spec.* 2. *Iris nostras*
hortensis. BLACKW. *T.* 69. *Iris vulg. Germanica f.*
sylvestris C. BAVH. *Pin.* 30. a. petalis reflexis vio-
laceis (coeruleis), erectis aequae coeruleis (albidis).
In *Germaniae editis*.

H. Iris petalis reflexis fuscopictis, erectis luteis.
Iris lutea variegata. LOBEL. *hist.* 34. *Iris* LINN.
Sp. 4. In *Pannonia*. Num Varietas?

H. Iris corollis barbatis, caule foliis brevioris uni-
floro. *pumila*. LINN. *Sp.* 6. *Iris humilis latifolia*
minor, flore purpureo. I. R. H. 361. In *Austria*.
Pannonia, Sibiria.

2. Petalis reflexis imberbibus.

1. *Iris corollis imberbibus, petalis interioribus sti-*
gmate minoribus, foliis ensiformibus. Pseudacorus.
LINN. *Sp.* 7. *Iris imberbis lutea, nervo folii emi-*
nente. HALL. *Helv.* 281. *Acorus adulterinus*,
BLACKW. *T.* 361. C. BAVH. *Pin.* 34. *Gelbe Wasser-*
Lilien. Ad ripas fossarum.

H. Iris corollis imberbibus, petalis internis longitudinaline stigmati, foliis ensiformibus. *foetidissima*. LINN. 8. Iris *foetidissima* f. *Xyris*. TOURN. 360. Spatula *foetida* f. *Xyris*. BLACKW. T. 138. In *Anglia, Gallia, Hetruria*.

H. Iris corollis imberbibus, germinibus trigonis, caule tereti, foliis linearibus. LINN. Sp. 9. HALL. *Helb.* 280. Iris *angustifolia non foetida altior*. C. BAY. Pin. 32. In *Marchia*. Nobis hortensis.

β. Iris sylvestris peramoena. LÖB. *ic.* 69.

H. Iris corollis imberbibus, germinibus sexangulis, caule ancipiti, foliis linearibus. *graminea*. LINN. Sp. 13. ROYEN. 18. In *Austria*.

H. Iris corollis imberbibus, foliis subulato canaliculatis caule brevioribus. *Xiphium* LINN. *Spec.* 16. Iris *bulbosa latifolia, caule donata*. C. B. Pin. 38. In *Hispania*.

SISYRINCHIVM. LINN. 908.

Spatha diphylla, universalis. Antherae stylo ad basin stigmati affixae.

H. Sisyrinchium caule foliisque ancipitibus. LINN. Sp. 4. Bermudianā *iridis folio, fibrosa radice*. DILL. *Elth.* T. 41. f. 48. In *Bermudis*.

IXIA. LINN. 54.

Petala sex aequalia. Stigmata convoluta. Stamina libera. Spathae vagae.

H. Ixia foliis ensiformibus, floribus remotis. LINN. *Spec.* 2. Bermudiana *florē aurantio maculata*.

ca. HALL. *Gott.* 68. ROLOF. *hort.* Kraus. p. 23.
ic. In India.

CROCVS. LINN. 53.

Corolla sexpartita, aequalis, ex spatha monophylla. Stigmata convoluta ferrata.

H. *Crocus* spatha univalvi radicali, corollae rubro longissimo. *sativus*. LINN. *Sp.* I. BLACKW. *L.* 144.
Crocus α . *vernus*, β . *autumnalis*. HALL. *Helv.* 281.
variorum colorum. Sponte in alpinis Helveticis, Pyrenaeis.

β . Flore hexastemone, ut plurimum spathaceae.

COLCHICVM. LINN. 415.

Habitus fere Croci. Tubus floris radicans. Styli tres, stigmata canaliculata.

Colchicum foliis planis lanceolatis erectis, *autumnale*. LINN. *Sp.* I. Colchicum. HALL. *Helv.* 283.
 α . purpureum β . album γ . ex albo & purpureo varium. Ubique in pratis; germen intra radicem fovet, dum autumnis floret, altero vere prorupturum & in capsulam inflatam abiturum.

GALANTHVS. LINN. 362.

Petala tria maiora parva, tria dimidio minora cylindrum formantia. Stigma simplex.

H. *Galanthus* *nivalis*. LINN. *Sp.* I. REY. 33.
Leucoium bulbosum trifolium minus. C. B. *Pin.* 156.
 In hortis; spontanea fere in pomario arcis Castellanburg. Patria alpina.

LEUCOIVM. LINN. 363.

Corolla campaniformis, petalis apice incrassatis.
Stigma fetaceum, simplex.

Leucoium spatha uniflora, stylo clavato. *vernum*.
LINN. *Spec.* 1. Leucoium petalis aequalibus. HALL.
Holo. 284. Narcisso-Leucoium *vulgare*. I. R. H.
In sylvis. v. g. *über Güssmar. im Eulenloch.*

NARCISSVS. LINN. 364. HALL. 284.

Ex tubo longo petala sex plana, inserta nectario cylindraceo monophyllo. Stigma breviter trifidum.

H. Narcissus spatha uniflora, nectarii limbo rotato brevissimo, (foliis ensiformibus). *poeticus*. LINN.
Sp. 1. Narcissus *albus circulo purpureo & croceo*.
C. B. *Pin.* 48. I. R. H. In *Helvetia. Gallia.*

H. Narcissus spatha uniflora, nectarii limbo campanulato erecto petalo aequali. *Pseudo-Narcissus*.
LINN. *Sp.* 2. Narcissus *sylvestris pallidus, calyce luteo*. C. B. *Pin.* 52. In *Angliae, Italiae* nemo-
ribus.

H. Narcissus spatha multiflora, nectario campanulato brevi, foliis subulatis. *Jonquilla*. LINN. *Sp.*
5. Narcissus *iuncifolius luteus minor* C. B. *Pin.* 51.
Sponte Monspeli.

AMARYLLIS. LINN. 367.

Petala sex lanceolata. Stigma trifidum tenue
Corolla in aliis aequalis, in aliis inequalis, bila-
biata.

H. Amaryllis spatha uniflora, corolla aequali, pistillo declinato. *Atamasca*. LINN. Sp. 2. Liliomarcissus ind. flore albo exterius rubente. 1. R. H. In Virginia.

H. Amaryllis spatha uniflora, corolla inaequali, genitalibus, declinatis, *formosissima*. LINN. Sp. 3. Lilio-Narcissus *Jacobaeus*, flore sanguineo nutante. DILL. Elth. T. 162. f. 196. In America meridionali.

H. Amaryllis spatha multiflora, corollis revolutis, genitalibus strictis. *Sarniensis*. LINN. Sp. 5. Narcissus *Japonicus*, rutilo flore. 1. R. H. Guernsey-Lilly. DOUGLASS in proprio libello. Sponte in Japonia, nunc in insula *Guernsey*.

ALOE. LINN. 389.

Corolla monopetala in fundo nectarifera ventricosa, aut limbo parvo sexfido, aut profundius sexpartita, modo simplex aequalis, modo difformis. Stamina receptaculo inserta, stigma modo simplex, modo trifidum. Capsula oblonga, trifidula.

H. Aloe floribus pedunculatis cernuis corymbosis, subcylindricis *perfoliata*. LINN. Sp. 1. Aloe *foliis caulinis dentatis, amplexicaulibus vaginantibus*. Cliff. 132. Ad hanc speciem tanquam varietates refert LINNAEUS sequentes, quae sunt.

α. Aloe Afr. caulescens, foliis glaucis brevissimis. COMMEL. Prael. T. 22. *varior*. T. 45.

β. Aloe

3. Aloe Africana caulescens, foliis glaucis caulem amplexantibus. COMMEL. *H. Amst.* II. T. 14.

HALL. *Gott.* 70.

7. Aloe Afr. foliis glaucis margine & dorso parte superiore spinosis, flore rubro. COMMEL. *Prael.*

24. *H. Amstel.* II. T. 12.

8. Aloe Afr. caulescens, foliis glaucis, longioribus, in apice dorso etiam spinosis. HALL. *Gott.* 72.

9. Aloe Afr. foliis glaucis, margine & dorso integro spinosis. COMMEL. *Prael.* T. 20. *H. Amst.*

II. T. 13, *Gott.* 71.

10. Aloe caulescens foliis minus glaucis, caulem amplexantibus. COMMEL. *Prael.* 18.

11. Aloe Afr. caulescens, foliis spinosis, maculis ab utraque parte albicantibus notata. COMM. *H.*

Amst. II. T. 5. HALL. *Gott.* 72.

12. Aloe Africana maculata spinosa maior. HALL. *Elb.* T. 15.

13. Aloe Africana humilis, spinis & verrucis obsita, COMM. *Prael.* T. 16. *Pl. rar.* T. 46. Conveniunt quidem stigmate simplici, sed differunt tubo limbo & colore floris.

14. Aloe floribus pedunculatis cernuis racemosis prismaticis, ore patulo. *variegata.* LINN. *Sp.* 2. Aloe

foliis canaliculatis, trifariisq. imbricatis radicatis erectis: angulis cartilagineis ternis. Hort. Cliff. 132.

Aloe Afr. humilis foliis ex albo viridi variegatis. COMM. *Prael.* 79. T. 28. *Pl. rar.* T. 27.

H. Aloe floribus pedunculatis ovatis cylindricis curvis *distichis*. LINN. Sp. 3. Aloe foliis *lanceolatis* formibus *patulis distichis*. Cliff. 132.

α. Aloe Afr. flore rubro, folio triangulari, marginali ab utraque parte albicantibus notato. COMM. Amst. II. f. 8. HALL. Gott. 75.

β. Aloe Africana sessilis, foliis cuneatis verrucosis. DILL. Elth. T. 16.

γ. Aloe Afr. foliis planis conjugatis, carinatis, verrucosis, caule & flore corallii colore. POIR. II. p. 131. MART. Cent. II. T. 11.

δ. Aloe Afr. arborescens, montana, non spinosa, folio longissimo, plicatili, flore rubro. COMM. H. Amst. II. T. 3.

H. Aloe floribus sessilibus horizontalibus infundibuliformibus aequalibus, limbo revolutus *hyacinthoides*. LINN. Spec. 4. Aloe foliis *lanceolatis planis rectis radicatis*. Cliff. 32.

α. Aloe Ceylanica pumila, foliis variegatis. COMM. Amst. II. T. 21.

β. Aloe Guineensis, radice geniculata, folio variegato. COMM. Prael. T. 33. H. Amst. II. f. 20.

H. Aloe floribus sessilibus ovatis crenatis, segmentis interioribus connatis. *spiralis*. LINN. Sp. 5. Aloe foliis *ovatis acuminatis caulinis quinquefarariam imbricatis*. Cliff. 132. Aloe Afr. erecta, folio *retundo, acuminato*. COMMEL. Prael. 32. HALL. Gott. 73.

H. Aloe floribus sessilibus triquetris, bilabiatis, labio inferiori revoluta, *retusa*. LINN. Sp. 6. Aloe
foliis

foliis rhomboidibus ventricosis quinquefariam positis radicatis triquetris plano-extantibus. *Cliff.* 132. Aloe Africana, brevissimo crassissimoque folio, flore viridi. *COMM. Amst. II. T. 6.*

H. Aloe floribus sessilibus infundibuliformibus bilabiatis, lacinjis quinque revolutis: summa erecta. viscosa. *LINN. Sp. 7.* Aloe foliis canaliculatis imbricatis, caulinis apice reflexo-patulis. *Cliff.* 132. Aloe Afr. erecta triangularis & triangulari folio viscosa. *COMM. Proel. T. 31. DILL. Elsb. T. 13. f. 13.*

H. Aloe floribus sessilibus bilabiatis: labio superiore erecto, inferiore patente. *pumila, LINN. Sp. 8.* Aloe foliis ovato subulatis, acuminatis tuberculis cartilagineis utraque dispersis. *Cliff.* 131.

a. Aloe Afr. margaritifera minor. *COMM. Amst. II. T. 14. DILL. Elsb. T. 16. f. 17.*

β. Aloe Afr. margaritifera minima. *DILL. Elsb. T. 16. f. 18.*

γ. Aloe Afr. humilis anethoides. *COMM. Proel. T. 27.*

δ. Aloe Afr. minima atrochorda, spinis herbaceis numerosis ornata. *BOERH. II. T. 131.*

Conveniunt petalis sex fere distinctis congruentibus, stigmate trisido.

H. Aloe floribus sessilibus reflexis imbricatis prismatica. *Varia. LINN. Sp. 9.* Aloe foliis linearibus radicalibus membranaceis. *Cliff.* 133. Aloe Afr. folio triangulari longissimo & angustissimo, floribus luteis foetidis. *COMM. II. T. 15.*

AGAVE. LINN. 390.

Corolla campaniformis. Stamina flore duplo longiora. Stigma trigonum.

H. Agave foliis dentato-spinosis, scapo ramoso. *Americana*. LINN. Sp. 1. Aloe falso in oblongum aculeum abeunte. C. B. Pin. 286. Aloe Americana. HALL. Gott. 69. TREW. Comm. litt. 1744. T. 3. In America.

II. Fructu intra florem posito.

a. solitariis nudis floribus.

I. floribus hexapetalis, aut ad basin divisis, campanulatis.

YUCCA. LINN. 388. Cordylina.

ROYEN.

Corolla vngribus cohaerens. Stylus nullus. Filamenta staminum superne crassiora.

H. Yucca foliis integerrimis. *gloriosa*. LINN. Sp. 1. Yucca foliis. ALLEN. BAYH. Pin. 91A In Canada, Peru.

H. Yucca foliis crenatis mutantibus. *dracunculis*. LINN. Spec. 1. Yucca Draconis folio serrata. BILL. Eltham. T. 324. f. 417. Cordylina foliis pungentibus crenatis. ROY. Lugd. 22. In America.

HEMEROCALLIS. LINN. 391.

Ex tubo brevi corolla fere infundibuliformis. Stamina declinata. Stylus filiformis.

H. Hemerocallis scapo ramoso, corollis monopetalis. *Lilioasphodelus*. LINN. Sp. 1. a. Lilioasphodelus.

Lilium luteum. CLVS. *Hist.* 1. p. 337. In *Sibiria Italia*.

β. *Lilium asphodelus phoeniceum*. CLVS. 1. c. LOB. 16. 23. In *India*.

LILIVM. LINN. 371.

Petala reflexa, ad basin singulorum linea nectarifera. Stylus stigmatibus triangularibus.

H. *Lilium* foliis sparsis, corollis campanulatis, intus glabris. *candidum*. LINN. *Sp.* 1. *Lilium album*. C. B. *Pin.* 76. BLACKW. *T.* 11. & offic. In *Parthia*.

H. *Lilium* foliis sparsis, corollis campanulatis erectis, intus scabris. *bulbiferum*. LINN. *Sp.* 2. *Lilium purpureo-croceum*. C. B. *Pin.* 77. In *Sibiria Austria Italia*.

Lilium foliis verticillatis, floribus reflexis corollis revolutis. *Martagon*. LINN. *Sp.* 5. HALL. *Flecht.* 289. *Lilium montanum* f. *syloestrum*. In *sylvis*.

H. *Lilium* foliis verticillatis, floribus reflexis, corollis campanulatis. *Canadense*. LINN. *Sp.* 6. *Lilium angustifolium*, flore flavo maculis nigris distincto. I. R. H. 371. In *Canada*.

FRITILLARIA. LINN. 372. Corona Imperialis TOURN. 197.

Fovea nectarifera ad basin petalorum. Stigmata triplex. Flores penduli.

H. *Fritillaria* foliis omnibus alternis. *Melagris*. LINN. *Sp.* *Fritill.* & *fellorum*. *dis. florent.* *Chiff.* 219. G 3 Fritil-

Fritillaria *ca. alba*. *P. purpureo variegata*. C. B. Pin. 64. In *Helvetia*. *Gallia*.

H. Fritillaria racemo comoso inferne nudo, foliis integerrimis. imperialis. LINN. Sp. 1. Corona imperialis. I. R. H. C. B. & auctor. In Oriente.

ERYTHRONIUM. LINN. 375

Ad basin trium petalorum interiorum tubercula
Sylus stigmatum triplici.

H. Erythronium. Dens Canis. LINN. Sp. 1. Dens Canis BAVH. Pin. 87. In Italia. Sibiria. Virginia.

TULIPA. LINN. 376

I. Stylus nullus. Stigma triangulare, trilobum, angulatis. Capula trigono-teres.

H. Tulipa flore erecto, foliis ovato-lanceolatis. Asiatica. LINN. Sp. 2. Tulipa B. V. M. & auctorum. In Oriente.

H. Tulipa flore suberectis, foliis lanceolatis. Vestris. LINN. Sp. 1. Tulipa minor lutea italica gallica. B. Pin. 63. Montpellier. In pomario Cattleburgensis praefecturae quasi spontanea.

2. floribus monopetalis.

HYACINTHUS. LINN. 381

Corolla monopetala, in aliis sexpartita, in aliis limbo breviter sexfido. Ad apicem germinis tres pori nectariferi.

H. Hyacinthus corollis campanulatis, sexpartitis apice revolutis. non scriptus. LINN. Sp. 1. Hyacinthus

sinthus oblongo flore coeruleus maior. C. B. Pin. 43.
In Anglia, Gallia. Helvetia.

H. Hyacinthus corollis infundibuliformibus semis-
sexisdis basi ventricosis. *orientalis.* LINN. Sp. 5.
Hyacinthus orientalis, C. B. Pin. 44. DOD. Pempt.
216. Varias. flore pleno in hordis. In Asia, Africa.

H. Hyacinthus corollis ovatis. *Muscari.* LINN.
Sp. 6. Hyac. *racemosus moschatum.* C. B. Pin. 43.
Muscari obsoletiori flore. CLVS. Hist. I. p. 178.
In Syria.

H. Hyacinthus corollis subovatis. *monstrosus.*
LINN. Spec. 7. Hyac. *Sannehus panicula comosa.*
COL. Ecphr. 2. p. 10. T. 12.

H. Hyacinthus corollis angulato-cylindricis, sum-
mis sterilibus longius pedunculatis. *comosus.* LINN.
Spec. 8. *Muscari latifolium, summis floribus in lon-
gis petiolis erectis.* HALL. Helv. 285. n. 1. Hyac.
max. botryoides, coma caerulea. I. B. II. p. 574.

POLYANTHES. LINN. 384.

Corolla infundibuliformis. Stamina ex faucibus
corollae orta.

H. Polyanthes floribus alternis. *tuberosa.* LINN.
Sp. 1. Cliff. 127. *Hyacinthus indic. tuberosa radice.*
CLVS. Hist. I. p. 176. *Tuberosa hortulanorum.* In
India.

POLYGONATVM. HALL. 286. Convallariae
LINN. spec.

Corollae campanulae limbus breviter sexfidus.
Bacca globosa immatura maculata.

1. *Polygonatum bifolium*, scapo florigero apophyllo.
HALL. *Helv.* 286. *Convallaria majalis* LINN. *Sp.*
1. *Lilium convallium*, C. B. *Pin.* 304. BLACKW.
T. 70. *Mayblume*. In sylvis.

1. *Polygonatum foliis alternis, pedunculis pendulis*
multifloris. HALL. *l. c.* 287. n. 4. *Convallaria multi-*
flora. LINN. *Spec.* 4. *Polygonatum maximum lati-*
folium, C. B. *Pin.* 303. BLACKW. T. 251. vulgo *Si-*
gillum Salomonis. In dumetis.

1. *Polygonatum foliis alternis, pedunculis pendu-*
lis unifloris. EN. *Helv.* 287. *Convallaria Polygo-*
natum. LINN. *Sp.* 3. *Polygonatum flore odoro-*
C. B. *l. c.* In praecipitiis circa arcem *Pless*.

1. *Polygonatum foliis verticillatis*. HALL. *l. c.* 288.
Convallaria verticillata. LINN. *Sp.* 2. *Polygonatum*
angustifolium non ramosum, C. B. *Pin.* 303. In
sylvis.

VNIFOLIUM. HALL. *Helv.* 288. *Convallariae*
Sp. LINN.

Priori proximo vicinum. Flores autem quadrifidus
Stamina quatuor. Fructus bilocularis.

1. *Vnifolium*. HALL. *l. c.* 288. *Convallaria foliis*
cordatis. LINN. *Sp.* 8. *Lilium convallium minus*
B. *Pin.* 304. In sylvis.

ASPARAGVS. LINN. 282.

Petala vnguibus coercentis. Stamina petalis in-
serta. Bacca globosa umbilicata.

n. Aspa-

H. *Asparagus* caule herbaceo erecto, foliis setaceis, stipulis paribus. *officinalis*. LINN. *Sp.* 1. *Asparagus sativus*. C. B. *Pin.* 489. BLACKW. *T.* 332. In arenosis. v. g. circa *Berolinum*: ad *Rhenum*.

H. *Asparagus* aculeis solitariis, ramis flexuosis, foliis fasciculatis angulatis muticis deciduis. *alba*. LINN. *Sp.* 6. *Asparagus aculeatus*, spinis horridus. C. B. *Pin.* 490. In *Hispania*.

3. floribus nudis planis.

ASPHODELVS. LINN. 379.

Corolla sexpartita. Valvulae sex basi petali insertae circa germen. Filamenta staminum ex valvulis orta.

H. *Asphodelus* caule folioso, foliis triquetris fistulosis. *luteus*. LINN. *Spec.* 1. *Asphodelus luteus*. C. B. *Pin.* 28. Hasta regia. BLACKW. *T.* 233. In *Sicilia*.

H. *Asphodelus* caule nudo, foliis strictis subulatis striatis subfistulosis. *fistulosus*. LINN. *Sp.* 2. *Asphodelus minor*. CLVS. *Hist.* 1. p. 197. In *Europa australi*. *Syria*: *Creta*.

ANTHERICVM. LINN. 380. Phalangium.

HAL. TOWN.

Corolla hexapetala. Filamenta staminum subulata. Stigma obtusum.

H. *Anthericum* foliis carnosis subulatis plantisculis. *alcoides*. LINN. *Sp.* 5. *Phalangium capense* *sedile* foliis *alcoiformibus* *pulposis*. DILL. *Esh.* *T.*

232. f. 300. Stamina plumosa. Ad Caput binas
Speci.

h. Anthericum foliis carnosiss teretibus, caule
fruticoso. *frutescens*. LINN. *Spec.* 4. Phalangium
Afr. foliis cepacis, floribus spicatis aureis. BOERH.
Ind. Pl. II. p. 133. Phalangium *Capense caulescens,*
foliis cepacis succosis. DILL. *T. 231. f. 298.* Sta-
mina barbata.

Anthericum foliis planis, scapo simplicissimo.
Liliago, LINN. *Sp.* 3. Phalangium *radicibus tere-*
tibus, caule nudo, foliis gramineis, petiolis flo-
rum non ramosis. HALL. *Hélu. 292. n. 11.* Phalan-
gium parvum flore non ramosum. C. B. *Pin. 29.* Fil-
amenta glabra. In asperis lapideis circa arcem
Plesi.

Anthericum foliis planis, scapo ramoso, corol-
lis planis, *ramosum*. LINN. *Spec.* 2. Phalangium
priori simile, petiolis ramosis. HALL. *l. c. n. 2.* Pha-
langium *parvo flore ramosum*. C. B. *l. c.* Flores
minores & viridibus lineis notati. In M. *Alten-*
stallberg.

β. Ornithogalum *spicis florum longissimis ra-*
mosis. GMEL. *Fl. Sib. T. I. Tab. 8.* Flos totus
albus.

ORNITHOGALVM. LINN. 377.

Corolla hexapetala. Filamenta staminum plana,
in aliis omnia, in aliis alterna apice trifido.

Ornithogalum scapo anguloso diphylo, peduncu-
lis umbellatis simplicibus, filamentis omnibus subu-
latis. *luteum*. LINN. *Sp.* 1. Phalangium *radice sub-*
rotun-

secunda, foliis 2, floribus umbellatis. HALL. *Hist.* 293. n. 5. *Ornithogalum luteum*. G. B. *Pin.* 1. In agris circa *Göttingam*. Omnis habitus hanc plantam hic retinet, etiam stamina a definitione generica differant.

G. *Ornithogalum floribus corymbosis*, pedunculis scapo altioribus, filamentis alternis emarginatis, *umbellatum*. LINN. *Sp.* 9. *Ornithog. pedunculis florum imis supremos superantibus*. HALL. *Helv.* 294. *Ornithog. umbellatum medium angustifolium*. G. B. *Pin.* 294. Sponte in *Helvetia* in agris.

H. *Ornithogalum floribus corymbosis*, pedunculis scapo humilioribus, filamentis emarginatis, *arabicum*. LINN. *Sp.* 8. *Ornithogalum arabicum*. C. B. *Hist.* II. p. 186. In *Aegypto*. *Arabia*.

G. floribus aggregatis vagina communi exceptis.

ALLIVM. HALL. LINN. 370. Ceba.

Porrum. Moly.

Petala concavo-erecta. Stigma acutum. Capsula brevis, trigona. Spatha bifolia.

I. Staminiibus alterne trifidis.

H. *Allium staminiibus alterne trifidis*, foliis gramineis, capite bulbifero, nucleis in bulbum connatis. HALL. *opusc. bot.* 331. *de allio*. n. 1. *Allium sativum*. LINN. *Sp.* 10. C. BAVH. *Pin.* 73. In *Sicilia*.

H. *Allium staminiibus alterne trifidis*, capite bulbifero, scapo ante maturitatem contorto. HALL. *All.* n. 2. *Allium Scorodoprasum*. LINN. *Sp.* 11. *B.*

ALLI

Allium genus *Opoteptodon* differt quibusdam. *Hist.* II. p. 539.

1. *Allium* staminibus alterne trifidis, capite bulbifero, foliis gramineis leviter serratis. HALL. *All.* n. 3. RVPP. *Jen. T.* 2. f. 2. *Allium arenarium*. LINN. *Sp.* 3. *Allium mont. latifolium*. CLVS. *Pann.* p. 212. In M. Kischäuser.

2. *Allium* staminibus alterne trifidis, foliis fistulosis, capite sphaerico, bulbifero, radice multiplici. HALL. *All.* n. 4. *Allium vineale*. LINN. *Sp.* 18. *Allium sylvestre*. C. B. *Pin.* 74. *Porrum sylv. vinearum*. C. B. 72. In agris circa Göttingam.

3. *Allium* staminibus alterne trifidis, foliis gramineis, floribus sphaerice congestis, radice scabulifera. HALL. *All.* 5. *Allium Ampeloprasum*. LINN. *Sp.* 1. *Allium sphaerico capite, folio latiore*. C. B. *Pin.* 74. In Minorca. In ins. Holms.

4. *Allium* staminibus alterne trifidis, foliis latissimis, capite sphaerico, caule ancipiti. HALL. *All.* n. 6. *Allium nutans*. LINN. *Sp.* 20. *Porrum scopulnudo, ancipiti, antequam floruerit, nutante, foliis ensiformibus, hinc paulo convexioribus*. GMEL. *Fl. Sib. T.* I. p. 55. T. 12. In Sibiria.

5. *Allium* staminibus alterne trifidis, foliis gramineis floribus sphaerice congestis, radice tunicata, cauli circumnata. HALL. *All.* n. 7. *Allium Porrum*. LINN. *Sp.* 2. *Porrum commune*. C. B. *Pin.* 72.

6. *Allium* staminibus alterne trifidis praelongis, capite sphaerico, atropurpureo. HALL. *All.* n. 8.

An

An. *Allium sphaerocephalum*. LINN. Sp. 14. *Moly atropurpureum*. SWEERT. T. 60. In Sibiria.

H. *Allium staminibus alterne trifidis*, caule ad terram ventricoso. HALL. *All. n.* 12. *Allium scapo nudo inferne ventricoso longiore, foliis teretibus*. (Cepa.) LINN. Sp. 25. *Cepa vulgaris*. C. B. Pin. 71.

2. *Staminibus omnibus simplicibus, subulatis.*

H. *Allium staminibus simplicibus*, caule medio ventricoso. HALL. *All. n.* 13. *Allium scapo nudo, folia adaequante, foliis teretibus ventricosis*. (*fistulosum*.) LINN. Sp. 27. *Cepa oblonga*. C. B. Pin. 71.

H. *Allium umbellatum foliis teretibus radicalibus*, scapi longitudine, floribus tubulosis. HALL. *All. n.* 14. *Allium Schoenoprasum*. LINN. Sp. 28. *Cepa sessilis*. TABERN. ic. p. 486.

H. *Allium foliis gramineis, umbella sphaerica, bicolore*. HALL. *All. n.* 16. *Comment. Soc. Gott. T. II.*

Allium radice senescente lignosa transversa. HALL. *All. n.* 19. *Allium scapo nudo ancipiti, foliis linearibus canaliculatis subtus subangulatis, umbella fastigiata*. (*angulosum*.) LINN. Spec. 22. *Allium s. Moly foliis Narcissinis*. CLVS. Pann. 221. 222. In M. Altenstollberg.

3. *Allium senescent*. LINN. Sp. 21. HALL. *Helv.* 297. n. 2. *Allium scapo ancipiti, teretiusculo, foliis ensiformibus, hinc paulo convexioribus*. Gmel. Sib. Lp. 53. T. XI. f. 2. In Sibiria.

Allium

Allium foliis radicalibus latissimis petiolatis, floribus umbellatis. HALL. *All.* n. 21. *Enum. Helv.* p. 297. *Allium vrsinum*. LINN. *Sp.* 33. LOB. *ic.* p. 159. *A. sylv. latifolium*. C. B. *Pin.* 74. Abunde in sylvis.

Allium foliis latissimis, radice oblonga, obducta reticulo. HALL. *All.* n. 20. *Allium Victorialis*. LINN. *Sp.* 4. *A. montanum latifolium maculatum*. C. BAVH. *Pin.* 74. *A. anguinum* CAM. *epit.* 329. In M. Kiffhäuser. In Alpes Helv. & Silvacis.

Allium umbellatum, foliis fistulosis, compressis, radice obducta reticulo. HALL. *Comm. Gott.* T. H. *Tab.* IX. p. 337. In Sibiria.

Allium foliis caulinis lanceolatis, floribus umbellatis, ex ala bulbiferum. HALL. *All.* 22. p. 382. *Allium*. LINN. *Sp.* 6. *Moly latifolium illiflorum*. C. B. *Pin.* 75.

Allium foliis radicalibus sessilibus, lanceolatis, floribus umbellatis. HALL. *All.* n. 23. RUYEN. *Engelb.* p. 57. *Allium Moly*. LINN. *Sp.* 16. *Moly latifolium luteum odore Allii*. C. B. *Pin.* 75. In Hungaria. M. Pyrenaeis.

Allium foliis teretibus, vagina bicorni, umbella pendula suavepurpurea. HALL. *All.* 386. n. 27. *Comm. icon.* In Sibiria. Prope Veronam.

Allium inodorum, umbella bulbifera, vagina bicorni, foliis carinatis. HALL. *All.* 391. n. 27. *Tab.* 2. *Allium carinatum*. LINN. *Spec.* 13. *A. sylvestre bicorni*, flore ex herbaceo albicante, cum triplici

in singulis petalis stria acropurpurea, RATT. Syn. III. p. 379. In apricis circa Gottingam.

CRINVM. LINN. 365.

In spatha bifolia flores umbellati. Corolla infundibuliformis, sexpartita, segmentis tribus vncinatis.

H. Crinum corollarum apicibus introsum vncinatis. *Americanum*. LINN. Sp. 3. Lilio-asphodelum *Americanus sempervirens maximus polyanthus albus*. DILL. Bibl. T. 166. f. 195. COMM. HEL. rar. 14. T. 14. In America.

Crinum foliis sublanceolatis planis, corollis obtusis. *africanum*. LINN. Sp. 4. Polyanthes sociatus umbellatis. Cliff. 126. In Aethiopia.

CLASSIS. IV.

TRIPETALOIDEAE.

Intermediae inter Liliaceas & Multifloras. Calyx trifolius. Petala tria. Sexus varius. Stamina semper in proportione ternaria. 3. 6. 9. 12. 24. Tubus tot, quot loculi aut filiculae. Fructus aut sexangularis, sexlocularis, aut plures filiculae triquetrae, in trigonum aut hexagonum positae. Perianthii florigeri involucri spathaceo excepti. Folia caulem vaginantia, in aliis liliacea integra, in aliis incisa, petiolata. Monocotyledones, gemmiparae. Omnes in paludosis crescunt. Vid. Comm. Dec. Gott. T. III. p. 432. f. 99.

BUTOMUS. LINN. 455. HALL. 299.

Involucrum trifolium umbelliferum. Calycis folia petalis similia colorata. Stamina novem. Siliquae sex distinctae, tubae apice bifido. Folia graminea.

Butomus. LINN. *Spec.* 1. HALL. *l. c.* Juncus floridus maior. C. B. *Pin.* 112. Gladiolus aquatilis. BOB. *Pempt.* 600. In fossis. P.

ALISMA. LINN. 418. Damasonium.

Involucrum trifolium flores colligens. Stamina sex. Capsulae plures in trigonum posita tuba simplici.

Alisma foliis ovatis acutis, fructibus obtuse trigonis. LINN. *Spec.* 1. Damasonium panicula verticillata, petiolis ramosis, siliquis brevissimis. HALL. *Helv.* 300. Plantago aquatica. C. B. *Pin.* 190. In fossis. P.

Alisma foliis gramineis & ovatis, caule repente, umbella pauciflora. Alisma natans. LINN. *Spec.* 4. Damasonium repens, potamogetonis rotundifolii folio. VAILL. *Act. Par.* 1719. p. 29. T. 4. fig. 8. Damasonium scapo geniculato, reptante, unisifero. BOEHM. *Fl. Lips.* p. 108. Rarius in inundatis circa Berolinum. Capsulae circiter octo. Flores ex spathis nonnullos prodeuntes semper vidi. P.

SAGITTARIA. LINN. 946. HALL. 300.

Monophyta. Mart. Stamina viginti, quatuor in capitula congestae. Fem. intra mares. Siliculae plurimae in globum collectae. Spathae trifoliae.

Sagit-

TRIPETALOIDAE.

243

Sagittaria foliis sagittatis, acutis LINN. *Spec. 1. Sagittaria aquatica. a. latifolia, angustifolia.* C. B. P. 194. Caulis verticillatus. In lacu ad Seeburg. P.

STRATIOTES. *Comm. Soc. Gott. T. III. p. 425.*

LINN. 607. *Aloides* BOERH. II.

p. 172.

Diphyta. Fl. Masc. Spatha bifolia flores plures colligens. Stamina duodecim, antheris erectis, corona filamentorum lanceolatorum cincta. Fem. Spatha bifolia uniflora. Corolla filamentorum circa tubas, ut in mare. Tubae sex, triangulares bifidas. Capsula oblonga, sexlocularis.

Stratiotes. *Comm. Gott. 1. c. p. 426. f. 99. Tab. IX. X. Stratiotes foliis ensiformi- triangularibus ciliatis, serratis.* *Aloides* LINN. *Sp. 1. Aloe palustris.* C. B. P. 286. In paludibus & stagnis profundioribus circa Berolinum. P.

HYDROCHARIS. *Comm. Soc. Gott. T. III.*

p. 431. LINN. 999.

Diphyta. Mas. Spatha bifolia triflora. Stamina novem, quorum tria interiora stylum e basi emittunt. Fem. Spatha nulla. Filamenta sex, lanceolata, ut in *Aloide*, posita circa tubas sex bifidas. Capsula ovata, sexlocularis. Caulis per flagella rept. Folia reniformia.

Hydrocharis. *Mascul. ramos.* LINN. *Spec. 1.*

MALL. *Helv. 30. 1. Comm. Gott. 1. c. T. III. f. 1-91. Nymphaea alba minor.* C. B. P. 193. *Microleuconymphaea.* BOERH. VI. *Lugd. 1. 10. p. 231.* In lacu ad Seeburg. P.

H

CLAS.

CLASSIS V.

RHQEADES. LINN.

Omnes androgynae. Calycis folia, vt plurimum decidua. Petala patentia. Stamina petalis multo numerosiora. Stylus nullus, sed tot stigmata plana, quot loculi capsulae polyspermae. Omnibus vires saporiferae.

NYPHIAEA. LINN. 579. HALL. 302.

Calyx quadrilaut quinquefolius, persistens. Petala numerosa. Stamina plurima. Stigma peltatum, radiatum. Capsula ovata.

Nymphaea calyce magno pentaphyllo, fructu pyriformi. HALL. *Helv.* 302. n. 1. LINN. *Sp.* 1. Nymphaea *lutea indur.* C. B. *Pin.* 193. BLACKW. *T.* 497. 499. In fossis suburbanis. P.

Nymphaea calyce tetraphyllo, fructu rotundo. HALL. *A. C. n. 2.* Nymphaea *alba.* LINN. *Sp.* 2. B. *Pin.* 193, BLACKW. *T.* 498. 499. In fossis suburbanis, & lacu *Seeburgensi.*

PAPAVER. LINN. 573. HALL. 303.

Calyx bifolius. Petala quatuor. Stigma peltatum, radiatum. Capsula multilocularis. Lacte albo manat.

Papaver calycibus, capsulis & foliis amplexicaulis incis. glabris. *sonniferum.* LINN. *Sp.* 1. Papaver *spontaneum* & *botanense* semine albo & nigro. C. B. *Pin.* 110. Rapi *nigrum.* BLACKW. *T.* 484. In hortis & cultis. Stigmata 10.

h. Papaver capsulis glabris, caudibus unifloris
 scabris foliosis, foliis pinnatis serratis. *orientale*
 LINN. *Spec.* 8. Papaver *orientale*, *viridissimum*, *flore*
magno. TOURN. *épin.* T. 127. In *Oriente*. Sigma-
 ta 16.

Papaver foliis pinnatifidis hispidis, fructu subro-
 tundo, glabro. HALL. *Hort.* 303. Papaver *rheas*.
 LINN. *Sp.* 4. Papaver *erectum*. B. *Pin.* 171.
 BLACKW. T. 2. In agris & hortis.

Papaver simile priori, sed fructu elliptico, &
 pilis cauli appressis, fortioribus. HALL. *Gott.* p. 88.
 Sponte in horto.

Papaver foliis pinnatifidis hispidis, fructu ellipti-
 co hispido. HALL. *Gott.* l. c. Pap. *capsulis clavatis*
hispidis, caule folioso multifloro. LINN. *Sp.* 2. Pa-
 paver *capitulo longiore hispido*. C. B. *Pin.* 172. In M.
Heynberg inter segetes. A.

h. Papaver caule ramoso, foliis lineari-pinnati-
 fidis, capsulis sulcatis hispidis. LINN. *Vps.* 136.
 Papaver *hybridum*. LINN. *Sp.* 1. Argemone *capitu-*
lo breviori. B. *Pin.* 172. In *Europa* australiori. A.

h. Papaver capsulis hispidis, scapo unifloro, nu-
 do, hispido, foliis simplicibus pinnato-sinuatis. nu-
 dicaule. LINN. *Sp.* 4. Papaver *erraticum nudicaule*,
flore flavo odorato. DILL. *Elth.* T. 224. f. 291. In
 M. Baldo. Wallia. *Sibiria*. B.

ARGEMONE. LINN. 374.

Papaveri valde similis. Calyx trifolius. Petala
 tria, quatuor, quinque, quot petala, tot valvae,

radix stigmatis & septe imperfecta capsulae spinosae.

H. Argemone capsulis quinquevalvibus, foliis spinosis. *Mexicana*. LINN. Sp. 1. Papaver spinosum. B. Prodr. 93. f. 92. CLVS. XCIII. In *Mexica, Jamaica, Europa australi*. A.

H. Argemone flore albo, saepe tripetalo. HALL. Gott. 89. Num Argemone capsulis trivalvibus. LINN. Sp. 2. ? A.

2 GLAUCIUM. HALL. 305. Chelidonii Sp. LINN.

Calyx & corolla, vt in Papauere. Stamina plurima basi connata. Siliqua longa, bilocularis, stigmate bifido.

H. Glaucium foliis tomentosis, imis pinnatis superioribus serratis, amplexicaulibus. HALL. Herb. 304. Chelidonium pedunculis unifloris, foliis amplexicaulibus sinuatis, caule glabro. LINN. Sp. 2. Papaver corniculatum luteum. B. Pin. 171. In *Europa Australi*. A.

H. Glaucium priori valde simile, sed flore dilute rubro, vngue flavo, & foliis fere glabris. Num Papaver corniculatum phoeniceum, pallescente flore. CLVS. XCII. ? Icon omnino convenit.

Glaucium foliis sessilibus pinnatifidis, caule hispido. Chelidonium corniculatum. LINN. Sp. 3. Glaucium hirsutum, flore phoeniceo. I. B. H. 254. Papaver cornutum, flore phoeniceo. CLVS. Hist. II. p. 91. In *Hungaria, Hispania, Montpetii*. A.

CHELIDONIUM. HALL. 305.

LVDW. 447.

Omnia prioris, sed stamina non connata, filiqua unilocularis, stigma capitatum, planta vndique fuc-
co aureo. *manit.*

Chelidonium foliis subrotundis. HALL. *Helo.* 305.

Chelidonium *pedunculis umbellatis. maius.* LINN. *Sp.*

1. Chelidonium *angulare.* BLACKW. *F.* 98. β. Chelid.

foliis laciniatis. CLVS. *Hist.* II. p. 203. Vbiq̃

in ruderatis. A.

CHRISTOPHORIANA. HALL. 306.

Actaea LINN. 568.

Calyx quadrifolius, caducus. Petala aliquot exi-
gua, quasi degenerantia stamina. Bacca globosa,
stigmatē oblique depresso.

Christophoriana. DOD. *Purg.* p. 231. CLVS. *Hist.*

2. p. 86. Actaea racemo ovato, fructibusque bac-

catis. *spicata.* LINN. *Sp.* 1. Aconitum *bacciferum.*

C. B. *Pim.* 192. In sylvis. & ad sepes. P.

CLASSIS. VI.

MULTISPERMAE.

In univēsum huius classis plantis stamina sunt
multo plura, ac petala, & post florem siliculae, in
aliis polyspermae dehiscētes, in aliis monospermae,
a semine non discedentes.

ORDO. VI.

Multifiliquae siliculis non dehiscentibus, monospermis, liberis.

THALICTRVM. LINN. 617. HALL. 307. 00

Calyx nullus, petala quatuor fugacia. Siliculæ fulcatae, tubis brevissimis. Flores paniculati.

H. Thalictrum foliis lanceolato-linearibus intergerminis. *angustifolium*, LINN. Sp. 9. Thalictrum caule folioso, foliis angustissimis non dentatis. HALL. Helv. 307. Thalictrum pratense angustissimo folio. C. B. Prædr. T. 146. In pratis Germaniae, v. g. circa Berolinum. P.

H. Thalictrum caule folioso, sulcato, panicula multiplici erecta. *flavum*, LINN. Sp. 10. HALL. H. 308. Thalictrum pratense, LINN. Lapp. 224. CLVS. 224. Thalictrum maius siliqua striata. BAVH. Pin. 336. Folia eundo latiora, sine tridentato. In Germaniae, Helvetiae pratis humidis. P.

H. Thalictrum foliis latis, rotunde tridentatis, caule nudo laxissime paniculato. HALL. Helv. 208. n. 4. Thalictrum foliis sexpartitis, siliquis cernuis, minus. LINN. Sp. 6. Thalictrum minus. B. Pin. 337. Thalictrum minus, rubeae folia, staminibus flavis. SEGVIER. T. XI. In Helvetia, circa Jenam. P.

H. Thalictrum floribus pentapetalis, radice tuberosa, *tuberosum*, LINN. Sp. 31. Thalictrum minus asphodeli radice, magno flore. 1. B. H. 271. In Hispania. P.

SPIRAEA. LINN. 554. *Filipendula* HALL. *Vlmaria*. *Barba Caprae*. *TOVRN.* *Aruncus*.
LINN. ed. pr.

Calyx stellatus. Petala quinque & stamina ex calyce orta. Capsulae nonnullae exiguae compressae, saepe intortae.

Spiraea foliis lanceolatis obtusis serratis nudis, floribus duplicato cymosis. Salicifolia. LINN. *Spec.* 1. *Spiraea Salicis folio*. 1. R. H. CLVS. *Hist. Lp.* 80. In Sibiria sponte. Fr.

Spiraea foliis supradecompositis, spicis paniculatis, floribus dioicis. LINN. *Sp.* 8. *Aruncus H. Cliff.* 463. *Filipendula sexu distincta foliis aequalibus.* HALL. 306. n. 1. *Barba caprae floribus oblongis.* 1. R. H. B. *P.* 163. Passim in sylvis. v. g. circa *Casselas.* P.

Spiraea foliis pinnatis, impari maiore lobato, floribus cymosis. Vlmaria. LINN. *Sp.* 10. *Filipendula foliis ex ovalibus acutis, extremo trilobato, minimis pinnulis intermistis.* HALL. l. c. n. 2. *Vlmaria.* CLVS. *Hist.* CXVIII. BLACKW. 465. *Barba caprae floribus compactis* B. *Pin.* 164. Copiose in prat. P.

Spiraea foliis pinnatis, foliolis vniformibus serratis, caule herbaceo, floribus cymosis. LINN. *Sp.* 9. *Filipendula foliis longis, angustis, extremo trilobato, minimis pinnulis intermistis.* HALL. 307. n. 3. *Filipendula vulgaris.* B. *Pin.* 163. BLACKW. *T.* 467. Super *Wernigerode.* Circa *Berolinum.*

ORDO. II.

Multisiliquae siliculis polyspermis dehiscenribus.

a. Floribus nectario instructis.

FRAXINELLA. HALL. 311. *Dictamnus*.
LINN. 468.

Calyx quinquefolius. Petala quinque. Filamentis decem flaminum glandulae adspersae. Siliculae quinque connatae, tubi unica.

Fraxinella. HALL. 311. *Dictamnus albus*. LINN. *Sp. i. Mat. med.* 208. BLACKW. *T.* 75. a. flore rubro & flore albo. In Germania, Gallia, Italia. P.

AQVILEGIA. LINN. 605.

Petala quinque patentia, totidem nectaria alterne cum petalis posita, infundibuliformia. Stamina plurima. Siliculae quinque.

Aquilegia nectariis incurvis, vulgaris. LINN. *Sp. i.* *Aquil.* foliis rotunde lobatis. HALL. *H.* 310. *Aquilegia*. BLACKW. *T.* 409. & *auctorum*. In sylvis & pomariis. P.

H. *Aquilegia nectariis rectis, flaminibus corolla longioribus.* *Canadensis*. LINN. *Sp.* 3. *Aquilegia pumila praecox*. *Canadensis*. LOR. *ic.* 26. In Virginia, Canada. P.

ACONITVM. LINN. 603.

Petala quinque inaequalia, supremum galeatum, sub quo in longis pedunculis nectaria duo corniculata. Siliculae tres & quinque.

Aconi-

Aconitum foliis petalis multifidis villosis, petalo supremo cylindraceo. LINN. *Spec.* 1. HALL. *H.* 312. n. 1. Aconitum *lycoctonum luteum*. C. B. *Pin.* 183. In sylvis v. g. circa Lützelengen. P.

H. Aconitum foliis multipartitis, laciniis linearibus incumben- tibus, squarrosis. *pyrenaicum*. LINN. *Sp.* 4. HORT. *Vpf.* 152. In Sibirja. Pyrenaeis. P.

H. Aconitum foliorum laciniis linearibus super- ne latioribus, linea exaratis. *Napellus*. LINN. *Spec.* 3. HALL. *l. c.* n. 2. Aconitum *caeruleum* f. *Napellus*. B. *Pin.* 183. In Helvetia. Gallia.

H. Aconitum foliis multifidis, laciniis semiparti- tis, superne latis, *variegatum*. LINN. *Sp.* 2. Acon. *caeruleum minus*. B. *Pin.* l. c. CLVS. Aconit. *lycoct.* IX. *Judenbergense*. p. XCVII. In Italia, Bobemia. P.

H. Aconitum floribus pentagynis. LINN. *Sp.* 1. Acon. *foliorum laciniis linearibus distinctis ubique eius- dem latitudinis*. LINN. *Mat. med.* 270. HALL. *H.* 313. Aconit. *salutiferum* f. *Anthora*. LINN. *Sp.* 5. In alpi- bus Helveticis. M. Thury prope Genevam. P.

DELPHINIVM. LINN. *Sp.* 602.

Calyx nullus. Petala quinque inaequalia, supre- mum in conicum calcar abit, quod simile fere calcar replet modo simplex modo duplex, intra petala in bracteam ovatam abiturum. Silicula una, & tres.

Delphinium nectariis monophyllis, caule subdi- viso. *Consolida*. LINN. *Sp.* 1. Delphinium *flore uni- siliquo*, foliis tenuissime divisis. HALL. *Helv.* 314.

Delphinium segatum, L. & H. *Consolida regalis arvensis*, B. Pin. 142. Inter segetes A.

H. *Delphinium nectaris monophyllis*, caule simplicis. *Atacis*, LINN. *Spec.* 2. *Delphinium sativum*, RIV. T. 123. *Consolida regalis hortensis*, B. Pin. 142. Siliquae villosae. A.

H. *Delphinium nectaris diphyllis*, labellis integris, floribus subsolitariis, foliis compositis linearimultipartitis, *grandiflorum*, LINN. *Sp.* 4. *Hort. Vpf.* 150. *Delphinium perenne, foliis tenuissime divisis, filiquis villosis*, HALL. *Gott.* 94. *Delphinium elatius subincanum perenne, floribus amplis azureis*, AMM. *Rutb. n.* 175. BOLOF, *Hort.* 61, T. 3. In *Sibiria*, P.

Delphinium nectaris diphyllis, labellis bifidis apice barbatis, foliis semiquinquelobis, subhirsutis, lobis latis, serratis. *Delphinium elatum*, LINN. *Sp.* 15. *Hort. Vpf.* 151. *Delphinium perenne, aconiti folio ampliari, floribus coeruleis longioribus pedunculis infidentibus, minus villosum*, AMM. n. 174, HALL. *Gott.* 93. *Aconitum lycoctonum flore Delph.* I. *Silvest. GLVS.* XCIV. Folia semiseptiloba, lobis latis, serratis ciliatis. In *Sibiria*.

H. *Delphinium nectaris diphyllis*, foliis multipartitis, glabris, laciniis linearibus superne latioribus, pinnatifidis. Pro distincta specie haberi mereatur. BINNAEVS autem in *diff. de plantis hybridis*, *Amoen. Acad. Vol.* III. p. illam esse credit plantam hybridam ornatam ex *Delphinio elato*, matre, & *Aconito Napello*, patre. Flos prioris speciei. P.

H. Del-

11. *Delphinium nectaris diphylis*, foliis palmatis, lobis integris. LINN. *Sp.* 6. *Mat. med.* 269. *Delphinium orient. platani folio*. HALL. *Gott.* 93. *Staphis-agria*. B. *Pin.* 324. BLACKW. *T.* 265. In *Hungaria. Italia, circa Monspetium*. P.

RESEDA. LINN. 207. Luteola.

HALL. 315.

Petala digitata, fissa, ad basin supremi squama nectarifera. Capsula valloocularis inter stylos aperta.

Reseda foliis lanceolatis integris, calycibus quadrifidis. LINN. *Sp.* 1. *Luteola*. HALL. *l. c.* BLACKW. *T.* 283. *Luteola herba salicis folio*. BAVH. *Pin.* 10. Petala tria, Styli tres, stamina plurima. In gramineis. v. g. circa *Pless*. A.

H. *Reseda foliis omnibus trifidis, inferioribus pinnatis. lutea*. LINN. *Sp.* 7. HALL. *H, l. c.* *Reseda foliis rariter pinnatis undulatis*. HALL. *Gott.* 94. *Reseda vulgaris*. B. *Pin.* 100. Calyx sexfidus, petala sex, styli tres. In *Thuringia & arenosis Europae australioris*. A.

H. *Reseda foliis pinnatis, floribus tetragynis. alba*. LINN. *Sp.* 6. *Reseda foliis pinnatis planis, flore spicato*. HALL. *Gott.* 93. *Reseda maxima*. B. *Pin.* 100. Calyx sexfidus, petala sex omnia trifida, stamina undecim. *Monspeli*. A.

H. *Reseda foliis integris, floribus odoratis, calyce corollam vix superante*. HALL. *Gott.* 95. *Reseda Aegyptiaca odoratissima*. Calyx sexfidus, petala sex, stamina plurima alba. In *Aegypto*. A.

H. *Reseda foliis integris trilobisque, calycibus maximis hexaphyllis. Phyteuma.* LINN. *Sp.* 8. *Reseda minor vulgaris.* I. R. H. 423. *Resedae affinis Phyteuma.* B. *Prodr. F.* 42. Calyx corolla multo maior, petala sex, styli tres, stamina pauciora, rubella. In *Gallia, Italia.* A.

PARNASSIA. LINN. 333. HALL. 316.

M. Squamae quinque intra petala cordatae, singulae 15 stylis globuliferis instructae. Stamina quinque. Capsula quadrangularis, ex apice hians.

P. *Parnassia.* LINN. *Sp.* 1. ROYEN. 420. HALL. *f. c.* *Gramen Parnassi.* LOB. *ic.* 603. *Gramen Parnassi albo simplici flore.* C. B. *Pin.* 309. In pratis viginosis. P.

NIGELLA. LINN. 606. HALL. 316.

Flos nudus. Petala quinque. Squamae octo, petiolatae, bifidae, lineatae; in singularum origine fovea nectarifera, cui lamina lanceolata applicatur. Capsulae quinque aut decem, liberae modo, modo connatae in totidem stylos revolutos abeunt.

Nigella flore foliis nudo, pistillis corollam aequantibus. HALL. *Helv.* 316. *Nigella pistillis quinis, petalis integris, capsulis turbinatis arvensis.* LINN. *Sp.* 3. *Nigella arvensis cornuta.* I. R. H. B. *Pin.* 145. Inter segetes, v. g. circa *Lütteleugen.* A.

H. *Nigella petalis subtricuspidatis, foliis subpilis, sativa.* LINN. *Sp.* 2. H. *Vps.* 154. *Nigella rapae & folio minore.* I. B. III. 1. p. 208. In *Palaeestina. Aegypto.* A.

Nigella. floribus involuctis foliis exceptis, *Damascena*. LINN. Sp. I. H. Kp. 53. *Melanchium sylvestre*. LOB. ic. 140. β. Var. flore pleno. Inter legetes *Europae australis*. A.

Nigella pinnatifida dens corollae longioribus, *orientalis*. LINN. Sp. 3. HALL. Gott. 96. *Nigella orientalis*, flore flava, semine alato plano. TOWN. Cor. 19. Capsulae 9 ad decem. Circa *Aleppum*. A.

GARIDELLA. LINN. 507.

Nigellae proxime affinis. Sed stamina decem, capsulae tres, styli vix vili, petala exigua.

H. *Garidella*. LINN. Sp. I. *GARIDELLA* Pruv. Eugl. *Nigellastrum* raris, foeniculaceis, foliis. MAGN. Hort. T. 143. *Nigella cretica*, folio foeniculi. C. B. Prud. 75. In *Provincia Creta*. A.

HELLEBORVS. LINN. gen. ed. prior. 461. HALL. 317. *Trollius*. *Isopyrum*. LINN.

Flos nudus. Intra orbem petalorum nectaria tonica, tubulata. Capsulae plurimae, compressae, liberae.

Helleborus vaginis pallidis, vix serratis, flore virente. HALL. *flor.* 317. n. 2. *Helleb. caule multifloro, folioso, foliis pedatis, (foetidus.)* LINN. Sp. 4. *Helleb. niger foetidus*. B. Pin. 185. *Helleboraster*. BLACKW. T. 79. In *Germania meridionali*. *Helvetia*. P.

Helleborus vaginis foliaceis, digitatis, flore virente. HALL. l. c. n. 1. *Helleb. caule multifloro, folioso, foliis digitatis. (viridis.)* LINN. Sp. 3. *Helleb.*

leb. *niger hortensis*, flore viridi. B. Pin. 185. Circa Grubenhagen, Viennam. P.

H. Helleborus scapo subunifloro subnudo, foliis pedatis. *niger*. LINN. Sp. 2. Helleb. *niger flore rufes.* D. P. 186. Helleborus *niger legitimus* Cava. Hist. I. p. 279. In Austria. Italia. B.

H. Helleborus caule plerumque unifloro, petalis et filiquis plurimis. HALL. Gott. 97. Herb. 218. Trollius corollis conniventibus, nectaris longitudine staminum. LINN. Sp. 1. Ranunculus flore globoso. D. P. 287. Trollius flos GESN. Hort. 144. Petala biseriata 14, nectararia novem. In montibus Helvetiae. P.

H. Helleborus foliis repetito multifidis, nectaris planis. HALL. Gott. 97. Helleborus folio fumarariae AMM. Rusb. 74. T. 12. Noppyrum stipulari subulatis, petalis acutis. LINN. Sp. 1. In Siberia. A.

β. nectario carentibus.

PAEONIA. LINN. 600.

Calyx quinquesolius, inaequalis, quinque petala, capsulae duae, tres, ovatae.

H. Paeonia foliis difformiter lobatis. HALL. Herb.

H. Paeonia *officinatis*. LINN. Sp. 1. Paeonia *fermina* C.B. BLACKW. T. 65.

β. Varietas flore pleno hort. Sponte in montibus Helvetiae.

Paeo-

H. Paeonia laciniis foliorum linearibus. Folia decomposita, pinnulis ultimis linear-lanceolatis, nervo depresso; flos nido folioso excipitur. Petala egregie phoenicea. Planta a priori diversissima & MARTINO missa, in *Vkrainia* habitat. P.

CALTHA. LINN. 623. *Populago* HALL. TÖVEN.

Petala quinque sine calyce. Capsulae strobilatae dispositae.

Caltha palustris. LINN. *Spec. 1. C. B. Pin.* 276. *Populago.* HALL. *H.* 219. TABERN. 750. *C. 16.* L. R. H. Ubique in vdis. Varietas *fl. pleno bore.*

ORDO. III.

Multisiliquae siliiculis non dehiscensibus monospermis, receptaculo communi exceptis. *Gymnosperrmae.*

a. Petalis & staminibus ex ambitu receptaculi ortis.

LIRIODENDRON. LINN. 609. *Tulipifera,*

Calyx trifolius petala sex. Stamina viginti quatuor in sinibus squamarum, stigmate globoso, in receptaculo conico.

H. Liriodendron. LINN. *Sp. 1. H. Vrf.* 354. *Tulipifera arbor Virginiana.* HERM. *Lugdb.* 612, *T. 613.* CATESB. *Car. 1. p. 48.* *T. 48.* RYTTNER. *H. Canon.* In *America septentrionali.* Fr.

MYOSURUS. LINN. 318.

Calycis folia quinque ex. ungue angusto, repanda. Petala totidem fere similia, minora. Stamina ad duodecim. Semina in receptaculo longissimo.

1. *Myosurus*. LINN. *Sp.* p. 284. HALL. *Gott.* 98. *Gauda muris*. LOBEL. ic. 49. DODON. *Pempt.* 112. In collibus apricis v. g. prope *Gottingam* in *M. Hagen*, A. 10.

ADONIS. LINN. 618.

Petala octo aut duodecim. Semina in receptaculo longissimo.

Adonis floribus octopetalis. fructibus subcylindricis. *annua*. LINN. *Sp.* 1. HALL. *H.* 318. *a. flore phoeniceo*. *β flore citrino*. Inter segetes *γ. petalis brevioribus flore rubro, maior*. hort. A.

H. Adonis flore dodecapetalo, fructu ovato, *vernalis* LINN. *Sp.* 2. Adonis *perennis dodecapetala* var. HALL. *H.* 319. *Helleborus niger tenuifolius*, *Bupthalmi flore*. C. B. *Pin.* 186. Circa *Halberstadt*. In *Borussia*. P.

RANUNCULUS. LINN.

Calycis folia & petala quinque, ad singulorum basin fovea nectarifera.

1. *Ranunculus* foliis ovato lanceolatis, petiolatis, caule declinato. *Flammula*. LINN. *Sp.* 1. HALL. *H.* 322. *Ranunculus longifolius palustris minor*. B. P. 180. *Flammula* *Ranunculus*. D. D. *Pempt.* 432. In vdis vbiq. P.

OTIV.

Ranun-

g. Ranunculus foliis lanceolatis, sessilibus, caule erecto. *Lingua. LINN. Sp. 3. HALL. Herb. 1. 2. 211. Ranunculus longifolius palustris major. C. B. Pin. 34. In lacubus ad Seckburg & Deucherhausen. P.*

h. Ranunculus foliis inferioribus palmatis, summis digitatis, fructibus oblongis. *Ranunculus Sp. 15. HALL. Herb. 1. 2. 211. Ranunculus palustris apertifolio, latior. C. B. Pin. 34. In lacubus ad Seckburg & Deucherhausen. P.*

i. Ranunculus foliis radicalibus reniformibus, orengatis incisus, caulibus digitatis, linearibus, caule multifloro. *auricomus. LINN. Sp. 13. Ranunculus auricomus folia subrotunda. C. B. Pin. 178. In sylvis. P.*

k. Ranunculus foliis omnibus quinatis lanceolatis, inciso-teretibus. *aconitifolius. LINN. Sp. 16. Ranunculus mont. aconiti folio albus. C. B. Pin. 182. In sylva Hercynica circa M. Bructerum. P.*

l. Ranunculus foliis ternatis, internodiisque foliis trifidis incisus, caule inferne ramoso. *astinatus. LINN. Sp. 18. Ranunculus grumosa radice ramosa. B. Pin. 181. Varietates plurimae ex colore flore pleno in hortis. In Asia. P.*

m. Ranunculus foliis compositis, caule simpliciter villosa subnodo unifloro, radice tuberosa. *An Ranunculus mont. leptophyllos, asphodeli radice. C. B. Ephr. 1. p. 312. T. 311. Icon egregie conveniens in horto flore pleno adest. P.*

n. Ranunculus radice simplici globosa. *LINN. Cliff. 230. HALL. Herb. 323. Gey. 180. Ranunculus polycephalus retroflexus, pedunculis sulcatis, caule erecto*

Ho. foliis compositis. LINN. Sp. 25. Ranunc. *bulbosus*. LQB. ic. 666. P.

1. *Ranunculus* caule ex geniculis radicato, foliorum lobis petiolatis. HALL. Herb. 324. n. 6. *Ranunculus calycibus patulis, pedunculis sulcatis, foliolonibus repentibus, foliis compositis.* LINN. Sp. 26. *Ranunc. prat. repens. hirsutus.* B. Pin. 179. Num Varietas? Uterque in pratis & gramineis. P.

2. *Ranunculus* caule erecto multifloro; folio pentagono, lobis sessilibus serratis. HALL. H. 324. n. 7. *Ranunc. acris.* LINN. Sp. 28. *Ranunc. pratensis erectus, acris.* B. Pin. 178. In pratis. P.

3. *Ranunculus* priori simillimus, foliorum lobis sessilibus laciniatis. HALL. Gott. 101. *Ranunculus polyanthemorum.* LINN. Sp. 27. LQB. ic. 666. P.

4. *Ranunculus* foliis tripartitis acute sinuatis tomentosis. LINN. Sp. 29. *Ranunc. magnus valde hirsutus, flore luteo.* L. B. Hist. 3. p. 417. In sylvis. Num varietates? P.

5. *Ranunculus* seminibus aculeatis, foliis superioribus decompositis linearibus. *arvensis.* LINN. Sp. 31. HALL. H. 323. n. 5. *Ranunculus arvensis echinatus.* B. Pin. 179. In agris. A.

6. *Ranunculus* seminibus aculeatis, foliis simplicibus palmatis incisis. *muricatus.* LINN. Sp. 32. *Ranunc. creticus echinatus latifolius.* F. ALP. Exot. 262. B. Pin. 180. In Europa meridionali. A.

7. *Ranunculus* ramis unifloris, foliis trifidis, siliquis incurvis. HALL. Gott. 99. *Ranunc. foliis siliiformi-ramosis, calycibus appendiculatis.* LINN. Sp.

25. *Ranunc. ceratoccephalus*, *seminibus falcatis*, in *spicam adactis*. MQRYS. II. 8. 4. T. 28. f. 22. Stamina pauca ad decem usque. In Europa australi. A.

Ranunculus foliis submersis capillaceis, emeris peltatis. *aquatilis*. LINN. *Spec.* 37 HALL. *Helv.* 328. n. 16. *Ranunc. aquat. folio rotundo & capillaceo*. B. PIN. 180. In stagnis & inundatis.

β. *Ranunculus* foliis capillaceis, circumscriptione vaga. HALL. *l. c.* n. 17. In aquis parum velocibus v. g. *im Rasenübren-Wasser*.

γ. *Ranunculus* foliis capillaceis, circumscriptione rotunda. HALL. *l. c.* n. 18. *Ranunculoideis folio circinato, tenuissime diviso*. VAILL. In aquis quietis. v. g. *beym Schützenhof*.

δ. *Ranunculus* foliis capillaceis, circumscriptione oblonga. HALL. *l. c.* n. 19. In aquis rapidioribus. v. g. *in der alten Leim*. Varietates esse videtur LINNAEO, LUDWIGIO, BOEHMERO, nec HALLERUS plane dissentit.

Ranunculus foliis linearibus, brevissime bifurcatis. HALL. *Rupp.* 104. MAPP. *Alsat.* 263. *cann. ia*. Nam satis a prioribus differt?

Ranunculus foliis coriatis angulatis petiolatis. LINN. *Spec.* 9. *Ficaria*. HALL. *Helv.* 321. *Chelidonium minus*. BLACKW. F. 51. *Chelidonia rotundifolia minor*. BAVH. *Pin.* 167. Ad sepes & in umbrosis. Calyx trifolius, petala 6. P.

HEPATICA. HILL. LINN. ed. pr. 457. Trinitas

HALL. 320. Anemones spec. LINN.

ed. ak.

Calyx trifolius. Flos bis tripetalos. Receptaculum papposum.

Hepatica. LINN. Vpf. 155. Mat. med. 277.

BLACKW. T. 207. Trinitas. HALL. l. 2. Anemone

foliis trilobis integerrimis. LINN. Sp. 1. Vbique

in dumetis. β . flore pleno, albo, rubro, coeruleo

hortens.

ANEMONE. LINN. 614. Pulsatilla

Involucrum foliaceum a flore remotum. Petala duorum triumve ordinum.

Anemone semine levi, petalis quinis, subrotundis, luteis. HALL. Helv. 329. n. 1. Anemone ranunculoides. LINN. Sp. 17. Ranunculus nemorosus luteus. B. Pin. 178. Primo vere in sylvis. P.

Anemone semine levi, petalis senis ellipticis. HALL. H. 330. n. 2. Anemone nemorosa. LINN. Sp. 15. Ranunculus phragmitis, vernus, albus. 1. R. H. In sylvis. P.

Anemone pedunculo nudo, seminibus seta plumosa terminatis. LINN. Cliff. 224. sylvestris. Sp. 10. Anemone sylvestris alba maior. B. Pin. 176. In sylva prope Hessendreisck & in monte super Lützelangen. P.

n. Anemone caule ramoso, petalis lanceolatis. ROYEN. Lugdb. 488. Anemone pedunculis alternis longissimis, fructibus cylindricis, seminibus

Anemone virginiana. (Virginia.) LINN. Sp. 12. HERM. Parad. T. 18. In Virginia. P.

G. Priori similis, sed constanter omnibus partibus altera duplo maior, femina villosissima, serius semper floret.

Anemone tubis caudatis, foliis omnibus duplicato pinnatis, pinnulis remotis. HALL. Helv. 331. n. 5. Anemone foliis caulinis ternis connatis supra-decompositis multifidis, seminibus hirsutis caudatis. (alpina.) LINN. Sp. 8. Pulsatilla flore albo. B. Pin. 177. In M. Bructero. P.

H. Anemone tubis caudatis, foliis tenuissime divisis, flore connivente nutante. HALL. Gott. 103. Pulsatilla foliis decompositis pinnatis, flore pendulo, limbo reflexo. LINN. Cliff. 223. Anemone LINN. Sp. 6. Pulsatilla flore minor nigricante. B. Pin. 177. In campis arenosis v. g. circa Berolinum. P.

H. Anemone (Pulsatilla) foliis simpliciter pinnatis, foliolis lobatis, flore erecto. LINN. Fl. Suec. edit. prior. 448. Anemone vernalis. LINN. Sp. 4. Pulsatilla involucre piloso nitido Fl. Lapp. 223. Pulsatilla apii folio, vernalis. B. Pin. 177. HALL. Helv. 332. cum icone. Petala villosa, tubae caudatae. P.

CLEMATIS. LINN. 616.

Petala quatuor. Antherae lateri filamentorum adnatæ. Styli subulati. Folia coniugata.

Clematis foliis pinnatis, foliolis cordatis, inaequaliter incis, serratis. Vitalba. LINN. Sp. 5. HALL. H. 334. Clematis sylvestris latifolia. B. Pin. 300. In sylvis & sepibus. P.

H. Clematis foliis compoſitis decapofitiſſimis
foliolis quibusdam trifidis. *Vienna*. LINN. *Sp.* 1121
Flammula ſcandens, flore violaceo, clauſa. DILL.
Elb. 144. *T.* 118. *f.* 444. Clematis purpurea re-
pens, petalis ſlorum coriaceis. RAIL. *Hiſt.* 1928. B.
Variet. flore pleno. In *Virginia*. P.

H. Clematis foliis pinnatis, foliolis ovato lanceo-
latis integerrimis, caule erecto. *recta*. LINN. *Sp.* 8.
Clematis ſ. Flammula ſurrecta alba. I. B. II. p. 127.
Flammula. GLVS. *Hiſt.* I. p. 124. In *Austria*. *Men-*
ſpelii. P.

H. Clematis foliis ſimplicibus ovato lanceolatis.
integrifolia, Clematis coerulea erecta. B. *Pin.* 380.
Clematis coerulea, Pannonica. GLVS. *Hiſt.* I. p.
123. P.

ORD. II.

SENTICOSAE.

In vniuerſum, his omnibus, quæ fructus ſimilitudo
cum prioribus iungit, petala quinque, ſtamina plu-
rima ex calyce orta, radix perennis, & vires adſtrin-
gentes. Præterea Geum, Comarum, Potentilla,
Fragaria, & Alchimilla habent calycem ſemidecom-
fidum aut ſemioctofidum, laciniis alternis minoribus,
omnes autem ſtylos lateri germinis inſertos.

GEVM. LINN. 564. Caryophyllata, ro var.

J. S. I. HALL. 335.

Styli longi, geniculati, pilofi.

Geum floribus erectis, fructu globoſo, triſtis
vincinatis nudis, foliis lyratis. *urbanum*. LINN. *Sp.* 21

Caryo-

Caryophyllata pinnis punctioribus rhomboides, rubrum sine punctis. HALL. L. c. n. 1. Caryophyllata vulgaris. B. Pin. 321. BLANKW. T. 253. Ad sepos. P.

β. *Caryophyllata Canad. foliis acutioribus flore lateo-malo. EV.*

γ. *Geum floribus erectis fructu globoso, arista vinctatis nudis, foliis ternatis. Virginianum. LINN. Sp. 1. Caryophyllata Virginiana, albo flore minore, radice inodora. HERM. Parad. T. III. In Virginia. P.*

Geum floribus nutantibus, fructu oblongo, arista plumosis, rivale. LINN. Sp. 3. Caryophyllata pinnis frequentioribus, extrema semitrifida, flore connivente nutante. HALL. Helv. 336. Caryophyllata aquatica, nutante flore. B. Pin. 321. 1. R. H. In pratis humidis & ad rivula. β. flore pleno in orto. P.

COMARVM. LINN. 563.

Petala segmentis calycis rubri triplo minora. Styli breves. Receptaculum carnosum.

Comarum. LINN. Sp. 1. HALL. H. 337. Quinquefolium palustre rubrum. B. Pin. 325. In vallis v. g. ad lacum prope Seeburg. P.

POTENTILLA. LINN. 559. HALL. 338.

Tormentilla LINN. Quinquefolium. T.

Calycis quinque segmenta patentia, quinque circa fructum maturum se complicantia. Styli breves.

1. *Potentilla foliis pinnatis, caule fruticoso, fructu nigro.* LINN. Sp. 1. *Potentilla fruticosa, foliis serratis.* A. medic. trífido, HALL. Gott. 106. *Pentaphylloides rectum frutescens.* WALTH. T. 17. In Anglia, Sibiria. Fr.

2. *Potentilla foliis pinnatis, pinnis perangulatis, quinquefidis.* HALL. Gott. 105. *Potentilla foliis bipinnatis, segmentis integerrimis, distantibus, subius tomentosis, caule decumbente.* (multifida.) LINN. Sp. 4. In Sibiria. Fr.

3. *Potentilla foliis pinnatis ferratis, caule repente.* Anserina, LINN. Sp. 2. *Potentilla repens, scapo nudo unifloro, pinnis plurimis confertis, minoribus intermissis.* HALL. H. 338. n. 2. Anserina BLACKW. T. 6. Ubique ad vias. P.

4. *Potentilla foliis pinnatis, bifidis, pinnis tredecim ovatis, crenatis, decurrentibus.* HALL. Gott. 106. P.

5. *Potentilla foliis pinnatis, pinnis septenis, subrotundis, crenatis, albo flore.* HALL. Gott. 107. *Potentilla foliis pinnatis ovatis, serratis, quinque crenatis, caule erecto.* (rupestris.) LINN. Sp. 6. *Pentaphylloides erectum.* L. B. Hist. II. p. 398. In Marchia. In monte Hundsrück. P.

6. *Potentilla foliis novenis palmatis, in apice ferratis.* HALL. Gott. 107. *Folia glabra, angustata, in apice acuta serrata, cuneiformia.* P. 107.

7. *Potentilla foliis radicalibus septenis, cuneiformibus, serratis, caulibus ternatis, ramis filiformibus, decumbentibus.* LINN. Cent. I. Plant. n. 35. 2. Folia

ex stipulis amplexicaulis digitatis palmata, ampla, media biuncialis, obverse ovata, in toto ambitu magnis dentibus profunde serrata, pilosa. Calyx villosus segmentis alterne integris & profunde tri-
fidis, corollae magnitudine. Petala flava. P.

III. *Potentilla* foliis digitatis lanceolatis, serratis utrinque subpilosis, caule erecto. LINN. Sp. 10. *Quinquefolium minus*. L. TABERN. 351. *Quinquefolium rectum, luteum*. B. Pin. 325. *Quinquefolium mont. erectum, bifidum, luteum*. GARIDEL. T. 83. Folia quinata, petala sulphurea. P.

IV. *Potentilla* foliis quinatis, palmatis, sericeis, serratis. HALL. Gatt. 197. *Potentilla alba*. LINN. Sp. 16. *Quinquefolium album minus*. B. Prodr. 325. Pedunculi florum ex radice. In *Morchia Branden-
burgensi*. P.

Potentilla foliis quinatis, longe petiolatis, caule repente. HALL. H. 240. n. 9. *Potentilla reptans*. LINN. Sp. 17. *Quinquefolium minus repens*. B. Pin. 325. BLACKW. T. 454. Caulis ex intervalis radicles emittit. In muris aliisque locis argillofis. P.

Potentilla foliis quinatis, obtuse in apice serratis, petalis cordatis. HALL. H. 341. n. 6. *Quinquefolium minus repens luteum*. I. R. H. C. B. L. c. *Quinquefolium minus*. TABERN. p. 123. Primo vere locis ficis. P.

Potentilla foliis quinatis angustissimis, subtus tomentosis, caule erecto. HALL. H. 341. n. 10. *Potentilla argentea*. LINN. Sp. 11. *Quinquefolium yedo argenteo*. B. Pin. l. c. *Quinquefolium petrae-*

an. matus. TABERN. p. 122. In maris & locis
secis. P.

1. *Potentilla foliis quinatis, flore tetrapetalo, cau-
le recto.* HALL. H. n. 11. *Tormentilla caule erecto.*
LINN. Sp. 1. *Tormentilla.* LINN. Mat. med. 249.
BLACKW. T. 445. *officin.* In sylvis & locis steri-
libus. P.

2. *Potentilla foliis ternatis hirsutis, caule erecto
umbellifero.* HALL. Gorr. 108. An *Pentaphylloides
matus erectum, fragariae folio, flore luteo.* MO-
RIS. Hist. 2. p. 193. 5. 11. T. 20. f. 2. Icon fa-
cte convenit. Flores parvi inter folia latentes, in um-
bellas dense congesti. Caulis & folia pilosa. A.

3. *Potentilla procumbens, foliis ternatis, hirsu-
tis.* HALL. Gorr. 1. c. *Potentilla Norwegica.* LINN. Sp.
19. *Potentilla foliis ternatis incis, caule diffuso.*
GRIFF. 195. In Norwegia & Canada. P.

4. *Potentilla foliis ternatis, ampliter crenatis,
subtus tomentosis.* HALL. Gorr. 1. c. *Potentilla fo-
liis ternatis incis, subtus tomentosis, caule hasten-
dente.* LINN. Sp. 20. P.

FRAGARIA. LINN. 558.
Stamina circiter viginti. Receptaculum pulpo-
sum, molle, esculentum.

1. *Fragaria flagellis reptans, vesca.* LINN. Sp. 17.
Mat. med. 245. HALL. H. 242. *Fragaria vulgaris.*
P. 326. BLACKW. T. 77. Vbiq. in sylvis.
Varietas α . frequentibus in placenta pilis & peduncu-
lis lanuginosis In Saxonia; β . *hortensis* fructu maiori.

GYMNOBOLYSPERMAE.

119

ALCHIMILLA. HALL. LINN. 153.

Aphanes. LINN. 154.

Calyx semioctofidus. Petala nulla. Stamina quatuor. Styli ad basin germinis inserti. Semen unum aut duo in fundo calycis.

Alchimilla foliis integris stellatis polygoniis. HALL. *Helv.* 184. Alchimilla foliis lobatis. (vulgaris.) LINN. Sp. 1. Alchimilla. BLACKW. T. 72. vulgaris. B. Pin. 319. In sylvis. P.

Alchimilla foliis mollioribus semitrilobis, lobis incisis. HALL. *Gott.* p. 31. Aphanes. LINN. *Spec.* 1. Perchepier. HALL. *Helv.* p. 184. In arvis abunde. A.

RVBVS. LINN. 557.

Calyx quinquepartitus. Styli lateri germinis inserti. Acini multi monospermi in baccam compositam congesti.

Rubus foliis ternatis nudis, caule aculeato. HALL. *Gott.* p. 31. LINN. Sp. 4. HALL. *H.* 341. n. 2. Rubus repens, fructu caefio. B. Pin. 479. In agris & dumetis v. g. circa Graezelii fabricas. P.

Rubus foliis ternatis & quinatis, subtus tomentosis, fructu rubro, villosus. HALL. *H.* 1. p. 3. Rubus Idaeus. LINN. Sp. 1. BLACKW. T. 289. Rubus Idaeus a. spinosus β. laetior. B. Pin. 479. In sylvis v. g. circa Pfaff. P.

Rubus foliis ternatis & quinatis, costa spinosa, fructu nigro & levi. HALL. *H.* n. 4. Rubus fruticosus. LINN. Sp. 5. Rubus vulgaris fructu nigro. B. Pin. 479. BLACKW. T. 45. β. Varietas fructu albo. Vbique in dumetis & sylvis. P.

ROSA.

ROS ACULEATA. § 6. HD.

Calyx carnosus, quinque segmentis coronatus. Germina multa in fundo calycis, qui in fructum mollem, coloratum abit.

Rosa aculeata, foliis odoratis subtus rubiginosis. HALL. *Helv.* 359. *eglantaria*. LINN. *Sp.* 2. Rosa sylvestris, foliis odoratis. B. *Pin.* 483. In sterilibus ad vias v. g. verius *Pless.* Fr.

G. Varietas flore flavo, *hera*.

Rosa foliis utrinque villosis, fructu spinoso. HALL. *Fl.* 350. n. 8. *villosa*. LINN. *Sp.* 3. Rosa sylvestris pomifera maior. B. *Pin.* 484. Passim in sepibus. Fr.

Rosa fructu levi, calycis foliolis divisis. HALL. *Fl.* 349. Rosa caule aculeato, petiolis inerimibus, calycibus semipinnatis. (canina.) LINN. *Sp.* 4. Rosa canina sylvestris. BLACKW. *T.* 18. Ubique ad sepes. Fr.

H. Rosa caule aculeato, pedunculis hispida, calycibus semipinnatis glabris. *centifolia*. LINN. *Sp.* 6. Rosa multiplex media. B. *Pin.* 482. Fr.

H. Rosa caule aculeato, pedunculis brevibus, calycibus semipinnatis glabris. *alba*. LINN. *Spec.* 7. Rosa alba vulgaris maior. B. *l. c.* In hortis. Fr.

CLASSIS. VI

POMIFERAE.

ORDO III

Pomiferae umbelliferae.

Petala & stamina, ut in priori classe, ex calyce oriuntur. Germen infra receptaculum floris. Conceptaculum feminum pulposum ex apice calycem persistentem semiquinquesidum praefert.

RIBES. LINN. 247.

Petala calycis segmentis minora. Stamina quinque. Tuba bifida. Bacca unilocularis.

Ribes inermis, racemis glabris pendulis, floribus planiusculis. *rubrum*. LINN. *Sp.* 1. *Mat. med.* 92. HALL. *H.* 346. Ribes *vulgare acidum*. BLACKW. 285. *Johannisbeeren*. Sponte in *Helvetia*, circa *Altdorfium*, & regionibus septentrionalibus. 9. Varietas hortensis fructu rubro & albidio. Fr.

Ribes inermis floribus planiusculis, racemis erectis. HALL. *l. c.* n. 2. Ribes *alpinum*. LINN. *Spec.* 2. Ribes *alpinum dulce*. 1. B. *Hist.* 2. p. 98. In sylvis *Heterophicis* super *Stollberg*. Fr.

Ribes inermis, foliis & nigro fructu maximis, olentibus. HALL. *Helv.* 347. n. 4. Ribes *inermis, racemis pilosis, floribus oblongis. (nigrum.)* LINN. *Sp.* 4. *Mat. med.* 100. *Grossularia non spinosa, fructu nigro*. B. *Pin.* 455. In pago *Rosdorf* ad rivum. Fr.

Ribes ramis aculeatis erectis, fructu glabro. LINN. *Cliff.* 82. HALL. *H.* 346. n. 1. Ribes *vna crispa*. LINN.

LINN. Sp. 7. *Grossularia sylvestris* A. B. BLACKW.
T. 277. Variet. *bort.* Sponte rarius ad sepes. Fr.

MYRTVS. LINN. 543.

Calyx erectus. Tuba simplex. Bacca trilocularis.

1. *Myrtus* floribus solitariis involucro diphyllis.
communis. LINN. Sp. 1. Mat. med. 224. *a.* *Myrtus*
minor vulgaris. Tarentina. *β.* *Myrtus latifolia* Ba-
tica. B. Pin. 469. BLACKW. 114. In Europa au-
strali. Fr.

SORBUS. HALL. LINN. 548. *Crataegus* L. 547.
Mespilus. L. 549.

Stamina viginti circiter. Tuba duae, tres, quin-
que, & totidem semina in bacca rotunda umbili-
cata.

1. *Sorbus* foliis pinnatis utrinque glabris fructu mi-
nimo. *aucuparia*. LINN. Sp. 1. HALL. Helv. 259.
Sorbus sylvestris. Tab. 1020. *Sorbus terminalis*.
BLACKW. 173. *Vogelbeerbäum*. Ubique in sylvis.
Tubae tres. Fr.

1. *Sorbus* (*Crataegus*) folio septangulo, subtus hir-
suto. HALL. H. 354. *Crataegus foliis cordatis acu-
tis, laciniis acutis serratis*. LINN. Cliff. 187.
terminalis. LINN. Sp. 2. C. B. & varior. auctorum.
In sylvis passim. Tubae duae. Fr.

1. *Sorbus* (*Crataegus*) spinosa, foliis obtusis, sae-
pe trifidis. HALL. H. 354. *Crataegus foliis obtusis,
bifidis, serratis*. (*Oxyacantha*). LINN. Sp. 8. Me-
spilus *sylv. apii folio, spinosa*. B. Pin. 454. *Spina*
alba.

alba. BLACKW. T. 149. Ad sepea. Tubae duae & tres. Fr.

H. Sorbus (*Mespilus*) *incanis*, foliis lanceolatis integerrimis subtus tomentosis, calycibus acuminatis. *Germanica*. LINN. *Sp.* 1. *Mespilus Germanica folia laurino non serrato*. B. *Pin.* 453. *Mespilus* BLACKW. 154. In hortis hic colitur. Tubae quinque. Fr.

H. Sorbus (*Mespilus*) foliis ovatis integerrimis. *Cotoneaster*. LINN. *Spec.* 6. HALL. *H.* 352. n. 2. *Chaenomespilus Gissleri*. CLVS. *Hist.* 1, p. Tubae duae, tres, & quinque. Fr.

PYRVS. LINN. 550. *Malus*. T. 406. *Cydonia*. 405.

Stamina viginti. Tubae quinque. Pomum carnosum in quinque loculis membranaceis semina oblonga fover.

Pyrus foliis ferratis, pomis basi productis. LINN. *Sp.* 1. HALL. *H.* 351. *Pyrus sylvestris*. B. *Pin.* 439. BLACKW. 453. In sylvis. Fr.

β. *Pyri domesticae* varietates *bortenses* plurimae.

Pyrus foliis ferratis, pomis basi concavis. *Malus*. LINN. *Sp.* 2. HALL. *H.* 351. *Malus sylvestris*. B. *Pin.* 435. BLACKW. 178. In sylvis. Fr.

β. *Mali domesticae* varietates *bortenses*.

H. *Pyrus* foliis integerrimis, (calyce ferrato). *Cydonia*. LINN. *Sp.* 4. *Malus Cotonea*. Fr.

PUNICA. LINN. 544. T. 407.

Calyx coloratus. Tuba simplex. Pomum magnū nonem loculis instructum.

H. *Punica. Granatum.* LINN. *Sp.* 1. *Malus Punica.* B. *Pin.* 438. *Granata.* BLACKW. 145. *Balaustia fl. pleno* B. l. c. BLACKW. 97. In Europa australi. Fr.

PSIDIUM. LINN. 541. Guaiava.

T. 443.

Calycis segmenta ovata. Stylus subulatus, longissimus. Bacca maxima villocularis.

H. *Psidium. Guaiava.* LINN. *Sp.* 1. *Psidium roseis tetragonis.* H. *Vpf.* 122. *Guaiava alba a. acida, & dulcis.* In India. Fr.

ORDO. II.

Pomiferae non umbilicatae.

Omnia priora. Sed ovarium intra calycem cavitate intermedia remotum, deciduum. In fructu carnoso nux villocularis. Tuba simplex.

PRUNUS. LINN. 546. Armeniaca. Cerasus.

Laurocerasus. Padus. T.

Calyx campanulatus. Stylus filiformis. Fructus oblongus aut subrotundus simili nuce.

1. *Prunus spinosa* foliis lanceolatis. LINN. *Sp.* 10. HALL. *Helv.* 355. *Prunus sylvestris.* B. *Pin.* 444. BLACKW. T. 305. Vbique in collibus aprieis. Fr.

H. *Prun-*

h. *Prunus pedunculis simplicibus foliis lanceo-*
lato-ovatis deciduis, domestica. LINN. Sp. 9.
Cliff. 186. *Mat. med.* 232.

g. *Pruni domesticae varietates hortenses.* In hac
 & priori fructus ovalis. Fr.

Prunus pedunculis simplicibus, foliis ovato-lan-
ceolatis, conduplicatis, Cerasus. LINN. Sp. 8. *Cera-*
sus foliis ovato lanceolatis. HALL. *Helv.* 156. n. 1.
Cerasus sylvestris. C. B. Sponte in sylvis. Fr.

g. *Cerasi hortensis varietates.* Fructus subro-

undus.
 h. *Prunus floribus sessilibus, foliis subcordatis.*
Armeniaca. LINN. Sp. 6. *Mala Armeniaca.* B. *Pin.*
 482. Fr.

h. *Prunus floribus racemosis, foliis semperviren-*
tibus dorso biglandulosis. Laurocerasus. LINN. Sp.
 4. *Padus sempervirens, foliis sessilibus.* HALL. *Germ.*
 112. *Laurocerasus.* CLVS. *Hist.* 1. p. 4. Ex *Optim-*
is ad nos advenit. Fructus subrotundus, calyx non
 deciduus. Fr.

Prunus floribus racemosis, foliis deciduis, basi
subtus biglandulosis. Padus. LINN. Sp. 1. *Padus*
HALL. H. 357. *Cerasus racemosa sylvestris, fructu*
non eduli. I. R. H. *Elzbeeren.* Circa *Northheim, No-*
ribergam, & in *Helvetia.* Fructus & calyx similis
 priori. Fr.

AMYGDALVS. LINN. 548. *Perfica.* T. 400.

Habitus prioris. Fructus subrotundus, villosus,
 Nux compressa, sulcis reticulata.

H. *Amygdalus foliorum serraturis omnibus acutis. Persica.* LINN. *Sp.* 1. *Persica Malus.* BLACKW. *T.* 101. B. *Pin.* 440. *Pferfigbaum.* Fr.

ORDO. III.

Pomiferae Hesperides.

CITRVS. LINN. 807. *Aurantium.* Citreum.
Limon. TOURN. 393-397.

Calyx minimus, quinquedentatus. Stamina multa filamenta inter se connata, ex thalamo orta. Stylus stigmate globoso. Fructus pulposus, novemlocularis.

H. *Citrus petiolis linearibus. Medica.* LINN. *Sp.* 1. ROYEN. 266. *Malus Medica.* B. *Pin.* 435. *Citria Malus.* BLACKW. *T.* 361. Fr.

H. *β. Limon vulgaris.* FERRAR. *Hesper.* 193. *Limonia Malus.* BLACKW. 362. Fr.

H. *Citrus petiolis alatis. Aurantium.* LINN. *Sp.* 2. *Malus Aurantia.* B. *Pin.* 346. *Aurantia* BLACKW. *T.* 349. Omnes sponte in vtraque India. Fr.

CLASSIS. VII.

MULTILOCVLARES.

Similes prioribus numero staminum, petalis liberis, patentibus. Fructus autem est capsula sicca, aut unilocularis, aut multilocularis.

PHILADELPHVS. LINN. Sp. 440. Syringa.

T. 389.

Petala quatuor. Tuba quadrifido stigmate. Capsula quadrilocularis calyce cincta.

H. *Philadelphus foliis subdentatis. coronarius.* LINN. *Sp. 1. Syringa alba.* f. *Philadelphus Athenæi.* I. R. H. B. *Pin. 399. Frutex coronarius.* CLVS. *Hist. 1. p. 55. Ad sepes & in hortis, quasi spontanea.* Fr.

CAPPARIS. LINN. 567.

Calycis folia quatuor, totidemque petala maxima. Capsula carnosâ unilocularis.

H. *Capparis aculeata. spinosa.* LINN. *Spec. 1. Mat. med. 259. Capparis spinosa, fructu minore, folio rotundo.* B. *Pin. 480. In Europa Australi.* Fr.

TILIA. LINN. 587. HALL. 357.

Calycis folia quinque colorata, totidemque petala. Tuba stigmate pentagono. Capsula quinquelocularis, sed unicum semen plerumque adolescit.

Tilia floribus nectario destitutis. LINN. *Sp. 1.* *Tilia foliis amplioribus, mollibus subhirsutis.* HALL. *Helv. 357. Tilia femina, folio maiori.* B. *Pin. 426.*

β. *Tilia folio glabro, duriori, minori.* HALL. *l. cit. n. 2. Tilia femina folio minori.* BAYH. *l. c. l. sylvis & ad vias.* Fr.

CORCHORVS. LINN. 599.

Calycis folia & petala quinque. Tuba crassa. Sin-
guâ cylindræa, quinquelocularis, maxima.

H. *Corchorus capsulis oblongis, ventricosis, Toloriorum infimis serraturis setaceis reflexis. olitorius.* LINN. *Sp.* 2. *Corchorus* PLINII. R. P. 317. CAMER. *Hort.* 47. T. 12. In *Asia, Africa, America.* A.

CISTVS. LINN. 598.

Calyx quinquefolius, duo folia alterna minora. Petala quinque fugacia. Stylus stigmate orbiculato. Capsula subrotunda.

H. *Cistus arborefcens, foliis lanceolatis supra laevibus, petioliis basi coalitis vaginantibus. Ladanifera.* LINN. *Sp.* 3. *Cistus ladanifera hispanica incana.* R. Pin. 467. *Cistus Ledon* L. *angustifolius.* CLVS. *Hist.* I. p. 77. In *Hispania.* Fr.

H. *Cistus arborefcens, foliis lanceolatis, sessilibus, vtrinqve villosis, trinerviis, alis nudis. Monspeliensis.* LINN. *Sp.* 5. *Cliff.* 205. *Cistus Ledon.* V. CLVS. *Hist.* 1. p. 79. *Cistus ladanifera Monspeliensium.* R. Pin. 467. Folia ficiora, ac in sequente, margine reflexa, inferne exalbida, vix pilosa, vilcida, odore forti, basi vaginantia, nervis lateralibus per maximam folii longitudinem decurrentibus. Petioli & calycis folioli pilis albis dense oblita. In *Gallia Narbonensi & Hispania.* Fr.

H. *Cistus arborefcens, foliis sessilibus basi coalitis vtrinqve villosis, rugosis, ellipticis. Cistus Ledon* IV. CLVS II p. 78. nostram plantam exacte refert, est *Cistus hirta.* LINN. *Cent.* I. n. 37., quo ille iconem *Glusianam* revocat, a nostra sit diversissima. Num *Cistus* HALL. *Goit.* n. 2.? Videtur. Folia vtrinqve pilosa, sine obtuso, qualia CLVSIVS pingit,

git, molliora ac in priori, viscida, nervis lateralibus vix ad dimidiam longitudinem folii decurrentibus. Fr.

H. *Cistus arborescens* foliis sessilibus utrinque villosis, rugosis, obverse ovatis, margine undulatis. *Cistus max. folio rotundo hirsutissimo.* C. B. P. *Pin.* 465. HÆLL. *Gott.* 114. *Cistus mas. folio rotundior.* L. B. *Hist.* II. pl. 21. Leon, ad quam DV. HAMMEL, *Traité des arbres & arbustes* T. 1. T. 64. Synonymum, I. p. citat, nostrae plantae egregie convenit, nisi quod petioli iusto longiores pingantur. Folia basi coalita, cum ramulis villosa, subtus incana, ad dimidium lata ac longa, rugosa, trinervia, neque viscida, ut in prioribus. Flos purpureus, solitarius. Fr.

H. *Cistus arborescens*, foliis petiolatis, cordato-subrotundis, mollibus, utrinque tomentosis, crenatis. Num. *Cistus mai.* III. CLVS. *Hist.* I. p. Icon DV. HAMMEL, *lib. cit.* T. 68. sic satis ob figuram foliorum ad nostram accedit. Arbuscula trunco firmiori. Folia tactu villi sericei instar pubescentia ad insertionem petioli hamulis sagittata. Ramuli incani. Nondum floruit. Cistum totus habitus prodit. Fr.

Tres species hic omissas vide sub fine libri.

HYPERICVM. LINN. 808.

Calyx quinquepartitus, quinque petala. Filamenta staminum multa, saepe coalita. Capsula subrotunda, tot loculis, quot styli adsunt.

Hypericum floribus trigynis, caule ancipiti, foliis obtusis pellucido-punctatis. *perforatum.* LINN. *Sp.*

34. HALL. *H.* 360. n. 1. *Hypericum vulgare*. B. *Pin.* 279. *Hypericum*. BLACKW. *T.* 15. Vbiq; in agris & sylvis. P.

Hypericum humifusum, foliis glabris perforatis, punctis nigris in margine notatis. HALL. *l. c.* n. 2. *Hypericum floribus trigynis, caulibus encipitibus prostratis*. LINN. *Sp.* 15. *Hypericum minus supinum*. B. *Pin.* 279. I. R. H. In atenosis v. g. M. Hagen; jenseit Rheinhausen im Wald. P.

Hypericum foliorum ora punctata, calycibus ferrato-glandulosis. HALL. *H.* 361. n. 4. *Gott.* 115. *Hypericum hirsutum*. LINN. *Sp.* 16. *Var. a.* Ascyton. f. *Hypericum bifolium glabrum, non perforatum*. B. *Pin.* 280. *Androsæum Campoclarense*. COLUMN. *Espbr.* 73. *T.* 74. Tubae tres. Circa Rheinhausen, in monte ad Borsenden, ad viam sub M. Pleß. In M. Altenßottberg. P.

Hypericum villosum, calycibus ferrato-glandulosis, foliis perforatis. HALL. *H.* n. 5. *Gott.* 116. *Hypericum* LINN. *Sp.* 16. *Var. β.* *Hyp. villosum erectum, caule rotundo*. I. R. H. *Androsæum alterum hirsutum*. COL. *l. c.* p. 75. Rarius ad sepes. P.

Hypericum floribus trigynis, calycibus ferrato-glandulosis, foliis cordatis glabris, pulchrum. LINN. *Sp.* 18. *Hypericum calyce ferrato, foliis connatis*. HALL. *Gott.* 116. *Helv.* 3. *Hyp. pulchrum* Tragi. I. B. III. p. 383. In sylva super Hardeggen. P.

Hypericum caule quadrangulo, foliis perforatis. HALL. *H.* n. 6. *Gott.* 116. *Hyp. quadrangulum*. LINN. *Sp.* 13. I. R. H. In fossis. P.

h. *Hypericum floribus trigynis, staminibus corolla longioribus, caule ancipiti fruticoso. bircinum.* LINN. *Sp.* 8. *Hypericum foetidum frutescens.* 1. R. H. DILL. *Eltb.* 182. *T.* 151. In *Europa australi*. Fr.

h. *Hypericum floribus trigynis, fructu baccato, caule fruticoso ancipiti. Androsæmum.* LINN. *Sp.* 5. *Androsæmum maximum frutescens.* B. PIN. 280. *Hypericum bacciferum.* BLACKW. *T.* 94. In *Anglia, Italia.* Fr.

h. *Hypericum floribus pentagynis, caule fruticoso, foliis ramisque cicatrilatis. Bulgaricum.* LINN. *Sp.* 1. *Myrto-Cistus Pennæi.* CLVS. *Hist.* 1. p. 68. In *Maiorca.* Fr.

CLASSIS. VIII.

COLVMNIFERAE.

Calyx sæpe duplex, interior quinquesfidus. Petala quinque obcordata, vnguibus connata cum tubo, ex filamentis staminum coalitis composito. Stylus per tubum stamineum adscendens fissus in tot stigmata, quot loculi in fructu circa axin. Omnibus vires emollientes.

HIBISCVS. LINN. 756. *Ketmia.* T. 26.

Calyx exterior multífidus, interior quinquesfidus. Stigmata quinque crassa. Fructus loculi quinque, polyspermi.

h. *Hibiscus foliis ovato-lanceolatis; superne inciso-serratis, caule arboreo. Syriacus.* LINN. *Sp.* 9.

Alcea arborescens. CAMER. H. T. 1242. *Syracca*. P. Pin. 316. In *Syria*. Fr.

H. *Hibiscus aculeatus*, foliis trilobis & quinquelobis, inaequaliter serratis. HALL. Gott. 117. *Hibiscus*. LINN. Sp. 13. Var. d. *Ketmia Indica*, foliis digitatis, flore magno sulphureo, umbone atropurpureo, petiolis spinosis. EHRET. T. 6. f. 1. *Alcea Benghalensis*. COMM. H. L. p. 35. T. 18. In *India*. A.

H. *Hibiscus foliis palmato-digitatis septempartitis*. *Manibot*. LINN. Sp. 13. *Ketmia folio Manibot serrato*, flore amplo sulphureo. DILL. Elth. 189. T. 156. f. 189. In *India*. A.

H. *Hibiscus foliis palmatis serratis*, fructu pyramidalis, longissimo. HALL. G. 118. An *Ketmia Zeylanica Fici folio*, perianthio oblongo integro. DILL. Elth. T. 157. f. 1201. *Hibiscus ficulneus*. LINN. Sp. 10. Folia conveniunt, sed in nostra flores non mutant. An potius huic referenda *Alcea Americ. annua flore flavo*, fructu pyramidalis sulcato. COMM. H. Amst. 1. p. 537. T. 19. *Hibiscus foliis quinquepartito-pedatis*, calycibus interioribus latere rumpentibus. LINN. Sp. 15. Caulis undique asper aculeis sparsis mitibus. Folia inferiora triloba, superiora quinqueloba, lobis serratis latioribus. Flos priori multo minor simili colore. Calyx interior spathaeformis brevissime quinque-dentatus. Eruus pyramidalis undique strigosus & aculeis echinatus. A.?

H. *Hibiscus foliis subpeltato-cordatis septemangularibus serratis hispidis*. *Abelmoschus*. LINN. Sp. 14. *Ketmia Aegyptiaca*, femine meschata. J. B. H.

100. EVRM. Ceyl. 194. Abelmöschvulgo, HERM. Lagab. 25. In India. P.

H. Hibiscus foliis imis cordato-lanceolatis, superioribus pentagonis serratis. HALL. Gott. 118. Hibiscus foliis inferioribus cordatis angulatis, superioribus subhastatis, floribus subnatis, pistilla cernuo. (Pentacarpos.) LINN. Sp. 19. cum descr. Ketmia palustris minor. ZANICH. Venet. 155. T. 91. In paludosis Venetiae. Corolla purpurea. P.

H. Hibiscus foliis tripartitis incis. calycibus inflatis. Trionum. LINN. Spec. 20. Aloea vescaria. B. Pin. 317. Ketmia vescaria Africana. TÖVRN. 1. R. H. 101. In Italia, Africa. A.

ALTHAEA. HALL. LINN. 749.

Alcea. LINN. 750.

Calyx duplex, exterior in Althaea inaequaliter novemfidus, in Alcea semifidus. Capsulae monospermae, fructum depressum faciunt.

H. Althaea foliis simplicibus tomentosis. officinalis. LINN. Sp. 1. Althaea Dioscoridis & Plinii. B. Pin. 315. Althaea BLACKW. T. 90. In subhumidis. v. g. ad fluvium Vnstrut. P.

H. Althaea foliis scabris, acutis, semitrilobis. HALL. Gott. 119. Althaea foliis inferioribus palmatis, superioribus digitatis, caule fruticoso. (cannabina.) LINN. Sp. 2. B. Pin. 316. In Hungaria, Italia. P.

H. Althaea foliis sinuato angulosis. HALL. l. c. (Alcea) rosea. LINN. Spec. 1. Malva rosea folio

subrotundo. B. Pin. 315. *bortensis*. DOD. Pompr. 652. floribus varii coloris. In *Hungaria*, *Palæstina*. B.

H. *Althaea foliis palmatis*, HALL. l. c. (*Alcea*) *ficifolia*. LINN. Sp. 2. *Malva rosea bortensis*, *folio ficus*. B. l. c. I. R. H. floribus varii coloris. B.

MALVA. HALL. LINN. 751.

Lavatera. LINN. 752.

Calyx in *Lavateris* semitrifidus, in *Malvis* trifolius. Fructus prioris.

H. *Malva caule arboreo*, foliis palmatis, obtusis, plicatis. HALL. Goss. 120. *Lavatera caule arboreo*, foliis septemangularibus tomentosis plicatis, pedunculis confertis axillaribus. LINN. Sp. 1. *Malva arborea Veneta dicta*, parvo flore. B. Pin. 315. Ad ripas Maris mediterranei. Fr.

H. *Malva (Lavatera) foliis quinqueangularibus*, acutis crenatis, lacinia media productiore. LINN. Cliff. 348. HALL. G. 120. *Lavatera caule fruticosa*, foliis quinquelobo-bastatis. (*Olbia*.) LINN. Sp. 3. *Althaea frutescens*, folio acuto, molli, virente, flore specioso. PLVKNET. T. 8. f. 1. *Althaea arborea olbia* in *Gallo Provincia*. LOB. ic. 653. Pedunculi axillares solitarii brevissimi. Ad ripas maris Mediterranei circa *Monspesium*. Fr.

H. *Malva caule arboreo*, foliis tomentosis, obtusissimis micantibus. HALL. l. c. *Lavatera micans*, LINN. Spec. 2. *Malva foliis mollibus undulatis*, superius micis sulphureis ad solem splendentibus demaris. MORIS. Hist. 1. p. 523. S. 5. T. VI. f. 9. Folia

filamentosa, subrotunda, plicata. Scipulae magnae ovatae. Pedunculi axillares solitarii, petiolis breviores. Calyx exterior trifolius. Habitus prioris. In *Hispania*. Fr.

H. Malva (*Lavatera*) caule fruticoso, foliis subcordatis subtrilobis, rotundatis crenatis, stipulis cordatis, pedunculis unifloris. *triloba*. LINN. Sp. 4. *Althaea frutescens, folio rotundiore incano*. B. Pin. 316. Pedunculi ares pluresve ex alis foliorum, petiolis breviores. Scipulae cordatae. Calyx exterior trifolius. In *Lusitania*. Fr.

H. Malva (*Lavatera*) foliis inferioribus cordato-subrotundis, superioribus lobatis crenatis, calycibus utrinque incis, caule herbaceo. ROY. Lugdb. 355. *Lavatera Thuringiaca*. LINN. Sp. 6. *Althaea Thuringiaca grandiflora*. DILL. Elth. 9. T. 8. f. 8. CAMER. Hort. 12. T. 6. In *Thuringia, Pannonia*.

H. Malva foliis subrotundis, capsulis scuto subpositis. HALL. l. c. *Lavatera foliis glabris, caule scabra herbacea, pedunculis unifloris, fructibus orbiculo testis. (trimestri.)* LINN. Sp. 8. *Lavatera foliis subcordatis obtusis angulatis serratis, caule herbacea*. ROY. 355. *Malva folio vario*. B. Prodr. T. 137. flore rubro & albo. In *Hispania, Gallia Narbonensi*. A.

H. Malva caule erecto, foliis semiorbiculatis crenatis, calyce exteriore diphylo. *Hispanica*. LINN. Sp. 11. *Malva rotundifolia glabra ex Hispania, flore amplo*. PLVKNET. Alm. 238. T. 44. f. 3. flore rubro & albo. In *Hispania*. A.

4. *Malva caule erecto, foliis pene palmatis obtuse serratis.* HALL. *Helv.* 362. *Malva caule erecto herbacea, foliis lobatis obtusis, pedunculis petiolisque pilosis.* (sylvestris.) LINN. *Sp.* 9. *M. sylvestris, folio sinuato.* D. PIN. 314. *vulgaris.* BLACKW. *T.* 22. Vbique in gramineis. A.

5. *Malva caule repente, foliis cordato-orbiculatis obsolete quinquelobis. rotundifolia.* LINN. *Sp.* 8. HALL. *H.* 363. *Malva sylvestris folio rotundo.* D. PIN. 314. Vbique ad vias. A.

6. *Malva floribus albis, parvis, conglomeratis.* HALL. *Gorr.* 107. *Malva caule erecto, foliis angulatis floribus axillaribus glomeratis.* (verticillata.) LINN. *Sp.* 12.

7. *Malva foliis crispis.* C. B. 315. MORIS. *S. V. T.* 15. In Syria. A.

8. *Malva humifusa, minima* SHERARDI. *TILLIE Hort. Pis.* *T.* 38.

9. *Malva foliis subcordatis laciniatis glabris, caule arborecente.* *Capensis.* LINN. *Sp.* 6. *Malva capensis frutescens grossulantes folio maiori hirsuta.* DILL. *Elab.* 209. *T.* 169. f. 207. In Africa. Fr.

10. *Malva caule erecto, foliis multipartitis scabriusculis.* *Alcea.* LINN. *Spec.* 13. HALL. *H.* 363. n. 3. *Alcea vulgaris maior.* C. B. 315. I. R. H. BLACKW. *T.* 309. Ad viam versus Bovenden in dumetis, ad rivum prope Lengtern. P.

11. *Malva foliis palmatis scabris, caule tomentoso fruticoso, pedunculis multifloris.* *brionifolia.* LINN.

LINN. Sp. 5. *Althaea frutescens, hyemalis foliis* in Prodr. 138. In Hispania. Er.

H. Malva caule erecto, foliis imis subrotundis, superioribus tenuiter divis. HALL. Pl. 364. Malva foliis radicalibus unguiformibus incis, caulibus quinquepartitis pinnato-multifidis, moschata. LINN. Sp. 14. Malva montana, f. *Alcea rotundifolia lapinaria* COL. Ecphr. 1. p. 148. T. 147. In Asia, Helvetia. Italia. P.

H. Malva foliis palmatis dentatis, corollis calycibus aequalibus. An Malva aegyptia. LINN. Sp. 5. descr. Pedunculi saepe biflori in nostra semper foliis longiores, ex alis solitarii, in summitate plures congesti. Corollae coerulescentes, calyci aequales. *Alceam minorem maritimam tenuifoliam pro-cumbentem* HERM. Parad. 9. T. 9. quae est Malva Tournefortiana. LINN. Cent. 1. n. 63. huc referrem ob figuram foliorum, nisi corollae calyce dimis maiores pingerentur. A.

SIDA. LINN. 747. Abutilon.

TOURN. 25.

Calyx ut plurimum simplex. Fructus capsula cornutis, cristatis.

H. Sida caule repente, foliis multifidis. Malva Caroliniana LINN. Sp. 7. Abutilon Carolinianum repens, alceae foliis, flore belvolo. DILL. Elth. 5. T. 4. f. 4. MARTYN. Cent. T. 34. Calyx exterior trifolius. Capsulae aristatae. In Carolina. A.

H. Sida foliis cordatis subrotundis, renatis tomentosis, pericarpis inflatis multilocularibus crenatis

tis repandis, *crispa*. LINN. Sp. 8. *Abutilon vesicarium crispum*, floribus melinis parvis. DILL. *Elth.* 6. T. 5. f. 2. In Carolina, Bahama.

H. Sida foliis cordato-lanceolatis integerrimis, *periplocifolia*. LINN. Sp. 5. *Abutilon periplocacutius* folio, fructu stellato. DILL. *Elth.* 4. T. 3. f. 3. In Ceylona.

H. Sida foliis inferioribus cordato-angulatis, superioribus hastatis, petalis integerrimis. ROYEN. *Lugd.* 349. LINN. *Vps.* 199. *Abutilon americanum*, flore coeruleo. HALL. *Gott.* 122. Caules debiles procumbentes. Anguli foliorum inprimis superiorum acuti, ora fusca, rarius dentata. Stigmata septem. A.

H. Sida foliis crenatis, inferioribus dentatis obtusis, superioribus acuminatis. ROYEN. 349. Sida *cristata*. LINN. *Spec.* 9. *Abutilon Lavaterae* flore, fructu cristato. DILL. *Elth.* 3. T. 2. f. 2. Caules erecti. Folia omnia crenata, inferiora obtusiora, subcordata, superiora saepius hastata, angulis posterioribus rotundatis. Flos priori multo maior. Stigmata viginti. Planta a priori diversissima. A.

H. Sida foliis subrotundo-cordatis acuminatis crenatis. *Abutilon*. LINN. Sp. 7. *Althaea Theophrasti*, flore luteo. B. Pin. 316. In Indiis. A.

H. Sida foliis ovatis serratis, floribus ad alas conglomeratis. *Abutilon Vlmifolium*, floribus conglomeratis ad foliorum alas. HALL. *Gott.* 123.

H. Sida foliis lanceolato-rhomboidibus serratis. *rhombifolia*. LINN. Sp. 2. *Malvinda unicornis*, folio
lio

to rhomboid, pinnatis. DILL. Elb. 346. T. 1715 f. 312. Althaea carpi folio, fuscis latis. BOCC. St. 11. T. 6. f. 3. Pedunculi petiolis longiores. A.

H. Sida foliis cordato-ovatis serratis; stipulis serratis, axillis subspinosis. *spinea. LINN. Sp. 1. Malvinda bicornis, sanguisorbae folio villosa & glabra. DILL. Elb. L. 14. T. 171. f. 219. Pedunculi petiolis fere breviores. A.*

H. Sida foliis inferioribus subrotundis, superioribus trilobis, crenatis, flosculis exiguis aggregatis, folio spathaeformi exceptis. Ita interim definire liceat plantam, cuius flores plures in involucrio spathaceo compresso aggregati latent. Corolla flava calyce piloso paulo maior. Ad utrumque latus singuli floris adstant duae ligulae lineares, capillaceae. Stamina pauca. Capsulae quinque monospermae; seminibus obesis, minusculis. Vberius hanc plantam alibi proxime describam.

VRENA. LINN. 754.

Calyx exterior semiquinquefidus, interior quinquefolius. Capsulae echinatae.

H. Vrena foliis angulatis. LINN. Spec. 1. Vrena Sinica, Xanthii facie, DILL. Elb. 340. T. 319. f. 412. In China. Fr.

NAPAEA. LINN. 748.

Calyx simplex, teres. Petala oblonga. Stamina plurima. Capsulae muticae.

H. Napaea pedunculis involucriatis angulatis, foliis scabris, floribus dioicis, *dioica. LINN. Sp. 1. Abutilon*

etiam foliis profunde dissectis, palmatis inaequalibus.
Sesuvium G. m. KHAET. To. 7. f. 1. & T. 8. HALL.
 Gert. 224. Flores feminae similibus effusis abon-
 tibus gaudent. In *Virginia*. P.

HERMANNIA. LINN. 742.

Calyx simplex. Stamina quinque. Stigma sim-
 plex. Fructus quinquelocularis.

H. *Hermannia* foliis lanceolatis, obtusis, serratis.
Hysopifolia. LINN. Sp. 4. *Althaea Africana* fru-
 tescent, floribus aureis cochlearis pendulis minori-
 bus. VOLKAM. Norib. T. 24. In *Africa*. Fr.

PENTAPETES. LINN. 757.

Calyx simplex, Stamina decem fertilia, binaria
 alterne posita, cum filamentis quinque sterilibus, co-
 loratis.

H. *Pentapetes* foliis hastato-lanceolatis serratis
phoenicea. LINN. Sp. 1. *Alcea Indica*, lucido hast-
 ato folio, flore blattariae phoeniceo. PLUK. Phyt. T.
 255 f. 3. Al

GOSSYPIUM. LINN. 755.

Calyx duplex. Capsula quadrilocularis. Semina
 lana involuta.

H. *Gossypium* foliis palmatis, lobis lanceolatis,
 caule fruticoso, arboreum. LINN. Sp. 3. *Gossy-
 pium arboreum*, caule laevi. BAVH. Pin. 430. In
India. Fr.

CLASSIS IX.

SVCCVLENTAE.

Caulis & folia succo aequo turgida & carnosa.

ORDO I.

CACTI.

Calyx concavus monophyllus. Corolla in plurima petala profundissime fissa, parieti calycis affixa. Stamina plurima calyci inserta. Fructus carnosus sub calyce.

CACTVS. LINN. 539. Cereus. Opuntia.

Melocactus. Tuna. Peireskia.

Planta aphyllus. Calyx tubulato-cavus. Stylus cylindraceus stigmate multifido. Fructus carnosus, multilocularis. Sola Peireskia foliis instructa.

H. Cactus subrotundus tectus tuberculis, ovatis, barbatis, *mammillaris*. LINN. Sp. 1. Echino-Melocactus *minor lactescens, tuberculis f. mammillis majoribus*. HERM. Parad. 136. T. 136. Ficoides *Americana sphaerica tuberculata lactescens, flore albo*. COMM. H. 1. p. 105. T. 55.

H. Cactus quadrangularis longus erectus, angulis compressis, *tetragonus*. LINN. Sp. 4. H. Vps. 119.

H. Cactus erectus sexangularis longus, *hexagonus*. LINN. Sp. 5. Cereus *altissimus Surinamensis*. HERM. Par. 116.

H. Cactus erectus longus subnovemangularis; angulis obsoletis, spinis lana brevioribus, *lanuginosus*.

L

LINN.

titon folia prof
Femina 5. mar.

Examina 5. mar. 11

Geft. 124. Flor. C -

tivis gaudent. I n n

HER 5.

Calyx simple
plex. Fructus

H. Hermann:

Hyssopifolia. L.
rescens, floribus. VOLKAM

rescens, floribus

bus. VOLKAM

PE-

Calyx simp
alterne posita
loratis.

alterne poſiſta

Lexatis, Inc.

н. Рентарс

phoenicea. L. I.

to folio, flore

255 f. 3. A

G.

Calyx dup
lana involuta.

lana involuta.

Н. Гофурі

caule fruticoso

pium arboreum

India. Fr.

19.

is indica *f. Opuntia Curassavica minima.*

H. 1. p. 107. T. 56.

actus prolifer ensiformi-compressus serrato-
 aus. *Phyllanthus*. LINN. Sp. 20. Cactus foliis
 nibus obtuse serratis. H. Cliff. l. c. *Cereus scolo-*
folio brachiato. DILL. *Eltb.* 73. T. 64. f. 74.

Cactus caule tereti arboreo; spinoso, foliis
 olato-ovatis. *Peireskia*. LINN. Sp. 21. *Peire-*
sculeata, flore albo, fructu flavescente. DILL.
 305. T. 227. f. 294.

ESEMBRYANTHEMUM. LINN. 552. DILL.

Eltb. p. 159. Ficoides TOURN.

Calyx semiquinquefidus carnosus. Fructus carno-
 us tot loculis, quot styli adfunt. Folia in plerisque
 opposita.

H. *Mesembryanthemum* foliis alternis ovatis obtu-
 is undulatis. *crystallinum*. LINN. Sp. 2. *Mesembryan-*
themum crystallinum, plantaginis folio undulato (mi-
cus argenteis adperso.) DILL. *Eltb.* 231. T. 180. f. 211.

H. *Mesembryanthemum* acaule, foliis linguiformi-
 us altero margine crassioribus. *linguiforme*. LINN.
 p. 34. *Mesembr. folio linguiformi latiore.* DILL.
Eltb. T. 184 f. 224. Corolla flava.

H. *Mesembryanthemum* foliis acinaciformibus
 omatis, angulo carinali scabris, ramis angulatis.
Mesembr. acinaciforme, flore amplissimo purpureo.
 DILL. *Eltb.* 1. f. 271.

β. *Mesembryanthemum falcatum maius, flore*
amplissimo. DILL. T. 212. f. 271.

H. Mesembryanthemum foliis acinaciformibus distinctis laevibus, ramis teretibus. LINN. Sp. 26. Mesembr. *falcatum*, *minium*, *flore purpureo parvo*. DILL. *Elth. T.* 213. f. 276.

H. Mesembryanthemum acaule, foliis semicylindricis connatis externe | tuberculatis. *rostratum*. LINN. Sp. 26. Mesembr. *rostrum ardeae referens*. DILL. *Elth. T.* 186. f. 229.

H. Mesembryanthemum caulescens, foliis subulatis semicylindricis recurvis connatis longis. *loreum*. LINN. Sp. 29. Mesembr. *foliis corniculatis longioribus*. DILL. *Elth. T.* 199. f. 253.

H. Mesembryanthemum foliis alternis subulatis triquetris longissimis. LINN. Sp. 35. Mesembr. *folio pugioniformi, flore amplo stramineo*. DILL. *Elth. T.* 210. f. 269.

H. Mesembryanthemum foliis subcylindricis acutis connatis arcuatis laevibus. *veruculatum*. LINN. Sp. 27. Mesembr. *foliis veruculiformibus, floribus melinis umbellatis*. DILL. *Elth. T.* 203. f. 259.

H. Mesembryanthemum acaule, foliis subteretibus connatis, floribus octogynis. LINN. Sp. 6. Mesembr. *calamiforme*. DILL. *Elth. T.* 186. f. 228.

H. Mesembryanthemum *foliis dolabriformibus*. LINN. Sp. 32. DILL. *Elth. T.* 191. f. 237. Corolla intus aurea micans, extus aurantia.

H. Mesembryanthemum foliis subulatis triquetris strictis acutis: punctis pellucidis obsoletis sparsis. *glaucum*. LINN. Sp. 28. Mesembr. *scabrum, flore sulphureo convexo*. DILL. *Elth. T.* 196. f. 248.

H. Me-

H. Mesembryanthemum acaule, foliis triquetris linearibus apice trifariam dentatis. LINN. Sp. 28. Mesembr. *bellidiflorum*. DILL. *Elth. T.* 189. f. 233.

H. Mesembryanthemum caulescens, foliis deltoidibus triquetris dentatis. LINN. Sp. 9. Mesembr. *deltoides* & *dorso* & *lateribus muricatis*. DILL. *Elth. T.* 195. f. 246.

β. Mesembryanthemum deltoides, foliis non dorso, sed lateribus muricatis. DILL. *Elth. T.* 195. f. 243.

H. Mesembryanthemum foliis linearibus obsolete triquetris distinctis: summis imbricatis laevibus. *splendens*. LINN. Sp. 25. Mesembr. *foliis confertis splendentibus, flore pallido*. DILL. *Elth. T.* 204. f. 260.

H. Mesembryanthemum foliis planis oppositis ovatis acuminatis connatis integerrimis. *tortuosum*. LINN. Sp. 39. Mesembr. *tortuosum, foliis semper-vivi expansis*. DILL. *Elth. T.* 182. f. 223. β. Simile prioris sed foliis congestis. *T.* 181. f. 222.

H. Mesembryanthemum articulis caulinis terminatis in folia acuminata subtus dentata. *uncinatum*. LINN. Sp. 14. Mesembr. *perfoliatum, foliis maiaribus triacanthis*. DILL. *Elth. T.* 193. f. 240.

H. Mesembryanthemum foliis semicylindraceis, calycibus quadrifidis. *noctiflorum*. LINN. Sp. 4. Mesembr. *noctiflorum, flore intus candido, extus stramineo odoratissimo*. DILL. *Elth. T.* 206. f. 263. Styli quatuor.

H. Mesembryanthemum foliis apice barbatis. *barbatum*. LINN. *Sp.* 10. Mesembr. *radiatum, humile, floribus minoribus*. DILL. *Eltb.* T. 190. f. 235.

β. Mesembryanthemum radiatum, ramulis prolixis, decumbentibus. DILL. *Eltb.* T. 190. f. 234.

γ. Mesembryanthemum radiatum, humile, foliis maioribus. DILL. *Eltb.* T. 190. f. 236.

H. Mesembryanthemum foliis semiteretibus, floribus sessilibus axillaribus. *geniculiflorum*. LINN. *Sp.*

3. Mesembr. *Capense geniculiflorum, Neapolitanum creditum*. DILL. *Eltb.* T. 205. f. 261.

H. Mesembryanthemum foliis subtriquetris scabris, corollis bicoloribus. *bicolor*. LINN. *Sp.* 22.

H. Mesembryanthemum tenuifolium, frutescens, flore croceo. DILL. *Eltb.* T. 202. f. 258. Petalainthus flava, extus rubella.

H. Mesembryanthemum foliis subulatis semiteretibus glabris internodio longioribus. *tenuifolium*. LINN. *Sp.* 17. Mesembr. *tenuifolium procumbens, flore coccineo*. DILL. *Eltb.* T. 201. f. 256.

AIZOON. LINN. 553. Ficoidea. DILL.

Calyx quinquepartitus. Corolla nulla. Stamina per phalanges calyci inserta.

H. Aizoon foliis cuneiformi-ovatis, floribus sessilibus. *Canariense*. LINN. *Spec.* 1. *Kali aizoides Canariense, procumbens, portulacae foliis, aspergine rorida perpetuo madidis*. VOLK. *Norib.* 236, T. 236. A.

n. Aizoon foliis lanceolatis, floribus sessilibus. *Hispanicum*. LINN. *Sp.* 2. Ficoidea *Hispanica annua*, folio longiore. DILL. *Elth.* T. 117. f. 143. A.

ORDO. II.

Petalis liberis & staminibus receptaculo insertis.

PORTULACA. LINN. 531.

Calyx bifolius. Petala quinque. Germen sub flore. Stigmata quinque. Capsula ovata, unilocularis.

n. *Portulaca* foliis cuneiformibus, floribus sessilibus, *oleracea*. LINN. *Spec.* 1. HALL. *Helv.* 392. *Portulaca sylvestris*. ROB. *ic.* 388. In *Europa australi*, v. g. ad ripam *Rheni*; in vtraque *India*. A.

n. *Portulaca* foliis subulatis alternis, axillis pilosis, floribus sessilibus. *pilosa*. LINN. *Sp.* 2. *Portulaca Curassavica*, lanuginosa procumbens, HERM. *Parad.* T. 215. COMM. H. I. p. 2. T. 5. In *America meridionali*.

TETRAGONIA. LINN. 551.

Calyx quadrifolius, coloratus. Corolla nulla. Styli quatuor. Capsula quadrangulatis sub flore.

n. *Tetragonia* foliis linearibus. *fruticosa*. LINN. *Sp.* 1. *Tetragonocarpus Afr. rutilans*, foliis longis & angustis. COMM. H. II. p. 205. T. 103. Fr.

CRASSULA. LINN. 352, DILL. Elth. 96.

Calyx quinquefolius. Petala quinque vnguibus connata. Stamina quinque, totidemque capsulae, ad basin nectariferae.

H. *Crassula* foliis planis cartilagineo-ciliatis basi connato-vaginantibus, *coccinea*. LINN. *Sp.* 1. *Cotyledon Afr. frutescens*, flore umbellato *coccinea*. COMM. *rar.* 24. T. 24. Fr.

H. *Crassula* foliis lanceolato-subulatis sessilibus connatis; canaliculatis subtus contextis, *perfoliata*. LINN. *Sp.* 2. *Crassula altissima perfoliata*. DILL. *Elth.* T. 96. f. 113. Albe *Afr. taulescens perfoliata glauca & non spinosa*. COMMET. *Pract.* 74. T. 23. Fr.

H. *Crassula* foliis subulatis obsolete tetragonis, *tetragona*. LINN. *Sp.* 3. ROY. *Lugdb.* 455. Fr.

H. *Crassula* foliis oppositis obtuse ovatis integerrimis: hinc angustioribus, *cultrata*. LINN. *Sp.* 4. *Crassula Anacamptoridis folio*. DILL. *Elth.* T. 97. f. 114. Fr.

H. *Crassula* *Portulacae* facie, arborescens. DILL. *Elth.* T. 101. f. 120. *Portulaca arborescens* vulgo. Genus satis certe nondum definiri potest, cum flores a nullo Botanico observari fuerint.

Alia etiam sunt genera, quae hucusque non sunt observata.

ORDO. III.

SVCCULENTAE *dyplostemones*.

Dyplostemones dicuntur, quibus stamina duplicatione ad petala aut segmenta petali sunt; ob summam autem analogiam variae plantae *isostemones* huc vocari debent.

COTYLEDON. LINN. 512. DILL.

Elth. 95.

Calyx & Corolla semiquinquefida. Capsulae quinque, singulis ad basin squama nectarifera.

H. Cotyledon foliis lineari-lanceolatis, integerrimis, floribus pendulis. Cotyledon *Afr. frutescens, folio longo & angusto, flore flavescente.* COMMEL. rar. T. 23. B. Cotyledon foliis depressis, cruciatis, WALTH. H. p. 16. c. 16.

H. Cotyledon foliis semiglobosis. *baemispbaerica.* LINN. Sp. 2. DILL. *Elth.* 112. T. 95. f. 111. Fr.

H. Cotyledon foliis laciniatis, floribus quadrifidis. *laciniata.* LINN. Spec. 6. Cotyledon *Afrq, folio crasso lato laciniato, flosculo aureo.* BOERH. Lugdb. 1. T. 288. Fr.

SEDVM. HALL. *Helv.* 392. LINN. 513.

Sempervivum. LINN. 518. Rhodiola.

LINN. 997.

Calycis folia quinque ad duodecim, totidemque petala. Stamina semper bis tot. Capsulae tot quot petala. In *Sadis* LINN. & *Rhodiola* squamae nectariferae ad basin capsularum.

H. Sedum caule ramoso, rofulis foliorum ciliatis obliquo. HALL. *Gutt.* 133. Sempervivum *caule arborecente laevi ramoso.* LINN. Sp. 1. Sedum *maius arborescens.* B. *Rin.* 282. *legitimum.* CLVS. *Hist.* II. p. 58. In *America & Europa* australiori. Fr.

Sedum rofulis sessilibus liberis conniventibus, foliis ciliatis, in apice purpureis. HALL. *H.* 392. n. 1.

L. 3

Sem-

libon folia profunde, dissecta, pedunculis multigloris.
femina & *mar.* KRETZ. To 7. f. 1. & T. 8. HALL.
 Gott. 124. Flores feminae staminibus effusis, abo-
 tivis gaudent. In *Virginia*. P.

HERMANNIA. LINN. 742.

Calyx simplex. Stamina quinque. Sigma sim-
 plex. Fructus quinquelocularis.

H. *Hermannia* foliis lanceolatis, obtusis, serratis.
Hyssopifolia. LINN. Sp. 4. *Althaea Africana* fru-
 tescent, floribus aureis cochlearis, pendulis minori-
 bus. VOLKAM. Norib. T. 24. In *Africa*. Fr.

PENTAPETES. LINN. 757.

Calyx simplex. Stamina decem, familia, binatis
 alterne posita, cum filamentis quinque sterilibus, co-
 loratis.

H. *Pentapetes* foliis hastato-lanceolatis serratis.
phoenicea. LINN. Sp. 1. *Alcea Indica*, lucido hastato
 folio, flore blattariae phoeniceo. PLUK. Phyt. T.
 245 f. 3. Al

GOSSYPIUM. LINN. 755.

Calyx duplex. Capsula quadrilocularis. Semina
 lana involuta.

H. *Gossypium* foliis palmatis, lobis lanceolatis,
 caule fruticoso. *arborescens*. LINN. Sp. 3. *Gossy-
 pium arborescens*, caule laevi. BAYH. Pin. 430. In
India. Fr.

CLASSIS IX.

SVCCULENTAE.

Caulis & folia succo aqueo turgida & carnosa.

ORDO I.

CACTI.

Calyx concavus monophyllus. Corolla in plurima petala profundissime fissa, parieti calycis affixa. Stamina plurima calyci inserta. Fructus carnosus sub calyce.

CACTVS. LINN. 539. *Cereus*. *Opuntia*.

Melocactus. *Tuna*. *Peireskia*.

Planta aphyllus. Calyx tubulato-cavus. Stylus cylindraceus stigmate multifido. Fructus carnosus, unilocularis. Sola *Peireskia* foliis instructa.

H. Cactus subrotundus tectus tuberculis, ovatis, barbatis. *mammillaris*. LINN. Sp. 1. *Echino-Melocactus minor lactescens, tuberculis f. mammillis maioribus*. HERM. *Parad.* 136. T. 136. *Ficoides Americana sphaerica, tuberculata lactescens, flore albo*. COMM. H. 1. p. 105. T. 55.

H. Cactus quadrangularis longus erectus, angulis compressis. *tetragonus*. LINN. Sp. 4. H. Vps. 119.

H. Cactus erectus sexangularis longus. *hexagonus*. LINN. Sp. 5. *Cereus altissimus Surinamensis*. HERM. *Par.* 116.

H. Cactus erectus longus subnovemangularis; angulis obsoletis, spinis lana brevioribus. *lanuginosus*.

L

LINN.

LINN. Sp. 8. An *Cereus Curassavicus erectus, laminae flavescente*. HERM. Parad. T. 115.?

H. Cactus repens subquinqueangularis, grandiflorus. LINN. Sp. 11. Cactus scandens angulis quinque pluribusque obtusis. H. Cliff. 182, c. descr. Cereus Americ. maior articulatus, flore maximo, nocte se aperiente. VOLK. Hesp. 1. T. 234. TREWII Icon. plant. rar.

H. Cactus repens decemangularis. flagelliformis. LINN. Sp. 12. Cactus scandens angulis decem obsoletis. H. Vpf. 121. Cereus minor scandens polygonus spinosissimus, flore purpureo. TREW. rar. pl. Det.

H. Cactus triangularis scandens articulatus. LINN. Sp. 13. Cereus erectus, foliis triangularibus profunde canaliculatis. PLVK. Alm. 147. T. 29. f. 3.

H. Cactus compressus, articulatus, ramosissimus, articulis ovatis, spinis setaceis. LINN. Cliff. 183. Cactus Opuntia. LINN. Sp. 15. Opuntia vulgo bebbiorum. 1. B. H. 1. p. 154. In America calidiori, & Virginia.

H. Cactus compressus articulatus, ramosus, articulis ovato-oblongis, spinis setaceis. LINN. Cliff. 183. Cactus Ficus indica. LINN. Sp. 16.

H. Cactus compressus articulatus ramosus, articulis ovato-oblongis, spinis subulatis. LINN. Cliff. 183. Cactus Tuna. LINN. Sp. 17. Tuna maior spinis validis flavicantibus, flore gilvo. DILL. Elth. 396. T. 295. f. 238.

H. Cactus tereti-compressus articulatus ramosus. LINN. Cliff. 182. Cactus Curassavicus. LINN. Sp.

19. *Ficus indica* f. *Opuntia Curassavica minima*.
COMM. H. 1. p. 107. T. 56.

H. *Cactus* prolifer ensiformi-compressus serrato-
repandus. *Phyllanthus*. LINN. Sp. 20. *Cactus foliis*
ensiformibus obtuse serratis. H. Cliff. l. c. *Cereus scolo-*
pendrii folio brachiato. DILL. *Eltb.* 73. T. 64. f. 74.

H. *Cactus* caule tereti arboreo, spinoso, foliis
lanceolato-ovatis. *Peireskia*. LINN. Sp. 21. *Peire-*
skia aculeata, flore albo, fructu flavescente. DILL.
Eltb. 305. T. 227. f. 294.

MESEMBRYANTHEMVM. LINN. 552. DILL.

Eltb. p. 159. Ficoides TOURN.

Calyx semiquinquefidus carnosus. Fructus carno-
sus tot loculis, quot styli adfunt. Folia in plerisque
opposita.

H. *Mesembryanthemum* foliis alternis ovatis obtu-
sis undulatis. *crystallinum*. LINN. Sp. 2. *Mesembryan-*
themum crystallinum, plantaginis folio undulato (mi-
tis argenteis adperso.) DILL. *Eltb.* 231. T. 180. f. 211.

H. *Mesembryanthemum* acaule, foliis linguiformi-
bus altero margine crassioribus. *linguiforme*. LINN.
Sp. 34. *Mesembr. folio linguiformi latiore*. DILL.
Eltb. T. 184 f. 224. Corolla flava.

H. *Mesembryanthemum* foliis acinaciformibus
connatis, angulo carinali scabris, ramis angulatis.
Mesembr. acinaciforme, flore amplissimo purpureo.
DILL. *Eltb.* T. 211. f. 271.

β. *Mesembryanthemum falcatum maius, flore*
amplo luteo. DILL. T. 212. f. 271.

communi applicatae. Stylus in quinque stigmata fissus.

I. Pedunculis unisloris.

1. *Geranium pedunculis unisloris, foliis palmatis.* HALL. H. 365. n. 1. *Geranium pedunculis unisloris, foliis quinquepartitis trifidis orbiculatis.* (*sanguineum.*) LINN. Sp. 38. In M. Altenstollberg. Prope specum Baumannianam. P.

H. *Geranium pedunculis unisloris, foliis triangularibus, quinquelobis, serratis.* HALL. Comm. Gott. T. I. p. 225. *Geranium pedunculis subunisloris, foliis quinquepartitis, acutis, foliolis pinnatifidis.* (*Sibiricum.*) LINN. Sp. 37. A.

II. Pedunculis bisloris, staminibus decem fertilibus.

1. *Geranium pedunculis bisloris, petalis integris longitudine calycis, caule prostrato, foliis reniformibus incis. rotundifolium.* LINN. Sp. 36. *Geranium foliis reniformibus semiquinquelobis, lobis trifidis.* HALL. Gott. 137. Helv. 365. n. 2. *Ger. folio malvae rotundo.* B. Pin. 318.

2. *Geranium foliis reniformibus, semisectifidis, caule erecto, umbellato flore.* HALL. H. 366. n. 3. *Geran. columbinum maius, flore minore coeruleo.* VAILL. Par. 79. T. 15 f. 1. Ad vias. A.

3. *Geranium foliis mollibus, reniformibus, semiquinquesidis, lobis semitridis obtusis.* HALL. H. 366. n. 5. *Geranium pedunculis bisloris folisque ramis alternis, caule ramoso diffuso, calycibus muticis.*

ticiis. (molle.) LINN. *Sp.* 33. *Ger. columbinum vil-
losum, petalis bifidis purpureis.* VAILL. *Par.* 79.
T. 15. *f.* 3. Ad vias & in agris. A.

*Geranium foliis ad nervum quinquefidis, pedun-
culis brevioribus, caule erecto.* HALL. *l. c. n.* 6. *Ge-
ranium pedunculis bifloris, foliis quinquepartito-
multifidis, caule glabro erectiusculo. (dissectum.)*
LINN. *Cent.* I, *Plant. n.* 62. VAILL. *T.* 15. *f.* 2.
In agris. A.

*Geranium foliis ad nervum vsque quinquefidis,
pedunculo longissime, prostrato caule.* HALL. *H.*
367. *n.* 7. *columbinum.* LINN. *Sp.* 35. VAILL. *T.*
15. *f.* 4. Circa *Rafemühle, Herperhausen.* A.

*Geranium pedunculis bifloris, foliis peltatis mul-
tipartitis rugosis pinnato-laciniatis acutis.* HALL. *H.*
n. 8. *pratense.* LINN. *Sp.* 29. *Ger. batrachioides f. Gra-
tia Dei Germanorum.* B. *Pin.* 318.

β. flore albo & coeruleo striato. In pratis. P.

*Geranium foliis rugosis, semitrilobis, lobis den-
tatis, lateralibus bipartitis, petalis purpureis.*
HALL. *l. c. n.* 9. *syriaticum.* LINN. *Sp.* 28. *Ger. ba-
trachioides folio Aconiti.* B. *Pin.* 317. In aquosis. P.

*H. Geranium pedunculis bifloris, foliis quinque-
lobis, lobis medio dilatatis, petalis emarginatis reticu-
latis. varicolor.* LINN. *Cent.* I. *n.* 61. *Ger. Romanum
varicolor f. striatum.* RAII *Hist.* 1063. In Italia. P.

*H. Geranium pedunculis bifloris, foliis caulinis
trilobis integris serratis, supremis subseffilibus.*
LINN. *Sp.* 27. *nodosum.* B. *Pin.* 318. In Gallia.

H. *Geranium foliis quinquelobis serratis, floribus paniculatis atropurpureis.* HALL. *Helv.* 368. n. 10. *Ger. phacum.* LINN. *Sp.* 26. *Ger. fuscum petalis reflexis.* I. R. H. In *Helvetia*, P.

H. *Geranium pedunculis bifloris, calycibus inflatis, pistillo longissimo. macrorrhizum.* LINN. *Sp.* 25. *Ger. batrachoides odoratum.* I. R. H. B. *Pin.* 318. P.

H. *Geranium foliis septilobis, lobis obtusis, dentatis, capsulis hirsutissimis.* HALL. *Gott.* 139. *Ger. Bobemicum.* LINN. *Cent.* 11. n. 176. *Ger. batrachoides Bobemicum, capsulis nigris hirsutis.* DILL. *Elth.* 159, T. 133. f. 160. P.

I *Geranium foliis quinque-trivepartitis, lobis pinnatifidis, calycibus hirsutis decemangulatis. robertianum.* LINN. *Sp.* 31. HALL. *H.* 369. n. 12. BLACKW. T. 480. Ad muros & rupes. P.

III. Pedunculis multifloris.

I *Geranium procumbens, pedunculis multifloris, calycibus pentaphyllis, foliis pinnatis, pinnis semipinnatis dentatis. cicutarium.* LINN. *Sp.* 21. α. HALL. n. 14. *Ger. cicutae folio minus & supinum.* B. *Pin.* 319. In hortis & agris. A.

H. *Geranium procumbens pedunculis multifloris, calycibus pentaphyllis, foliis pinnatis, pinnis incisis dentatis.* HALL. *H.* n. 13. *Gott.* 140. LINN. *Sp.* 21. β. *Ger. moschatum.* B. I. c. Riv.

H. *Geranium procumbens, pedunculis bi- & trifloris, foliis ovatis hirsutis crenato-incisis obtusis.* ROYEN. p. 451. n. 12. PLUK. *Phyt.* 301. f. 4.

Haec

Haec esse videtur species, quam descripsi in *Comment. Soc. Sc. Gott. T. V, Planta supina. Petioli & pedunculi breves, crassiusculi. Corollae plurimae apetalae mihi sunt visae, in paucioribus vix vnum alterumve petalum album, calyce minus, occurrit. Capsulae hirsutae. Stamina quinque sterilia.*

H. *Geranium pedunculis multifloris, foliis cordatis sublobatis, incis. malacoides. LINN. Sp. 22. Ger. folio Altheae. B. Pin. 318, malacoides. LOR. ic. 662. A.*

β. *Geranium Cbium, vernalum, Caryophyllatae folio. T.*

H. *Geranium pedunculis multifloris, foliis lobatis, lobis acutis serratis. HALL. Gott. 140. gruinum. LINN. Sp. 23. Ger. latifolium, acu longissimae B. Prodr. 138. Ger. Creticum. RIV. In Creta.*

μ. *Geranium pedunculis multifloris, foliis pinnatis, pinnatifidis, obtusis. cicutum. LINN. Cent. I. n. 60. HALL. Helv. n. 15. Ger. apulum Coriandri-folium. COL. Ecphr. I. p. 136. T. 135. In Europa australi. Pinnae minimae decurrentes maioribus intermixtae. Folia supina. Acus longissima. A.*

IV. *Pedunculis multifloris, calycibus monophyllis.*

H. *Geranium foliis hirsutis, duplicato pinnatis, pinnulis incis, floribus maculosis, radice subrotunda. HALL. Gott. 141. triste. LINN. Sp. 1. CORNV.T. Canad. 109. T. 110. Ger. noctu olens foliis Myrrhidis. BREYN. Cent. I. p. 126. T. 58.*

H. *Geranium calycibus monophyllis, foliis duplicato pinnatis, pinnulis laxè multifidis.* HALL. *G.* 141. *myrrhifolium.* LINN. *Sp.* 5. *Ger. myrrhidis folio, flore magno striato.* BREYN. *Cent.* 1. p. 129. *T.* 59. COMMEL. *H.* II. p. 125. *T.* 63. P.

H. *Geranium calycibus monophyllis tetrapetalon, foliis bipinnatis.* HALL. *l. r.* *Ger. Afr. Coriandri folio, floribus incarnatis.* HERM. *Lugdb.* 279. *T.* 280. RIV. *T.* 106. Corolla fere papilionacea. P.

H. *Geranium calycibus monophyllis, foliis dentatis, trifidis, medio lobo semitrilobo, lateralibus minoribus incis. HALL. l. c. Ger. fulgidum.* LINN. *Sp.* 2. *Ger. Afr. Alceae folio, flore coccineo, fulgidissimo.* DILL. *H. Elth.* 156. *T.* 130. f. 137. TILL. *Pis.* 68. *T.* 26. Fr.

H. *Geranium calycibus monophyllis, caule nodoso, foliis glaucis sinuatis.* HALL. p. 142. *Geran. gibbosum.* LINN. *Sp.* 8. *Ger. Afr. noctu olens Aquilegiae foliis.* HERM. *Lugdb.* 284. *T.* 285. RIV. *T.* 99. Fr.

H. *Geranium calycibus monophyllis, foliis cucullatis, dentatis.* *cucullatum.* LINN. *Sp.* 7. DILL. *Elth.* 155. *T.* 129. f. 156. *Geran. Afr. ibisci folio rotundo, carlinae odore.* HERM. *Lugdb.* 274. *T.* 275. Folia inferiora reniformia, non angulata. P.

H. *Geranium procumbens, calycibus monophyllis, foliis cordatis, glabris, obtusis, serratis.* HALL. *l. c.* *Geran. grossularioides.* LINN. *Sp.* 16. *Ger. Afr. vnae crispae folio, floribus exiguis rubellis.* HERM. *Lugdb.* 287. *T.* 289. RIV. *T.* 107. P.

H. *Geranium calycibus monophyllis, foliis hirsutissimis, oblongis, semiquinquelobis, undatis.* HALL. l. c. *Geran. capitatum.* LINN. *Spec.* 13. *Ger. Afr. frutescens, malvae folio laciniato, odorato.* HERM. *Lugdb.* 277. T. 278. RIV. T. 98. Fr.

H. *Geranium calycibus monophyllis, foliis oblongis, semiseptifidis, ferratis, mollissimis.* HALL. l. c. *Ger. Afr. Betonicae folio laciniato, floribus incarnatis.* HERM. *Lugdb.* 278. RIV. T. 100.

H. *Geranium calycibus monophyllis, foliis cordiformibus semitrilobis, ferratis, hirsutis.* HALL. p. 143. *Ger. vitifolium.* LINN. *Sp.* 12. *Ger. Afr. arbor. vitis folio, odore melissae.* DILL. *Elth.* 153. T. 126. f. 153. Fr.

H. *Geranium calycibus monophyllis, foliis cordiformibus, obtusis, ferratis, mollissimis.* HALL. l. c. *Ger. odoratissimum.* LINN. *Sp.* 15. *Ger. Afr. humile, folio fragantissimo molli.* DILL. *Elth.* 157. T. 131. f. 138.

H. *Geranium calycibus monophyllis, foliis pinguibus tomentosis, reniformibus obtuse crenatis.* HALL. l. c. *Geran. inquinans.* LINN. *Spec.* 3. *Ger. Afr. arbor. Malvae folio pingui, flore coccineo.* DILL. *Elth.* T. 125. f. 151. MART. *Cent.* 3. f. 3. RIV. T. 97. Fr.

H. *Geranium calycibus monophyllis, foliis cordato-orbicularibus incis, zona notatis, zonate.* LINN. *Sp.* 11. a. *Ger. foliis reniformibus obtuse dentatis, nigram circulum inscriptis.* HALL. 144. *Ger. Afr.*

arborescens Alchimillae folio, flore rubro. COMM. Prael. 51. T. 1. Fr.

β. *Geranium simile prioris, foliis album circum inscriptis.* HALL. 143. Geran. Afr. *Alchimillae folio maculato.* RIV. T. 102. Fr.

h. *Geranium calycibus monophyllis, foliis pilosis reniformibus, quinquelobis, lobis tridentatis.* HALL. G. 144. Ger. *alchimilloides.* LINN. Sp. 14. Ger. Afr. *alchimillae hirsuto folio, flosculis albidis.* HERM. Lugdb. 282. T. 283. Fr.

h. *Geranium calycibus monophyllis, foliis glabris subovatis carnosis crenatis. acetosum.* LINN. Sp. 10. Ger. Afr. *frutescens, folio crasso & glauco, acetosae sapore.* COMM. Prael. 54. T. 4.

h. *Geranium calycibus monophyllis, foliis pentagoniis integerrimis glabris peltatis. peltatum.* LINN. Sp. 10. Geran. Afr. *foliis inferioribus asari, superioribus saphidis agrinae maculatis splendidibus, & acetosae sapore.* COMM. Prael. 52. T. 2. Fr.

LINVM. LINN. 349.

Calycis folia & petala, & stamina quinque, totidemque styli. Fructus fere globosus, decemlocularis.

♂ *Linum calycibus capsulisque mucronatis, petalis crenatis, foliis lanceolatis alternis, caule subsolitario. vssitatissimum.* LINN. Sp. 1. *Linum ramis foliisque alternis lineari-lanceolatis, radice annua.* HALL. H. 370. n. 1. L. *sativum.* B. Pin. 214. Vbi que colitur. A.

β. Li-

β. *Linum sativum, latifolium, Africanum, fructu maiore.* I. R. H. 339.

η. *Linum calycibus capsulisque obtusis, foliis alternis lanceolatis integerrimis. perenne.* LINN. *Sp.* 2. HALL. *H. n. 2.* *L. perenne minus, coerulcum, capitulo maiore.* MORIS. *H. H. p.* 573. I. R. H. In *Sibiria, Anglia, Helvetia.* P.

Linum calycibus acuminatis, foliis sparsis lineari-setaceis retrorsum scabris. tenuifolium. LINN. *Sp.* 5. *Lin. perenne, foliis angustissimis, calyce aristato.* HALL. *H. 371. n. 3.* *Linum sylv. angustifolium flore magno.* I. R. H. 340. In siccis apricis, v. g. *über der Papiermühle, am Wege nach Robringen.* &c. P.

η. *Linum foliis oppositis lineari-lanceolatis, floribus terminalibus pedunculatis. Africanum.* LINN. *Sp.* 15. ROYEN. 434. *Linum Afric. luteum, foliis coniugatis.* BOERH. *p.* 284. In *Africa.*

Linum foliis coniugatis, petalis albis acutis. HALL. *H. 372. n. 4.* *Linum foliis oppositis ovato-lanceolatis, caule dichotomo, corollis acutis.* LINN. *Sp.* 17. *Linum catharticum.* BLACKW. *T.* 268. In pratis & gramineis. A.

Linum ramosissimum, flore tetrapetalo. HALL. *G.* 147. *Linum foliis oppositis, caule dichotomo, floribus tetrandris tetragynis. (Radiola.)* LINN. *Sp.* 18. *Chamaelinum vulg.* VAILL. *Parif.* 33. *T. 4. f. 6.* In sabulosis, v. g. *Cellenfibus & circa Berolinum.*

DROSERA. LINN. 351. Rorella

HALL.

Calyx semiquinquefidus, petala subovata. Stamina & styli quinque. Folia roriferis villis ciliata.

1. *Drosera scapis radicatis, foliis orbicularis, rotundifolia.* LINN. Sp. 1. *Rorella caule simplici, foliis rotundis retusis.* HALL. H. 372. n. 2. *Ros Solis, folio rotundo.* B. Pin. 357.

2. *Drosera scapis radicatis, foliis oblongis, longifolia.* LINN. Sp. 2. *Rorella foliis ellipticis.* HALL. 46. n. 1. *Ros solis L. Salsoroga.* THAL. Herc. 116. T. 9. f. 2. Vtraque in paludosis circa M. Bructerum.

FAGONIA. LINN. 475.

H. *Fagonia spinosa, foliis lanceolatis planis linearibus.* LINN. Sp. 1. *Trifolium spinosum creticum.* CLVS. CCXLII. C. B. Prodr. 142. In Oriente. P.

TRIBULVS. LINN. 476.

Petala quinque. Stylus simplex. Fructus ex quinque capsulis aculeatis compositus.

H. *Tribulus foliolis sexiugatis subaequalibus, terreftris.* LINN. Sp. 3. *Tribulus terrestris.* LÖB. ic. n. p. 841. In Europa australi. A.

MELIA. LINN. 473. Azedarach. T. 387.

Calyx minimus. Nectario cylindrico decem stamina se inferunt. Stylus simplex.

H. *Melia* foliis decempartitis. *BAHL. G.* 161. *Melia foliis bipinnatis.* LINN. *Sp.* 1. *Azedarach.* DOD. *Pempt.* 848. 1. R. H. 616. In *Syria.* Fr.

ZYGOPHYLLVM. LINN. 474. *Fabago.*

T. 135.

Petala ovata. Circa stamina decem totidem squamae erectae, conniventes. Stylus simplex.

H. *Zygophyllum* foliis petiolatis, capsulis prismatico-pentaëdricis. LINN. *Sp.* 1. *Fabago Belgarum.* 1. R. H. *Capparis Fabago.* DOD. *Pempt.* 471. In *Syria.* P.

DYPLOSTEMONES ODORAE.

Omnes, uti priores, dyplostemonas, flores in plerisque numero partium quinario- & quaternario ludunt. Loculi fructus tot, quot petala. Omnes graveolentes.

RVTA. LINN. 469.

Petala concava. Capsula punctata quadri- aut quinqueloba, cruce insculpta.

H. *Ruta* foliis decompositis. *graveolens.* LINN. *Sp.* 1. *Ruta bortenfis latifolia.* 1. R. H. B. *Pin.* 336. BLACKW. T. 7. Petala oris laciniatis. P.

PEGANVM. LINN. 530. *Harmala.* TOWN.

Rutae affinis. Sed stamina quindecim, capsula trigona, trilocularia. Stigmata tria.

H. *Peganum* foliis multifidis. LINN. *Sp.* 1. *Harmala* DOD. *Pempt.* 121. *Purg.* 420. LOB. p. 55. In *Europa australiori & Syria.* P.

MONOTROPA. LINN. 477. Hypopitys.

HALL. 411.

Habitus aphyllus. Flores terminales quinquepartiti, laterales quadripartiti.

1 Monotropa floribus lateralibus octandris, terminali decandro: LINN. *Sp.* 1. Hypopitys *spica mutante.* HALL. *H.* p. 411. Orobanche *Hypopitys.* C. B. *Prodr.* 31. In umbrosis sylvis parasitica radicum.

ADOXA. LINN. 450. Moschatellina HALL. 412.

TOURN. T. 68.

Flores in spicas cubicas collecti.

1 Adoxa. LINN. *Sp.* p. 367. Moschatellina. HALL. *H.* 412. In dumetis versus *Eichenkrug.*

PARIS. LINN. 449. HALL. 412.

Omnes partes floris numerum quaternarium observant. Fr. bacca.

1 Paris foliis quaternis. LINN. *Spec.* p. 367. HALL. *H.* 412. Herba Paris. L. E. H. BLACKW. T. 286. Solanum *quadrifolium bacciferum.* B. Pin. 167. In umbrosis sylvis & dumetis.

DYPLOSTEMONES ALSINACEAE.

Calyx quinquefolius. Petala quinque, patentia. Stamina bis tot, paucioribus exceptis. Tubae tres, quatuor & quinque. Capsula unilocularis. Folia opposita.

ALSINE.

ALSINE. HALL. H. 385.

Capſula conica. Calycis folia obtuſiora.

I. Petalis integris ; tubis trinis.

Alſine foliis lanceolatis, trinerviis, petalis integris. HALL. H. n. 18. *Arenaria foliis ſubovatis acutis petiolatis nervoſis. (trinervia.)* LINN. Sp. 4. Alſine *plantaginis folio*. 1. B. Hiſt. III. 364. In ſylvis montanis. A.

Alſine foliis lanceolatis, petalis integris calycis magnitudine. HALL. H. n. 17. *Arenaria ſerpyllifolia*. LINN. Sp. 16. Alſine *minor multicaulis*. B. Pin. 251. In vallis & ſylvis macilentis. A.

Alſine foliis peranguiſtis, ad baſin lateſcentibus. HALL. H. n. 15. *Arenaria foliis ſubulatis, caule paniculato, petalis calyce brevioribus*. LINN. Sp. 8. Alſine *tenniſolia*. 1. B. VAILL. Par. T. 3. f. 1. In M. Heymberg. A.

Alſine foliis faſciculatis, petiolis ſimplicibus, petalis integris. HALL. H. n. 9. *Spergula foliis oppoſitis ſubulatis laevibus, caulibus ſimplicibus. (nodosa.)* LINN. Sp. 3. Alſine *nodosa Germanica*. B. Pin. 251. In vdis circa Seeburg. A.

Alſine foliis coniugatis, anguiſtis, linearibus, petalis integris. HALL. H. n. 14. *Arenaria*. LINN. Sp. 7. Alſine *saxatilis & multiflora, capillaceo folio*. VAILL. T. 2. f. 3. Circa Herzberg & in ſylva Hercynica.

Alſine tetrapetala, foliis anguiſtis, in origine lateſcentibus. HALL. H. n. 20. *Sagina ramis procum-*

bentibus. LINN. Sp. 1. *Alfina minima flore fugaci*.
J. R. H.

Alfina foliis filiformibus, vaginis ad foliorum
ortum squalentibus. HALL. H. n. 8. *Arenaria ru-*
bra. LINN. Sp. 6. *Spergula purpurea*. J. B. III. p. 722.
In agris siccioribus.

1. β. Eadem pulposior & altior, locis salis. v. g. in
prato prope *Herle*.

II. Petalis integris, rubis quinque.

Alfina foliis verticillatis, petiolis ramosis, petalis
integris. HALL. H. n. 7. *Spergula arvensis*. LINN.
Sp. 1. *Alfina Spergula dicta maior*. J. B. Pin. 251.
In arvis. A.

III. Pentastemon.

Alfina pentastemon, petalis serratis, umbellifera.
HALL. H. n. 6. *Holosteum floribus umbellatis*.
LINN. Sp. 3. DILL. nov. gen. p. 130. T. 6. TABERN.
ic. p. 233. In vallo urbis, hortis & agris. A.

Alfina pentastemon, petalis bipartitis, foliis ova-
to-cordatis. HALL. H. n. 5. LINN. Sp. 1. *Alfina*
media. J. B. Pin. 250. A.

IV. Petalis bipartitis, decastemon.

Alfina foliis cordatis, petalis bipartitis. HALL. H.
n. 4. *Stellaria foliis cordatis petiolatis, pedunculis*
ramosis, (*nemerum*.) LINN. Sp. 1. *Alfina atrissima*
nemerum. J. B. Pin. 251. *Alfina maior*. TABERN. p.
707. In fossis. A.

Alfina foliis ovato-lanceolatis, petalis bipartitis.
HALL. H. n. 1. *Alfina longifolia, uliginosis locis*
pro-

proveniens. I. B. III. p. 365. *Alfine aquatica media*.
I. R. H. Super *Hardeggen*; in palude inter *Harste*
& *Barenfen*.

Alfine foliis gramineis, petalis bipartitis. HALL.
H. 2: *Stellaria foliis linearibus, integerrimis*. LINN.
Sp. 5: *Gramen floridum minus*. TABERN. ic. 232.
In agris & palustribus inter Juncos. A.

Alfine foliis gramineis, asperis, petalis semibifidis. HALL. *H. n. 1*: *Stellaria Holostea*. LINN. *Sp. 4*:
Gramen Fuchsis. I. B. III. p. 361. In sylvis & dumetis.

CERASTIUM, LINN. 518, *Myosotis*.
HALL. 383.

Petala bifida. Capsula ut plurimum longior, incurva.

Cerastium (Myosotis) foliis oblongis, obtusis, petalis calyce maioribus. HALL. *H. n. 3*: *arvensis*. LINN. *Sp. 5*: *Myosotis arvensis hirsuta, flore maiore*. VAILL. *Par. 141. T. 30. f. 4*. I. R. H. *Ad vias*. A.

Cerastium (Myosotis) foliis ovatis, acuminatis, flore calyci aequali. HALL. *H. n. 4*: *Myosotis hirsuta, parvo, albo flore*. I. R. H. VAILL. *Par. T. 30. f. 1*. A.

Cerastium (Myosotis) floribus pentandris, petalis emarginatis. LINN. *Sp. 3*: *Myosotis exigua pentastemon*. HALL. *H. 5*: *Myosotis arv. hirsuta minor*. VAILL. *Par. 142. T. 30. f. 2*. Filamenta quinque sterilia, quinque antherifera. Styli tres. A.

Cerastium (Myosotis) foliis lanceolatis, caule dichotomo ramosissimo, capsulis erectis. LINN. *Sp.*

6. *Myosotis Hispanica segetum.* I. R. H. *Alfene corniculata.* CLVS. *Hist.* II. p. 184. A.

7. *Cerastium (Myosotis) viscosa, foliis ovatis.* HALL. *H. 26.* *Cerastium erectum villoso-viscosum.*

LINN. *Spec.* 2. *Myosotis hirsuta altera viscosa.*

PAILL. *Par.* 142. T. 30. f. 1. In macilentis. A.

8. *Cerastium (Myosotis) foliis connatis.* LINN.

Sp. 1. *Myosotis orient. perfoliata, lychnidis folio.*

DILL. *Eltb.* 295. T. 217. f. 284. A.

DYPLOSTEMONES CARYOPHYLLAEAE.

Calyx monophyllus, tubulosus, ore quinquesido.

Petala quinque, vnguibus linearibus, bractea re-

flexa, latiori. Stamina decem, alterne maiora. Styli

duo, tres, quinque. Capsula oblonga, unilocularis.

Folia opposita, simplicia. Secundum Hallerum ge-

nera minora ex tubis constitui possunt.

VISCAGO. HALL. DILL. *Cucubalus.*

LINN. *Silene.* LINN.

Styli tres. Summitates saepe viscidae.

I. Petala simplicia sine corolla auricularum.

Cucubali. LINN.

9. *H. Viscago (Cucubalus) calycibus campanulatis, petalis distantibus, fructu colorato, ramis divaricatis.*

LINN. *Sp.* 1. *Alfene scandens baccifera.* B. *Pin.* 250.

Cucubalus Plinii vulgo. Circa *Jenam.* P.

Visca-

Viscago (*Cucubalus*) calycibus subglobosis glabris reticulato-venosis, capsulis trilocularibus, corollis serratis, bifidis. LINN. *Sp.* 2. HALL. *H.* 2. *Lychnis sylv. quae Beben album vulgo*, B. *Pin.* 205. In collibus saxosis, ficcis. P.

H. *Viscago* calycibus ovatis glabris venoso-reticulatis, floribus subsessilibus, petalis calyce parum maioribus, denticulatis, radice annua. *Silene* LINN. *Sp.* 13. *Viscago vesicaria Cretica, flore parvo purpurascente*. DILL. *Elsh. T.* 317. f. 409. A.

H. *Viscago* (*Cucubalus*) petalis bipartitis, floribus secundis decumbentibus, pedunculis solitariis oppositis erectis, caule simplicissimo. *Tataricus*. LINN. *Sp.* 8. *Lychnis septemtrionalium, foliis byssopi, floribus vno versu dispositis*. HALL. *G.* 150. P.

H. *Viscago* foliis imis petiolatis, ovatis, floribus paniculatis, petalis emarginatis. HALL. *Gott. ed. prior. & ic. in app. & edit. alt. p.* 150.

H. *Viscago* foliis imis petiolatis, floribus verticillatim spicatis, petalis indivisis, sexu distincta. HALL. *H. n.* 5. *Cucubalus Otites*. LINN. *Sp.* 10. *Lychnis viscosa, flore muscosa*. B. *Pin.* 206. In *Marchia*. Prope arcem *Regenstein*.

H. *Viscago* foliis imis petiolatis, floribus verticillatim spicatis, petalis indivisis, androgyna. HALL. *G. p.* 165.

Viscago (*Cucubalus*) floribus spicatis alternis subsessilibus, corollis obsoletis nudis. LINN. *Sp.* 12. *Lychnis sylv. alba, floribus reflexis*. MAGN. *Monsp.* II. *T.* 170.

II. Petala denticulata, *Silene* LINN.

H. Viscago calyce duplo petalis longiore, floribus umbellatis. HALL. *H.* n. 3. *Silene floribus fasciculatis fastigiatis, foliis superioribus cordatis glabris.* (*Armeria.*) LINN. *Sp.* 24. In *Valefia. Gulja. A.*

I. Viscago foliis imis petiolatis, ovatis, petalis bifidis se convolventibus. HALL. *Helv.* n. 4. *Silene foliis lanceolatis, caule paniculato, floribus nutantibus, calyce striato, corolla involuta.* LINN. *Sp.* 7. In saxosis circa *Gottingam.*

H. *Silene* petalis bifidis, corolla subcoadunata, floribus secundis, calycibus erectis raris pilosis, pedunculis oppositis trifloris, ramis alternis. *amoena.* LINN. *Sp.* 8. *H. Vps.* 114. In *Sibiria. A.*

H. *Silene* petalis bifidis, caule fruticoso, foliis lato-lanceolatis glabris, panicula trichotoma. *fruticosa.* LINN. *Sp.* 9. *H. Vps.* 111. n. 2. *Lychnis sylv. myrtifolia Beben albo similis.* B. *Pin.* 205. *Ocymoides fruticosum.* CAMER. *Hort.* 109. *T.* 33. Fr.

H. Viscago (*Silene*) foliis radicalibus cochleariformibus obtusissimis, caule subverticillato. *gigantea.* LINN. *Sp.* 10. *Lychnis Graeca, Sedi arborescentis folio & facie, flore albo.* TOURN. *Cor.* 23. WALTH. *H.* p. 32. *T.* XL. P.

H. Viscago caule foliisque hirsutis ovato lanceolatis, floribus raris, petalis semibifidis. *Lychnis pcymastri facie, flore viridi.* HERM. *Parad.* *T.* 199.

H. *Viscago* foliis inferioribus obverse ovatis, superioribus lineari-lanceolatis, calycibus erectis subsiccatis, petalis emarginatis. *Silene crotica*. LINN. *Sp.* 20. *H. Vps.* 112. *Viscago foliis inferioribus bellidis, superioribus tunicae, calyce strictiore & turgidior.* DILL. *Elth.* 422. T. 314. f. 404. 405. In *Creta*.

H. *Viscago* (*Silene*) foliis obverse ovatis, floribus spicatis alternis secundis sessilibus, petalis bifidis inaequalibus. *nocturna*. LINN. *Sp.* 4. *Viscago birta noctiflora, floribus obsolete spicatis.* DILL. *Elth.* T. 310. f. 400. In *Pensylvania*. A.

H. *Viscago* (*Silene*) hirsuta, petalis dentatis, floribus erectis, fructibus reflexis subsessilibus alternis. *Lusitana*. LINN. *Sp.* 2. *Viscago hirsuta Lusitanica, bellato flore.* DILL. *Elth.* T. 311. f. 401. In *Lusitania*. A.

H. *Viscago* (*Silene*) petalis integerrimis subrotundis, fructibus erectis alternis. *quinguevulnera*. LINN. *Spec.* 3. *Lychnis hirsuta minor, flore variegato.* HALL. *Gott.* 152. In *Hispania*. A.

H. *Viscago* (*Silene*) calycibus fructiferis pendulis inflatis, angulis decem scabris. *pendula*. LINN. *Sp.* 14. *Viscago birta Sicula, lychnidis aquaticae facie, supina.* DILL. *Elth.* T. 312. f. 404. Folia inferiora spatulata, superiora lanceolata, caulis debilis. In *Creta, Sicilia*. A.

H. *Viscago* (*Silene*) calycibus fructus globosis acuminatis striis triginta foliis glabris. *conoides*. LINN.

LINN. *Sp.* 11. *Lychnis sylvestris latifolia, calycibus turgidis striatis.* B. *Pin.* 205. A.

Viscago (Silene) erecta laevis, calyceibus subglobosis glabris venosis, corollis inapertis. rubella. LINN. *Sp.* 18. *H. Vps.* 112. *Viscago Lust. flore rubello vix conspicuo.* BILL. *Elst.* T. 314-f. 406. A.

LYCHNIS. HALL. 376. *Lychnis* LINN. 517.
Agrostema LINN. 516.

Styli quinque. *Agrostema*. Segmenta calycis coriacei flore longiora; *Lychnis* LINN. Calyx subrotundus inflatus, filamenta staminum alis aucta.

H. *Lychnis* (*Agrostema*) tomentosa, foliis ovato-lanceolatis, petalis coronatis emarginatis. LINN. *Sp.* 3. HALL. *H. n.* 1. *Lychnis coronaria Dioscoridis sativa.* B. *Pin.* 203. In *Helvetia. Italia.* A.

1 *Lychnis (Agrostema) hirsuta, calycibus corollam aequantibus, petalis integris nudis.* LINN. *Sp.* 1. HALL. *H.* 2. *Lychnis segetum maior.* B. *Pin.* 204. *Nigellastrum.* DODON. *Coron.* p. 49. Inter segetes. A.

1 *Lychnis* sexu distincta, foliis ovatis, acutis. HALL. *H.* 3. LINN. *Sp.* 6. *Lychnis sylvestris purpurea simplex.* β. *eadem alba.* B. *Pin.* 204. γ. Variet. *fl. pleno.* *Ocymastrum rubrum & album.* TABERN. 299. *Alba circa Rosdorf ad viam.*

H. *Lychnis* floribus umbellatis, petalis breviter quadrifidis. HALL. *Gott.* 1531. *Lychnis floribus fasciculatis fastigiatis. (Chalcedonica.)* LINN. *Sp.* 1. *Flos Constantinopolitanus.* DOD. *Pempt.* 178. *flore coccinea aut incarnato.* In *Oriente.* A.

Lychnis

Lychnis petalis glandulosa, longilaminis, fructu subrotundo. LINN. Sp. 2. HALL. H. 378. *Caryophyllus pratensis*, flore laciniato simplici s. floribus cæci aut flore pleno. B. Pin. 219. In pratis. B. H.

Lychnis petalis integra, fructu quinqueloculari. *Viscaria*. LINN. Sp. 3. *Lychnis fida viscosa rubra angustifolia*. B. Pin. 205. Circa Ellingerode & Blankenburg. In agro ubi viscaria præcipue colitur.

SAPONARIA. LINN. 499.

Calyx oblongus. Ungues petalorum angusti, capsula cylindrica. Styli duo.

Saponaria calycibus cylindricis, foliis ovato lanceolatis, officinalis. LINN. Sp. 1. *Saponaria foliis trinerviis, floribus ut bellatis*. HALL. H. 2. *Saponaria vulgaris*. In vallo urbis. P.

Saponaria caule brachiato, foliis ovatis acuminatis sessilibus, calyce pentagono turgido. HALL. H. 3. *Vaccaria*. LINN. Sp. 2. *Lychnis segetum rubra foliis persiliatis*. B. Pin. 204. Inter segetes. A.

Saponaria caule dichotomo, foliis lanceolatis, petiolatis, floribus solitariis, calycibus cylindricis villosis, orientalis. LINN. Sp. 4. H. Vpf. 106. n. 2. *Lychnis orientalis annua supina, antirrhini folio, flore minimo purpurascente*. DILL. Elth. T. 167. f. 204. A.

GYPSOPHILA. LINN. 498.

Calyx campanulatus. Petala ovata, unguibus minimis. Filamenta staminum subulata.

1. *Gypsophila* foliis lanceolatis scabris, corollis
revolutis, (ramis in flore lentibus.) paniculata.
LINN. Sp. 3. In Sibiria.

2. *Gypsophila* foliis lanceolatis subtrinerviis re-
gis. altissima. LINN. Sp. 4. Gen. non. 1403. H. Kpf.
147. In Sibiria. P.

3. *Gypsophila* foliis lanceolato-linearibus ciliatis
obsolete triquetris, laevibus obtusis secundis.
glata. LINN. Sp. 5. *Saponaria petalis ovatis; fo-*
liis glaucis pulposi linearibus. HALL. Fl. Jen. Rupp.
187. T. 2. f. 11. *Gymphytum petraeum.* THAL. Herc.
p. 117. In M. Altenstollberg. P.

4. *Gypsophila* foliis lanceolatis semimplexicau-
libus perfoliatis. LINN. Sp. 6. *Spergula multiflora;*
foliis inferioribus saponariae, superioribus Beberga-
milibus. DILL. Eth. T. 276. f. 357. In Hispania. P.

5. *Gypsophila* foliis lineari-lanceolatis planis, caule
dichotomo, pedunculis bifloris, petalis emarginatis.
rigida. LINN. Sp. 6. *Saponaria.* HALL. H. 8. *Lych-*
nis annua minima, flore carneo lineis purpureis dis-
similis. 1. R. H. Inter segetes exeunte aestate. A.

6. *Gypsophila* caule dichotomo, foliis subulatis
calycibus quinqueangularibus, squamis ad basin ac-
cessoriis. *Caryophyllus muralis minimus.* B. Pin.
211. Videtur esse *Gypsophila.* LINN. Sp. 7. Etenim
squamae ad basin calycis adsint, summa tamen
cum priori specie analogia, et vita continua impediunt,
quo minus ad *Dianthus saxifragum* LINN. refer-
ri possit. A.

НАБЛЮДЕНИЯ

H. *Dianthus floribus aggregatis fasciculatis, Equan-*
mis calycinis ovato-fimbriatis tubum aequantibus,
foliis lanceolatis. barbatus Linn. Sp. i. Caryophyl-
lus barbatus bortenfis latifolius & angustifolius.
 B. Pin. 206.

Dianthus floribus aggregatis fascicularis, squamis calycinis lanceolatis villosis, tubum aequantibus.
Armeria. LINN. Sp. 3. Tunica. HALL. 3. Caryophyllus *barbatus sylvestris*, p. Pen. 208. *Armeria sylvestris altera*. LOB. ic. p. 448. A.

h. *Dianthus floribus foliis, squamis, calycinis*
subovatis brevissimis, corollis crenatis. Caryophyl-
lus. LINN. Sp. 5. Caryophyllus hortensis. f. plen.
3, Pin. 208. BLACKW. T. 85. Varietates flore
pleno.

H. *Dianthus floribus solitariis squamis calycinis subulatis, tubum aequantibus, corollis crenatis.*

LINN. Sp. 8. *Caryophyllus Chinensis supinus, Linn. coli folio, flore unica.* TOURN. Art. Per. 1705. p. 348. f. 5. A.

H. *Dianthus floribus subsolitariis, squamis calycinis lanceolatis quaternis, corollis crenatis. glaucus.* LINN. Sp. 5. *Tunica ramosior, flore candido cum corolla purpurea.* DILL. Elth. 400. T. 298. f. 348. In Anglia. P.

Dianthus floribus solitariis, squamis calycinis subovatis brevissimis, corollis multifidis fauce pubescentibus. plumarius. LINN. Sp. 9. *Tunica.* HALL. n. 5. *Caryophyllus flore tenuissime dissecto.* B. Pin. 209. In palcis nemorosis.

H. *Dianthus floribus solitariis, foliis subulatis, squamis calycinis mucicis, floribus profunde fimbriatis, lacteis.* G. Met. Flad. Gott. p. 156. P.

Dianthus floribus subsolitariis, squamis calycinis tubo parum minoribus, corollis serratis, punctatis. Caryophyllus simplex supinus latifolius. B. Ph. 108. HALL. G. 156. *Caryophyllus minor pubellus, supinus, maculis aureis argenteisve dispersis.* BOB. ic. 444. *Prope Eichenkrug, Rheinhausen & alibi in arenis.*

H. *Dianthus caulibus unifloris, squamis calycinis ovatis, corollis multatis, foliis linearibus. arenarius.* LINN. Sp. 107. *Tunica foliis brevibus rigidis, caule uniflora.* HALL. H. n. 8. *Caryophyllus sylvestris, flore unico.* B. Pin. 209. BOB. ic. 465. *sylvestris.* L. CLVS. I. p. 282. DILL. Elth. 402. In Marchia.

DIPLOSTEMONES. CALYCFLOREAE.

Calyx monophyllus. Petala patentia & stamina bis tot calyci inserta. Stigmata tot, quot loculi fructus sub calyce positi.

LYTHRUM. LINN. 532. *Salicaria*.

TOURN. 129.

Calyx tubulosus, segmentis sex maioribus, sex minoribus alternis, introsum flexis. Petala sex.

Lythrum (Salicaria) foliis oppositis cordato lanceolatis, floribus spicatis duodecandris. LINN. *Sp.* 1. HALL. *H. n.* 1. *Salicaria vulgaris purpurea foliis oblongis.* B. R. H. *Lythrum spicata purpurea.* B. *Pin.* 246. Ad ripas aquarum. P.

Lythrum (Salicaria) foliis alternis linearibus, floribus hexandris. *Hyssopifolia.* LINN. *Sp.* 5. HALL. *Fl. Jen. Rupp. T.* 2. f. 3. B. *Pin.* 246. Auf Aeckern hinter der Graetzels Mühle.

PEPLIS. LINN. 402.

Calyx campanulatus, caeterum priori similis. Petala fugacia, exigua. Stamina sex.

Peplis foliis coniugatis. *Portula.* LINN. *Spec.* 1. *Glaux palustris, flore striato clauso, foliis portulacae.* PETIT. *Gen.* 43. *T.* 43. MICH. *gen.* 21. *T.* 43. VAILL. *Par.* 80. *T.* 15. f. 5. In paludibus M. Hagen.

HEUCHERA. LINN. 283.

Calyx semiquinquefidus. Stamina quinque. Capsula ovata birostris.

1 H. Heuchera. *Americana*. LINN. *Sp. p.* 226. *Cortusa Americana*, flore squalide purpureo. HERM. *Parad.* 131. *T.* 131. In *Virginia*. P.

MITELLA. LINN. 496.

Calyx campanulatus. Petala quinque multifido-capillaria. Stamina decem.

1 H. Mitella scapo diphylo. LINN. *Sp. i.* *Cortusa Americana*, spicata flore, petalis fimbriatis. HERM. *Parad.* 190. In *America septentrionali*. P.

AGRIMONIA. LINN. 534.

Calyx bifolius sub fructu, cui insidet calyx exiguus quinquefidus. Stamina circiter duodecim.

1 Agrimonia foliis caulinis pinnatis, fructibus hispidis. *Eupatoria*. LINN. *Sp. i.* HALL. *H. p.* 467. *Agrimonia*. B. *Pin.* 321. BLACKW. *T.* 21. Ad vias secas. P.

β. Agrimonia odorata. B. P. l. c.

OENOTHERA. LINN. 424. Onagra,

TOURN.

Capsulae cylindricae insidet calyx tubulosus, limbo quadri-partito. Petala cordata. Stylus stigmatibus quadrifido.

H. Oeno-

H. *Oenothera* foliis ovato-lanceolatis planis.
LINN. Sp. 1. *Onagra latifolia*. 1. R. H. *Lysima-*
chia lutea corniculata. B. Pin. 245. In *Marchia*. B.

H. *Oenothera* foliis lanceolatis, capsulis ventri-
toso-conicis. An *Onagra angustifolia*, caule rubro,
flore minori? TOURN. Inst. 302. Flores & fructus
densius congesti, caulibus & calyx rubro tinctus. Pru-
ctus priori fere aequalis, sibs autem triplo
minor.

H. *Oenothera* foliis glabris, flore suave purpu-
reo. Comm. Soc. R. Sc. Gott. 1751, p. 224. In *Vir-*
ginia.

H. *Oenothera* foliis lanceolatis undulatis. *mollis-*
sima. LINN. Sp. 2. *Onagra bonariensis villosa*, flo-
re mutabili. DILL. Eltb. 297. T. 219. f. 286.

EPILOBIVM. LINN. 246.

Capsulae cylindricae papposis seminibus foetae
insistit flos similis priori, sed calyx est quadrisolius,
subo carens.

Epilobium spicatum, flore difformi, foliorum
nervo eminente, ora leviter serrata. HALL. H. n. 1.
Epilobium flore difformi, staminibus & pistillis
declinatis. LINN. Fl. Suec. ed. pr. 304. *Chamae-*
nerium vulgare latifolium. 1. R. H. B. Pin. 245.
In sylvis.

Epilobium foliis ovato-lanceolatis, serratis, le-
vissime hirsutis. HALL. H. n. 5. *Epilobium foliis*
oppositis ovatis dentatis. (*montanum*). LINN. Sp. 4.
Chamaenerium glabrum maius. 1. R. H. In umbrosis.

1. *Epilobium* foliis ellipticis obtuse lanceolatis, totum leve. HALL. *Gott.* p. 162. *Epilobium foliis vaginatis lanceolatis, integris petalis, caule erecto. (palustre.)* LINN. *Sp.* 6. *Chamaenerium glabrum angustifolium.* DILL. *Cat. Giesl.* p. 208. In palude inter Barenfen & Gladebeck.

2. *Epilobium* foliis longis, dentatis, villosis, flore minore. HALL. *H. n.* 7. *Epilobium hirsutum.* LINN. *Sp.* 2. β . *Chamaenerium villosum, maius, parva flore.* I. R. H. In fossis.

3. *Epilobium* foliis longis dentatis, villosis, flore maiore. HALL. *H. n.* 8. *Epilobium hirsutum.* LINN. *Sp.* 2. α . *Chamaenerium villosum, flore magno purpureo.* I. R. H. Ad rivas.

DIPLOSTEMONES BICORNES.

Calyx exiguus. Corolla monopetala limbo quinquefido. Stamina aut receptaculo, aut basi corollae tantum affixa. Antherae binas fetas exferunt. Tuba simplex. Fructus quadri- aut quinquelocularis.

VACCINIUM. LINN. 434.

Corolla campanulata, limbo quadrifido. Germen infra florem. Fr. Bacca umbilicata.

Vaccinium foliis integerrimis revolutis, ovatis, caulibus repentibus filiformibus, corolla ad basin fere quadrifida. LINN. *Sp.* 11. *Oxycoccus* HALL. *H.* p. 413. *Vitis Idaea palustris.* B. Pin. 471. Corollae segmenta per aetatem ira revolvuntur, ut flos tetrapetalos videatur. In turfaceis circa M. Berg. *fieri.* Fr.

Vac-

Vaccinium foliis perennantibus obverse ovatis, integerrimis, revolutis, subtus punctatis. HALL. *H.* n. 1. *Vaccinium Vitis Idaea*. LINN. *Sp.* 10. *Vitis Idaea foliis subrotunder non crenatis, baccis rubris.* B. Pin. 470. 1. R. H. Vbiq̃ue in sylva *hercynica*. Fr.

Vaccinium foliis ovalibus integerrimis, basi ciliatis deciduis. *virginosum*. LINN. *Sp.* 3. *Vaccinium foliis annuis exalbidis.* HALL. *Helv.* 2. *Vitis Idaea* II. CLVS. *Hisp.* l. p. 61. T. 61. In sylva *Hercynica*. Fr.

Vaccinium caule angulato, foliis serratis annuis. LINN. *Cliff.* 148. HALL. *H.* n. 3. *Vaccinium Myrtillus* LINN. *Spic.* 1. BLACKW. *T.* 463. *Vitis Idaea foliis oblongis crenatis, fructu nigro.* B. Pin. 470. *Myrtillus* MATH. p. 231. In sylvis. Fr.

ARBUTVS. LINN. 488. TOURN. 368. *Vua vrsi.* T. 370.

Priori similis, sed flos quinquefidus, & fructus intra florem.

H. *Arbutus* caule erecto, foliis glabris serratis, baccis polyspermis, tuberculis seminum exasperatis. *Vnedo*. LINN. *Sp.* 1. *Arbutus folie serrato*. B. Pin. 460. *Arbutus*. CLVS. *Hisp.* p. 77. In *Europa* australi. Fr.

Arbutus caulibus procumbentibus, foliis integerrimis. *Vua vrsi*. LINN. *Sp.* 4. HALL. *H.* n. 2. *Vua vrsi* CLVS. *Hisp.* l. p. 63. TABERN. p. 1080. I. R. H. In *Gallensi* ducatu. Fr.

ANDROMEDA. LINN. 485.

Corolla ovata, quinquefida, capsula sicca.

Andromeda foliis alternis lanceolatis margine reflexis, LINN. *Lapp.* 163. *T.* 1. *f.* 3. HALL. *H.* p. 416. Andromeda *Polii folia*, LINN. *Sp.* 5. Erica *folio rorismarini*, RAII. p. 1489. In torfaceis *Hercynitis*.

ERICA. HALL. 418.

Calyx duplex, vterque quadrifolius, interior coloratus. Corolla profunde quadrifida.

Erica foliis imbricatis, glabris. HALL. *H.* 419. Erica *vulgaris, glabra*, LINN. *Sp.* 1. B. *Pin.* 483. I. R. H. flore interdum albo, Vbique in sterilibus. Fr.

TETRALIX. HALL. 418. TOURN. 373.

Calyx simplex coloratus, quadrifolius. Corolla breviter quadrifida, doliiforma.

Tetralix foliis hirsutis, quaternis. HALL. *G. p.* 164. HALL. *H.* 418. Erica *herbacea*, LINN. *Sp.* 3. *procumbens*. B. *Pin.* 486. Erica *corios folio* VIII. CLVS. *Hist.* I. p. 44. In torfaceis *Cellensibus*.

MYRSINE.

M. Myrsine *Africana*, LINN. *Sp.* p. 196. Vitis *Idaea Aethiopica, myrti Tarentinae folio, flore atropurpureo*. COMM. *H. L.* p. 123. *T.* 64. In *Africa*.

Flores congesti alares. Calyx exiguus, quinquepartitus, Corolla semiquinquefida maculata. Stami-
na

na quinque medio flori sub segmentis inserta. Stylus stigmatibus trifido fimbriato, extra florem pendulo. Folia utrinque ovata, subintegra, versus apicem acute serrata. Fr.

PYROLA. LINN. 490.

Petala quinque, concava. Stamina decem. Capsula pentagona.

Pyrola foliis ovatis acuminatis, serratis, spica unilaterali. HALL. *H. n.* 1. *Pyr. secunda.* LINN. *Sp.* 3. *Pyrola folio mucronato serrato.* B. *Pin.* 181. *Pyrola II. tenerior* CLVS. *Pann.* 506. In sylvis passim.

Pyrola racemosa, staminibus & pistillo rectis. HALL. *Gott.* 165. *Hort. n. 2. minor.* LINN. *Sp.* 2. *Pyrola minor.* DILL. *Cat. Gieff. p.* 94. In sylvis *Hercynicis* circa *Ilfeld* & alibi.

Pyrola staminibus adscendentibus, pistillo declinato, rotundifolia. LINN. *Sp.* 1. HALL. *H. n.* 3. *Pyrola rotundifolia maior.* B. *Pin.* 191. In sylvis circa *Eichenkrug, Hardeggen.*

Pyrola scapo unifloro. uniflora. LINN. *Sp.* 6. HALL. *H. n.* 4. *Pyrola flore singulari.* RIV. *Pentap. irreg. T.* 35. *Pyrola IV.* CLVS. *Pann. p.* 508, 509.

TAMARIX. LINN. 337. Tamariscus, HALL.

TOVRN.

Petala quinque. Stamina decem, aut quinque. Stigmata tria plumosa. Capsula triquetra, unilocularis.

Tamarix floribus decandris. germanica. LINN. *Sp.* 2. HALL. *H. p.* 419. *Tamariscus fruticosa, folio*

lio crassiore f. Germanica. B. Pin. 485. LÖB. ic. 318.
BLACKW. F. 331. Ad Rhenum, Danubium aliosque
fluvios.

OLIGOSTEMONES TRIHILATAE.

LINN.

Calyx quinquepartitus, aut quinquefolius. Petala quinque saepe ex calyce orta. Stamina vario numero quinque, septem, octo. Germen saepe nectarium cingit. Stylus simplex. Fructus bi- aut trilocularis alatus, echinatus, angulatus.

TROPAEOLVM. LINN. 421. Cardamin-
dum. TOURN. 244.

Calyx monophyllus, pone nectario conico instructus. Petala calyci inserta inaequalia. Stamina octo declinata. Stylus stigmatibus trifido. Capsulae tres angulatae congestae.

Tropaeolum foliis integris, petalis acuminato-fetaceis. *minus.* LINN. *Sp.* 1. *Cardaminum minus* & *vulgare.* REVILL. *Peruv.* 3. p. 14. F. 8. *Nasturtium indicum minus.* B. Pin. 306. LÖB. ic. 166. f. 1. 2.

ESCVLVVS. LINN. 420. Hippocastanum.
TOURN. 382. Pavia. BOERH. 260.

Petala calyci inserta. Capsula echinata trilocularis. Stylus simplex. Flores racemosi.

H. Esculus floribus heptandris. *Hippocastanum.* LINN. *Vps.* 92. *Cliff.* 142. *Castanea equina.* OLIVS.
Hist.

Hist. I. p. 7. Exotica quidem, sed nunc fere spontanea ad vias & in hortis. Folia digitata. Fr.

H. Esculus floribus octandris. *Pavia*. LINN. *Sp. 2.*
Pavia. BOERH. *Lugd. bat. 2. T. 260. Cliff. 143.* In
Carolina. Fr.

ACER. LINN. 1023.

Calyx coloratus. Petala quatuor & stamina octo
in circulo tumido balsamico, cui germen immer-
sum. Stilus stigmatibus duplici. Capsulae duae alis ma-
ximis auctae. Sexus varius.

Acer foliis semitrilobis, lobis obtuse incis-
tis. HALL. H. n. 1. Acer foliis lobatis obtuse emarginatis.
(campestre.) LINN. *Spec. 1.* Acer campestre. O-
minus. B. Pin. 431. Acer minus *Hist. I. p. 8. 9.*
Vbiq. ad fapes. Fr.

Acer foliis quinquangulis, vndique ferratis.
HALL. H. 2. Acer foliis quinquelobis inaequaliter
ferratis, floribus ramosis. (Pseudo-platanus.)
LINN. *Spec. 2.* Acer montanum candidum. B. Pin.
439. A. R. H. Acer latifolium. CLVS. *Hist. I. p. 10.*
In sylvis. Fr.

Acer foliis quinquangulis, acutissime & rariter
dentatis. HALL. H. n. 3. Gört. 168. Acer foliis
quinquelobis acuminatis acute dentatis glabris, flo-
ribus corymbosis. (Platanoides.) LINN. *Sp. 5.* Acer
maior. CAMER. *Epit. 63.* In M. Hengst.

CARDIOSPERMUM. LINN. 447.

Calyx trifolius. Petala quatuor. Nectarium qua-
drifolium circa germen. Stamina octo. Capsula in-
flata triloba.

H. *Cardiospermum*. *Halicacanthum*. LINN. *Sp.* 17
Vesicaria. REX. *Tetrap. irreg.* T. 144. *Corindrum*
ampliori folio, fructu maiore. 1. R. H. 431. In
India. A.

STAPHYLAEA, LINN. 336. *Staphylodendron*.
 TOVRN. 386.

Calyx coloratus. Petala quinque. Circulus balsamicus circa germen. Capsulae tres inflatae, connatae.

Staphylaea foliis pinnatis. *pinnatis*. LINN. *Sp.* 1.
 HALL. p. 423. *Staphylodendron*. TABERN. p. 1022.
 CAMER. *Epit.* 171. 1. R. H. In hortis & ad sepea
 fere spontanea. Fr.

BERBERIS. LINN. 399. TOVRN. 385.

Petala sex, ad basin binis glandulis instructa. Baccas disperma.

Berberis pedunculis racemosis. *vulgaris*. LINN. *Sp.*
 1. Berberis *spinis triplicibus*. CLIFF. 122. HALL.
 H. p. 242. Berberis. BLACKW. T. 165. *dumetorum*.
 1. Pin. 454. 1. R. H. In dumetis. Fr.

TELEPHIUM. LINN. 339. TOVRN. 128.

Petala quinque calyci aequalia. Capsula triquetra.

H. *Telephium*. LINN. *Sp.* p. 271. *Telephium*
Dioscoridis Imperati. *Telephium legitimum*. CLVS.
Hist. II, p. 67. In Gallo-Provincia. P.

TRAPA.

TRAPA. LINN. 146. Tribuloides.

TOVRN. 431.

Calyx quadripartitus. Petala quatuor. Capsula
ossea spinis quatuor crassis, ex calyce mutato ena-
tis, armata.

Trapa natans, LINN. *Sp. P.* 120. *Trapa petio-
li, foliorum, natantium, ventrisaosi*. CLIFF. 481.
Tribuloides HALL. *Helv.* 468. *Tribulus aquaticus*,
CAMER. *epit. p.* 715. & aliorum. In stagnis. P.

PTELEAE *similis*.

Frutex *Pteleae* affinis, foliis ternatis, crenatis,
fructu trihilato. Ita interim vocare liceat fruticem,
qui noui generis nomen forte meretur. Frutex hu-
milis vix bipedalis, ramis patulis. Folia ternata
utrinque lanceolata; medio maiori. Flores cymosi.
Calycis exigui foliola quinque angusta. Petala quin-
que ovato lanceolata, cava, patentia, calyce mul-
to maiora, ex albo viridia. Stamina quinque, fila-
mentis basi crassioribus, hirsutis. Germen intra co-
rollam retractum; stylo breui, stigmate quadrifi-
do, abicit in capsulam quadrilocularem, quatuor
extantibus latis alis auctam. Saepè calycis folia, pe-
tala & stamina quatuor; stigmata tria, capsulae alas
& loculi tres. Vides affinitatem & diversitatem a
Ptelea.

ISOSTEMONES CYMOSAE. LINN.

Flos fructui innatus. Calycis segmenta, petala
aut corollae segmenta, & stamina quinque. Stilus
brevis stigmate tumido. Fructus pulposus, aut baccat.
Flores in cymas & racemos congesti.

EVO-

EVONYMVS. LINN. 240.

Petala & stamina ex placenta communi. Stigma obtusum. Capsula pulposa, quadrangularis, quadrilocularis.

Evonymus foliis oblongo-ovatis. HALL. H. 523. n. 1. Evonymus floribus plerisque quadrifidis. (caropano.) LINN. Sp. 1. Evon. vulgaris, granis rubentibus. R. Pin. 428. GAMER. Epis. 102. Ad sepes. Fr.

CELASTRVS. LINN. 239. Evonymoides.

Fabrica priori similis: sed stigma trifidum, & capsula trigona.

H. Celastrus spinis nudis, ramis teretibus, foliis acutis, pyracanthus. LINN. Sp. 5. Lycium aethiopicum, pyracanthae folio. COMM. H. L. p. 163. T. 84. In Africa. Fr.

RHVS. LINN. 331. Toxicodendron.

Cotinus. TOURN.

Calyx erectus, petala erecta. Stamina brevissima. Stigmata tria, sessilia. Capsula monosperma.

H. Rhus foliis pinnatis, obtusiusculis serratis ovatis, subtus villosis. Coriaria. LINN. Sp. 1. Rhus corticiorum. LOB. ic. p. 11. p. 98. BOB. Pempt. 779. Sumach vulgo. In Europa australi. Fr.

H. Rhus foliis ternatis, foliolis petiolatis ovatis nudis integerrimis, caule radicante. radicans. LINN. Sp. 6. Hedera trifolia Canadensis. CORNVT. Canad. T. 97. In Virginia. Canada. Fr.

H. Rhus

α. *Rhus foliis simplicibus obovatis. Cotinus.*
 LINN. *Sp.* 42. *Coccygia. Desfontainii.* LOB. *ic.* II.
 p. 99. *Cotinus coriaria.* DOD. *Pempt.* 780. In *Au-*
stria. Italia, Servia. Fr.

VIBURNUM. LINN. 332. *Tinus. Opulus.*

.. Calyx exiguus, Petalum campanulatum, Stigma-
 ta & bacca, vix in priori.

α. *Viburnum foliis integerrimis ovatis, ramifica-*
tionibus subtus villosis glandulosis. *Tinus.* LINN. *Sp.*
 1. *Tinus.* II. *clvs. Hist.* I. p. 49. *Laurus Tinus.*
f. sylvestris. I. B. *Hist.* III. p. 418. In *Europa au-*
strali. Fr.

α. *Viburnum foliis cordatis, crenatis venosis,*
subtus tomentosis. *Lantana.* LINN. *Sp.* 5. HALL. *H.*
 462. *Viburnum vulgo.* B. *Pin.* 249. I. R. H. *Lan-*
tana DOD. *Pempt.* 701. In *Thuringia. Europa au-*
strali. Fr.

Viburnum foliis lobatis, petiolis glandulosis.
 LINN. *Spec.* 7. *Opulus* I. R. H. LINN. *Cliff.* 109.
 HALL. *H.* p. 463. *Sambucus aquatica.* B. *Pin.* 456.
 Flores in radio cymae steriles, petali segmento
 exteriori maximo.

β. Varietas floribus omnibus amplioribus, steri-
 libus, in cymas globosas congestis. *Opulus flore*
globoso. I. R. H. *Sambucus arbor rosea.*

SAMBUCVS. LINN. 334.

Petalum rotatum, bacca di- & trisperma.

1 *Sambucus caule arboreo, floribus umbellatis.*
 HALL. *H. n.* 1. *Sambucus nigra*. LINN. *Spec.* 3.
 BLACKW. *T.* 151. *Sambucus fructu in umbella nigro.* B. *Pin.* 456. Ad sepes. Fr.

1 *Sambucus caule herbaceo, floribus umbellatis.*
 HALL. *H. n.* 2. *Sambucus Ebulus*. LINN. *Sp.* 1.
 BLACKW. *T.* 488. *Sambucus humilis f. Ebulus.*
 B. *Pin.* 456. Rarius in gramineis & hortis, P.

1 *Sambucus caule arboreo, floribus racemosis.*
 HALL. *H. n.* 3. LINN. *Sp.* 4. *Sambucus racemosa rubra.* BAVH. *Pin.* 456. In fylvis v.g. *Eulenloch* & *versus Befigebausen.* Fr.

CASSINE. LINN. 333.

Corolla patens. Bacca trilocularis stigmatibus umbilicata.

H. Cassine foliis subovatis integerrimis. *Maurocena*. LINN. *Sp.* 2. *Frangula sempervirens, foliis rigido subrotundo.* DILL. *Elzb.* 146. *T.* 121. *f.* 147. *Cerasus Capensis, fructu rubro, folio fere obtuso.* PETIV. *Gaz.* *T.* 57. *f.* 4. In *Aethiopia*, Fr.

CAPRIFOLIUM. HALL. 464. Lonicera.

LINN. 210.

Petalum ex longo tubo bilabiatum, in aliis fere aequaliter quinquesidum. Stilus stigmate tumido. Bacca trilocularis.

1 H. Caprifolium foliis supremis coalescentibus, floribus bilabiatis. HALL. *Gott.* 192. *Lonicera floribus verticillatis terminalibus sessilibus, foliis summis connato-perfoliatis.* LINN. *Sp.* 1. Periclymenum

perfoliatum. B. Pin. 302. LOBEL. ic. 692.
Caprifolium italicum. DOD. *Pempt.* 411. RAV. T.
 122. In Europa australi. Fr.

H. *Caprifolium* foliis connatis, floribus tubulo-
 sis. HALL. l. v. *Lonicera speciosus nudis verticillatis*
terminalibus, foliis summis connato-perfoliatis.
 BERN. Spec. 2. *Periclymenum perfoliatum Virgi-*
nianum sempervirens & florens. HERM. *Lugd.*
 484. T. 483. *Virginianum*. RIV. *Monop.* irreg.
 T. 116. Fr.

Caprifolium floribus racemosis, foliis distinctis.
 HALL. H. n. 5. *Lonicera rapunculoides ovatis imbricatis*
terminalibus, foliis omnibus distinctis. LINN. Sp. 3.
Periclymenum non perfoliatum Germanicum. B. Pin.
 302. In sepibus & sylvis. v. g. versus Seeburg. *Offe-*
rod. Fr.

H. *Caprifolium* fructu gemino rubro, foliis gla-
 bris, cordatis. HALL. Gott. l. 6. AMM. Ruth n. 262.
Lonicera pedunculis bifloris, baccis distinctis, fo-
liis cordatis obtusis. (Tatarica.) LINN. Sp. 5. In
 Sibiria. Fr.

Caprifolium floribus geminis, foliis ovatis, la-
 nuginosi, integerrimis. HALL. Hefu. n. 4. *Lonicera*
pedunculis bifloris, baccis distinctis, foliis integ-
rimis pedifloris. (Xylosteum.) LINN. Sp. 6.
Chamaecerasus dumetorum, fructu gemino rubro.
 B. Pin. 451. In sepibus. Fr.

H. *Caprifolium* floribus geminis, foliis ovatis,
 nervosis, ciliatis. HALL. H. n. 9. *Lonicera pedun-*
culis bifloris, baccis coadunatis globosis, stylis indi-
visis.
 O 2

vifis. LINN. *Sp.* 9. Periclymenum *vestitum*, fructu
töruleo. CHEV. *Pann.* p. 85.

DIERVILLA. LINN. ed. pr.

Calyx persistens, segmentis longis, subulatis, Ca-
psula ovata, bilocularis.

Diervilla. LINN. *Cliff.* 63. T. 17. Diervilla *Aca-*
disensis fruticosa, flore luteo. TOWN. *Alt. Par.*
1706. T. 7. f. 1. Lonicera *perdis terminalibus*,
foliis serratis. LINN. *Sp.* 14. In *America septen-*
trionali.

UMBELLIFERAE.

Caulis fistulosus. Folia alterna, plerumque com-
posita. Petioli ex eodem puncto enati flores in
umbellis proferunt. Umbellis foliola subiecta invo-
lucris nomine veniunt. Semina duo ad mediam axin
sibi applicata in apice communem gerunt placentam,
ex qua evasuntur, in paucissimis calyx ex-
iguus, in omnibus quinque petala, totidem stami-
na, styli duo simplices.

Umbella obscurior.

PHYLLIS. LINN. 286.

Calyx bifolius minimus. Fructus turbinato-ob-
longus, angulatus.

H. Phyllis stipulis dentatis. LINN. *Sp.* 1. *Vale-*
rianella Canariensis frutescens simplex nobis dicta.
DILL. *Elth.* 405. T. 299. f. 386. In *Canariis*.

HYDRO-

HYDROCOTYLE. LINN. 288.

Umbella simplex. Involucrum quadrifolium. Fructus orbiculatus.

Hydrocotyle foliis peltatis orbiculatis, vndique emarginatis. LINN. Cliff. 88. HALL. H. 425. vulgaris. LINN. Sp. 1. 1. R. H. Cotyledon aquatica. LOB. ic. 387.

II. *Umbella manifesta.*

Seminibus gibbis striatis, involucre nullo.

ANETHVM. LINN. 326. TOURN. 169.

Foeniculum. T. 164. HALL. 425.

Umbella multiplex; petala aequalia. Foliorum ultimae pinnulae tenuissimae.

H. *Anethum fructibus compressis. graveolens. LINN. Sp. 1. Mat. med. 146. Anethum bortenfe. B. Pin. 147. Inter Hispaniae fegetes.*

H. *Anethum fructibus ovatis. Foeniculum. LINN. Sp. 2. Foeniculum vulgare Germanicum. B. Pin. 147. HALL. H. 425. In Helvetia; Gallia Narbonensi.*

APIVM. LINN. 329.

Umbella vniversalis radius paucioribus; peculiaris pluribus. Petala aequalia.

Apium foliolis caulinis cuneiformibus semitrifidis, ora lorum circumferrata. graveolens. LINN. Sp. 2. HALL. H. 427. Apium palustre seu officinarum. B. Pin. 154. Apium BLACKW. T. 443. β. Apium dulce

dulce, *Celleri italorum*. I. R. H. 305. et in fossis
Europae. 2. in hortis culta.

SMYRNIVM. LINN. 325.

Umbella vniversalis inaequalis, partialis erecta,
flores subaequales.

H. *Smyrniun foliis caulinis ternatis petiolatis serratis. Olusatrum*. LINN. *Sp.* 2. *Smyrniun Diof.*
B. *Pin.* 154. Ad rupes *Angliae & Galliae*.

AEGOPODIVM. LINN. 330. Podagraria.

HALL. 427.

Umbella multiplex, petala inaequalia, femina
magna.

1. *Aegopodium foliis caulinis summis ternatis*.
LINN. *Sp.* p. 265. Podagraria. HALL. l. c. RIV. T.
46. *Angelica sylvestris minor*. B. *Pin.* 115.

PASTINACA. LINN. 324.

Petala aequalia flava. Fructus compresso-planus
margine cinctus.

1. *Pastinaca foliis simpliciter pinnatis*. LINN. *Sp.* 1.
HALL. H. 446. *Pastinaca sylvestris latifolia*. B. *Pin.*
155. In gramineis.

CARVM. LINN. 327. Carvi. T. 160.

Petala subaequalia. Semina ovato-oblonga.

1. *Carum*. LINN. *Sp.* 263. *Mat. med.* 138. HALL.
H. 428. CAMER. *Epis.* 516. *Carum pratense, Carvi officinarum*. B. *Pin.* 158. Pinnulae circa costam se
decussant. In pratis.

PIM-

PIMPINELLA. LINN. 328. *Tragofelinum*. T.

Vmbella laxa. Petala subaequalia.

Pimpinella foliis pinnatis : foliolis radicalibus subrotundis ; summis linearibus. *saxifraga*. LINN. Sp. 1. *Tragofelinum pinnis semibilobis, circumseratis*. HALL. H. n. 1. *Pimpinella saxifraga maior*. B. Pin. 264. I.

β. *Pimpinella (Tragofelinum)* pinnis laciniatis, superioribus tenuissime divisis. HALL. H. n. 3. *Pimpinella saxifraga minor*. B. Pin. 150. In locis siccis. Num varietas prioris.

γ. *Pimpinella* foliis radicalibus trifidis incis. *Anisum*. LINN. Sp. 4. *Anisum vulgare*. OLVS. Hist. 2. p. 202. BLACKW. T. 374. *Anisum herbariis*. B. Pin. 159. In *Aegypto*. A.

δ. Seminibus gibbis striatis, involucro alterutro aut utroque.

SESELI. HALL. LINN.

γ. *Seseli* foliis glaucis, foliolis longe linearibus, radice praelonga, seminibus subhirsutis. HALL. Comm. Soc. Gott. T. II. p. 347. vbi vide fusio-rem descriptionem. GMEL. Fl. Sibir. T. I. p. 207.

OENANTHE. LINN. 314.

Vmbella difformis. Flores radii *mares* inaequales, maiores, disci *androgyni* aequales. Fructus calyce coronatus.

Oenanthe stolonifera, foliis caulinis planatis fistulosis teretibus. *fistulosa*. LINN. *Sp.* 1. HALL. *H.* 432. n. 1. *Oenanthe aquatica*. B. *Pin.* 162. RIVIN. 66. in fossis & pratis v. g. versus *Maschmühle*. P.

H. *Oenanthe umbellularum* pedunculis marginalibus longioribus ramosis masculis. *prolifera*. LINN. *Sp.* 3. *Oenanthe prolifera apula*. B. *Pin.* 163. MORIS. *Hist.* III. p. 289. S. 9. T. 7. f. 5. A.

H. *Oenanthe* fructibus globosis. *globulosa*. LINN. *Sp.* 4. *Oenanthe Lusitanica*, semine crassiori globoso. I. R. H. 313. P.

H. *Oenanthe* foliolis radicalibus ovatis incis, caulibus integris linearibus longissimis simplicioribus. *pimpinelloides*. LINN. *Sp.* 5. *Oenanthe apiifolia*. B. *Pin.* 162.

PHELLANDRIVM. LINN. 315.

Flores inaequales. Fructus ovatus, fere glaber, calyce coronatus.

Phellandrium foliorum ramificationibus divaricatis. LINN. *Sp.* 1. Phellandrium foliis ad angulos obtusos ramosis. HALL. *H.* p. 432. RIV. *T.* 64. In fossis.

CORIANDRVM. LINN. 318.

Flores radii maiores inaequales androgyni, fructu sphaerico calyce coronato. Flores disci mares minores.

H. *Coriandrum* fructibus globosis. *sativum*. LINN. *Sp.* 1. *Mat. med.* 135. *Coriandrum maius*. B. *Pin.* 158. RIV. *T.* 69. 70. In Italia. A.

H. *Coriandrum* fructibus didymis. *testiculatum*.
LINN. *Sp.* 2. *Coriandrum sylvestre foetidissimum*. B.
Pis. 158. Prope *Vlmam*. A.

AETHVSA. LINN. 317. *Cynapium*. RIV.

Involucrum parziale trifolium, laterale, reflexum. Petala inaequalia.

Aethusa Cynapium. LINN. *Sp.* p. 256. *Cicuta-
ria fatua*. LOBEL. ic. 280. *Cynapium*, RIV. *T.* 75.
In hortis.

AMMI. LINN. 297.

Involucrum univernale pinnatifidum.

H. *Ammi* foliis inferioribus pinnatis lanceolatis
serratis, superioribus multifidis linearibus. LINN. *Sp.*
1. *Ammi maius*. B. *Pin.* 159. *Ammiofelinum*. RIV.
T. 84. In *Europa* australi. A.

CONIVM. LINN. 299. *Cicuta*,
TOVRN.

Involucrum utriusque foliola lata. Fructus subrotundus striatus.

Conium seminibus striatis. *maculatum*. LINN. *Sp.*
1. *Cicuta*. BLACKW. *T.* 451. HALL. *H.* p. 434.
Cicuta maior. B. *Pin.* 160. *vulgaris*. CLVS. *Hist.* II.
p. 200. In ruderalis, ad sepes.

SIVM. HALL. 435. LINN. 310. *Cicuta*,
LINN. 316.

Involucrum univernale latifolium reflexum. Petala aequalia.

1 *Sium* foliis pinnatis, serratis, umbella terminali. *latifolium*. LINN. *Sp.* 1. HALL. *H. n.* 1. C. B. *Pin.* 154. I. R. H. *Sium*. DODON. *Pempt.* 589. RIV. *T.* 77. In fossis ad loca salsa. v. g. prope *Harste*; & *Salz der Helden*.

1 *Sium* foliis pinnatis, umbellis axillaribus sessilibus, nodiflorum. LINN. *Sp.* 2. *Sium foliis pinnatis, inferioribus simplicibus, superioribus trifidis, argute dissectis*. HALL. *H.* 2. *Sium aquaticum ad alas floridum*. I. R. H. In fossis.

• H. *Sium* foliis pinnatis, floralibus ternatis. *Sisarum*. LINN. *Spec.* 3. *Sisarum Germanorum*. B. *Pin.* 155. *Sisarum*. RIV. *T.* 77. A.

1 *Sium* pinnis multifidis, serratis, nervo folioso. HALL. *H.* 6. *Sium foliolis linearibus decurrentibus connatis*. LINN. *Spec.* 6. *Falcaria*. RIV. *T.* 47. In agris super *Weende*.

1 *Sium* foliis serratis, pinnulis tripartitis. HALL. *H. n.* 5. *Cicuta umbellis folio oppositis, petiolis marginatis obtusis. (virosa.)* LINN. *Sp.* 1. *Cicuta aquatica*. WEPFER in proprio libro *T.* 1-4. LINN. *Lapp.* 103. In lacu ad *Seeburg* & *Denckershausen*.

SISON.

Sison foliis ternatis. *Canadense*. LINN. *Sp.* 3. HALL. *Gott.* 175. *Myrrhis Canadensis trilobata*. MORIS. *Hist.* III. p. 301. S. 9. *T.* 11. f. 4. Huic speciei involucrium utrumque deest.

LIGUSTICVM. LINN. 308.

1 H. *Ligusticum* foliis multiplicibus, foliolis superne incis. *Levisticum*. LINN. *Sp.* 1. *Mat. med.*

119. *Ligusticum vulgare*. B. Pin. 157. In montibus *Liguriæ*. P.

H. *Ligusticum foliis biternatis. Scoticum*. LINN. Sp. 2. *Ligusticum humilius Scoticum, apii folio*. I. R. H. PLVKN. Alm. 217. T. 96. f. 2. *Seseli maritimum Scoticum*. HERM. Parad. T. 227. RIV. T. 68. In *Scotia*.

ATHAMANTA. LINN. 301. Meum

TOURN. 165.

Athamanta foliolis capillaribus, seminibus glabris, striatis. Meum. LINN. Sp. 8. Meum. HALL. p. 426. *Meum vulgare, foliis Anethi*. B. Pin. 148. In sylva *Hercynica* prope *Andreasberg & Altenau*.

BUPLEVRVM. LINN. 291. HALL. 436.

Involucrum universale interdum deest: parziale quinquefolium aut quinquefidum. Semina profunde sulcata. Flores lutei. Folia simplicia.

Bupleurum foliis ovatis perfoliatis. HALL. H. 437. *Bupleurum involucris universalibus nullis, foliis perfoliatis*. LINN. Sp. 1. *Perfoliata vulgatissima arvensis*. B. Pin. 277. BLACKW. T. 95. Vbique inter segetes.

Bupleurum foliis imis petiolatis, superioribus ovatis perfoliatis. HALL. H. n. 2. *Bupleurum longifolium*. LINN. Sp. 5. *Perfoliata montana latifolia*. B. Pin. 277. In sylva prope arcem *Pless*.

Bupleurum foliis petiolatis falcato-ellipticis acutis. HALL. H. n. 11. *Bupleurum falcatum*. LINN. Sp. 6. *Bupleurum longis & angustis foliis incurvis*. H.

H. Ox. III, p. 300. *Über der Papiermühle & in M. Altenßollberg.*

H. Bupleurum foliis gramineis, calyce peculiari quinquefido aristato rigido. *HALL. H. n.* 6. *Bupleurum odontites.* *LINN. Sp.* 7.

CRITHMVM. *LINN.* 303.

H. Crithmum foliolis lanceolatis carnosis. *maritimum.* *LINN. Sp.* 1. *Crithmum.* *f. Foeniculum marinum.* *B. Pin.* 288. *LOB. ic.* 392. Ad littora maris.

Seminibus, gibbis, striatis, margine alatis.

PEVCEDANVM. *LINN.* 302.

H. Peucedanum foliis quinquies tripartitis filiformibus linearibus. *officinale.* *LINN. Sp.* 1. *Mat. med.* 126. *Peucedanum germanicum.* *B. Pin.* 149. In *Germania* australi.

H. Peucedanum foliis pinnatifidis, foliolis longe trifidis oppositis. *LINN. Sp.* 3. *Seseli pratense, Silaus forte Plinio.* *B. Pin.* 162. *Siler pratense.* *DOD. Pempt.* 310. Super *Wernigerode.*

H. Peucedanum foliis duplicato pinnatis, pinnulis longis linearibus, dissitis. An *Carvisolia.* 1. *B. III.* p. 171. Descripsi hanc plantam in *epistola* ad III. *WERLHOFFIUM Observ. bot.* p. 12. Caulis striatus. Petioli foliorum membranacei longe amplexicaules. Umbella utraque densissima. Involucrum utrumque decaphyllum, radiis appressum, iisque brevius. Petala alba, extus flava. Semina convexa striata, alba cincta. Circa *Berolinum* in sylvis vliginosis.

SELINUM. HALL. 443. LINN. 300.

Selinum sublaetescens, radice vnica, *palustre*, LINN. *Sp.* 2. HALL. *H.* 3. *Thysselinum angustifolium*. RIV. *T.* 20. *Seseli palustre laetescens*. B. *Pin.* 162. *Prodr.* 85. *In Landolfshäuser Sumpf.* Circa *Berolinum*.

Selinum pinnis pinnatis, pinnulis appendiculatis, circumscissatis, triangularibus, HALL. *Helv.* n. 41. *Oreoselinum apii folio*. In praecipitiis M. *Plessberg*.

Selinum caule alato, foliolis bi- & trifidis. *Angelica foliis duplicato pinnatis, foliolis angustis, saepe trifidis*. BOEHM. *Lips* 107. *An Peucedanum*. HALL. 3. *Angelica tenuifolia*. RIV. *T.* 18. DIEL. *Nov. gen.* p. 226. Caulis inferius alis membranaceis notatus. Foliola apice alba. Involucrum universale nullum, aut unifolium, partiale multifolium lineare, petiolis brevius. Semen tribus sulcis profunde exaratum, margine alatum.

ANGELICA. LINN. 309.

H. *Angelica* foliorum impari lobato. *Archangelica*. LINN. *Spec.* 1. *Mat. med.* 120. *Angelica sativa*. B. *Pin.* 155. *Archangelica*. CLVS. *Hist.* 2. p. 295. *Vulgaris in hortis. Sponte in alpi- bus Lapponticis.*

Angelica foliis aequalibus ovato-lanceolatis serratis. *sylvestris*. LINN. *Sp.* 2. HALL. *H.* 445. *Angelica sylvestris maior*. BAVH. *Pin.* 155. *Vbique in pratis.*

H. An-

H. *Angelica foliolis aequalibus, ovatis, inciso-ferratis.* LINN. *Sp.* 4. HALL. *Gart.* 179. *Angelica lucida Canadensis.* CORNVT. *Canad.* 196. *T.* 197. In *Canada.*

IMPERATORIA. LINN. 321.

H. *Imperatoria. Ostruthium.* LINN. *Spec.* 259. RIV. *T.* 7. LOB. *sc. p.* 700. BLACKW. *T.* 279. B. *Pin.* 156. *Magistrantia.* CAMER. *Epit.* 592. In *alpi-
bus Helveticis.*

Seminibus quatuor & pluribus alis armatis.

ASTRANTIA. LINN. 290. HALL. 439.

*Umbella simplex, involucri aequalis. Flores
males & androgyni. Semen calyce coronatum.*

1 *Astrantia foliis quinquelobis, lobis trifidis. ma-
ior.* HALL. *H.* 439. *Astrantia.* RIV. *T.* 67. *Astran-
tia maior, corona floris purpurascens.* I. R. H. In
M. *Altenstollberg.* P.

LASERPITIVM. LINN. 306.

HALL. 440.

1 *Laserpitium foliolis cordatis, inciso-ferratis. lati-
folium.* LINN. *Sp.* 1. HALL. *H.* n. 1. *Libanotis
latifolia maior.* C. B. *Pin.* 159. *Seseli Aethiopicum
herba.* DOD. *Pempt.* 312. In M. *Altenstollberg,* circa
Elbingerode, super Wernigerode. P.

H. *Laserpitium foliolis lanceolatis integerrimis
petiolatis. Siler.* LINN. *Sp.* 6. *Ligusticum, quod
Seseli officinarum.* B. *Pin.* 162. *Siler montanum.*
DOD. *Pempt.* 310. CLVS. *Hist.* CXEV. *Seseli*
BLACKW. *T.* 426. In *alpi-
bus Helvetiae.* P.

H. *LasERPitium* foliolis lanceolatis integerrimis, extremis coalitis. *Prutenicum*. LINN. *Sp.* 4. *LasERPitium foliis radicalibus patentissimis*. ROYEN. 103. *LasERPitium dancoides Prutenicum, viscoso semine*. BREYN. *Cent.* 167. *T.* 84. In *Marchia, Borussia*. Circa *Jenam*.

Seminibus plantis margine cinctis.

SPHONDYLIVM. HALL. 447.

LINN. 307.

Sphondylium pinnatum, pinnis hirsutis semiquinquefidis. HALL. *H. n.* 1. *Heracleum Sphondylium*. LINN. *Sp.* 1. *Sphondylium vulgare hirsutum*. BAYH., *Pin.* 157. *Branca* vrsina TABERN. p. 92. Vique in pratis.

β. *Sphondylium* foliis angustioribus. RIV. *T.* 5. HALLERVS a priori distinguit. *Enum. Gott.* 148.

TORDYLIVM. LINN. 293.

Semina plana margine crenulato.

H. *Tordylium* involucris umbella longioribus. *Syriacum*. LINN. *Sp.* 1. *Tordylium Syriacum*. RIV. *T.* 3. *humilius, semine granulato maiore*. MORIS. *Hist.* III. p. 317. f. 9. *T.* 16. f. 7. In *Syria*. A.

H. *Tordylium* umbellulis remotis, foliis pinnatis, pinnis subrotundis laciniatis. *apulum*. LINN. *Sp.* 3. *Tordylium apulum minimum*. COLUMN. *Ecpbr.* l. p. 122. *T.* 124. *Seseli creticum minimum*. B. *Pin.* 161. In *Italia*.

H. Tor-

H. *Tordylium umbella conferta radiata*, foliolis lanceolatis inciso-ferratis. (*maximum*.) LINN. *Sp.*

g. *Tordylium*. RIV. *Pentap.* I.

Seminibus aculeatis.

CAUCALIS. HALL. 448. LINN. 294.

Corolla universalis radiata.

1. *Caucalis involucris latifoliis*, petalo extremo maximo. HALL, n. 1. *Caucalis involucris singulis pentaphyllis*: foliolo unico duplo maiore. (*grandiflora*.) LINN., *Sp.* 1. *Caucalis arvensis echinata*, magno flore. B. *Pin.* 152. *Echinophora pycnocarpus*. COLVMN. *Ecpbr.* l. p. 91. T. 94. Inter segetes circa Webnde. A.

H. *Caucalis seminibus cylindricis asperis*. HALL. G. 181, cum descr. *Anthriscus Creticus*. RIV. T. 38.

1. *Caucalis involucro universalis nullo*, laciniis foliorum perangustis. HALL. H. n. 2. *Caucalis seminum aculeis triglochidibus uncinatis*; pilis verticillatis bispidis. (*leptophylla*.) LINN. *Sp.* 6. *Caucalis arvensis echinata parvo flore & fructu*. B. *Pin.* 152. In agris.

1. *Caucalis umbella universalis trifida*, partialibus pentaśpermis, foliis pinnatis, pinnis ferratis. LINN. *Cliff.* 91. ROY. *Lugdb.* 95. *Tordylium latifolium*. LINN. *Sp.* 5. *Caucalis arvensis echinata latifolia*. B. *Pin.* 152. In agris.

1. *Caucalis seminibus hispidis*, foliis hirsutis obruse lobatis. HALL. *Gott.* 181. *Helv.* 449. *Tordylium Anthriscus*. LINN. *Sp.* 6. *Caucalis semine aspero*,
floscu-

flosculis rubentibus. BAVH. *Pin.* 152. *Prodr.* 80. In sylvis & ad sepes.

H. *Caucalis umbellis sessilibus simplicibus.* LINN. *Cliff.* 91. HALL. *Gott.* 182. *Tordylium nodosum.* LINN. *Sp.* 7. *Caucalis nodosa, echinato semine.* B. *Pin.* 153. *Prodr.* 80. In Gallia, Italia. A.

ARTEDIA. LINN. 295.

Semen in nostra alis profunde serratis percursum. Involucrum pinnatifidum.

H. *Artedia feminibus aculeatis, muricata.* LINN. *Sp.* 2. *Caucalis maior daucoides Tingitana.* HERM. *Parad. p.* 111. *cum ic.* HALL. *Gott.* 182. *Echinophora Tingitana.* RIV. *T.* 26. A.

SANICVLA. LINN. 289.

Umbellae minimae globosae. Sexus Caucalidi similis.

Sanicula foliis radicalibus simplicibus, flosculis omnibus sessilibus. *Europaea.* LINN. *Sp.* 1. *Sanicula.* RIV. *T.* 30. BLACKW. *T.* 63. *officinarum.* B. *Pin.* 319.

Seminibus hirsutis.

DAUCVS. LINN. 296.

Involucrum pinnatifidum, umbella cava.

Daucus feminibus hispidis. *Carota.* LINN. *Sp.* 1. HALL. *H.* 451. *Daucus vulgaris.* CLVS. *Hist.* II. *p.* 198. I. R. H.

β. *Daucus sativus*, radice lutea aut aetiorubente. I. R. H. 307. α. Vbique in agris. β. in hortis colitur.

H. *Daucus* seminibus nudis. *Visnaga*. LINN. Sp. 2. *Gingidium umbella oblonga*. B. Pin. 151. In Europa australi. A.

LIBANOTIS. HALL. 450.

Semina villosa. Involucra multifolia.

1. *Libanotis* pinnulis circa costam decussatis. HALL. H. 450. n. 1. *Athamanta foliis bipinnatis planis, umbella haemisphaerica, seminibus hirsutis*. LINN. Sp. 1. *Daucus montanus apii folio*. B. Pin. 150. Prodr. 77. In M. Altenstallberg.

H. *Libanotis* foliis multifidis, planis. HALL. Gest. 183.

Seminibus rostratis.

CHAEROPHYLLUM. HALL. 452.

LINN. 320.

1. *Chaerophyllum* seminibus laevibus nitidis, nervo foliorum subhirsuto. HALL. H. 2. *Chaerophyllum flosculis omnibus fertilibus, caule aequali*. LINN. Sp. 1. *Chaerophyllum sylvestre, perenne, cicutas folio*. I. R. H. RIV. T. 43. BLACKW. T. 236. *Cicutaria vulgaris*. I. B. III. p. 181. Vbique in pratis.

H. *Chaerophyllum* seminibus nitidis ovato subulatis, umbellis ad nodos sessilibus. BOEHM. Lipp. 492. *Scandix* LINN. 3. *Chaerophyllum sativum*. B. Pin. 152. In hortis colitur.

Chae-

Chaerophyllum, feminibus brevibus hispide.
HALL. H. n. 1. Scandix. LINN. Sp. 4. *Myrrhis syl-*
vestris aequicolorum. COLUMN. Ecphr. 1. p. 110.
T. 112. *Caudalis folio Chaerophylli*. RIV. T. 34.
Circa Hannoveram.

MYRRHIS. HALL. 453. Scandix *spec.*

LINN. 319.

Semina striata.

Myrrhis radice lignosa, perenni, foliis hirsutis,
feminibus flavis, obscure striatis. HALL. Gott. 184.
Myrrhis perennis alba minor, foliis hirsutis, semi-
ne aureo. HALL. Rupp. Jen. 282. T. 5. *Myrrhis*
minor. LOBEL. *Uber Wernigerode*.

Myrrhis perennis, feminibus strictis longissimis,
HALL. H. n. 4. *Chaerophyllum* LINN. Sp. 4. *Myr-*
his Brockenbergensis. THAL. p. 77. *Myrrhis pal-*
ustris. RIV. T. 50. *Cicutaria palustris latifolia*. C. B.
Pin. 161. In pratis Hercynicis.

Myrrhis caule inferius hirsuto, superius glabro, tra-
dice verna globosa, ramosa. HALL. H. 3. *Chaero-*
phyllum LINN. Sp. 2. *Myrrhis tuberosa*
coniophyllum. J. R. H. *Myrrhis foetans*. RIV. T. 149.
Ad viam versus *Eldgehausen* in dumetis.

Myrrhis pinnulis semipinnatis, obtusis, ordinari-
um decreascentibus. HALL. H. 11. *Chaerophyllum*
caule maculato scabro, geniculis tumidis. HALL.
Sp. 3. *Myrrhis annua, semina striata, laevi*. J. R. H.
315. BLACKW. T. 243.

H. *Myrrhis seminaibus profunde sulcatis, angula-*
tis, maximis. Scandix odorata. LINN. Sp. 1. Odorata
P 2 rata

rata RIV. T. 56. *Myrrhis maior odorata*. B. Pin. 160. In Italia. P.

SCANDIX. HALL. 454.

Semina ventricoso-fubulata.

† Scandix foliis omnibus tenuissimis divisis. HALL. l. c. Scandix feminibus laevibus rostro longissimo. LINN. Spec. 2. Scandix semine rostrato, vulgaris. B. Pin. 152. Pecten veneris. CAMER. epit. 302. LOB. ic. 726. In agris vbique.

H. Scandix feminibus subulatis hispidis, floribus radiatis, caulibus laevibus. australis. LINN. Sp. 5. Scandix cretica minor. B. Pin. 152. Anisomarathrum. COLUMN. Ecpbr. 1, p. 89. T. 90. In Gallia Narbonensi. Creta. A.

Seminibus receptaculo communi paleaceo insertis.

ERYNGIVM. LINN. 287. TOURN. 173.

† Eryngium ramosissimum, pinnis foliorum pinnatis, alatis. HALL. H. n. 1. Eryngium foliis radicalibus pinnatis tripartitis. (campestre.) LINN. Sp. 7. Eryngium vulgare. B. Pin. 386. CLVS. Hist. II. p. 157. BLACKW. T. 297. Ad viam versus Boven. P.

H. Eryngium foliis radicalibus ovalibus, planis, crenatis, floribus pedunculatis. planum. LINN. Sp. 3. Eryngium Pannonicum latifolium. CLVS. Hist. II. p. 158. Ad Danubium. P.

H. Eryngium foliis radicalibus subrotundis plicatis spinosis, capitulis pedunculatis. LINN. Sp. 5. Eryn-

STELLATAE.

219

Eryngium maritimum. B. Pin. 386. CLVS. Hist. II. p. 169.

Dipetalae.

CIRCAEA. LINN. 24.

Petala & stamina duo fructui biloculari innata.

Circaea caule erecto, racemis pluribus. Lutea. LINN. Sp. 1. *Circaea foliis longioribus, obiter crenatis*. HALL. H. n. 1. *Solanifolia Circaea dicta maior*. B. Pin. 168. In umbrosis circa Plessi. Gleich. Um die Quelle bey der Webender Papiermühle.

Circaea caule ascendente, racemo unico. alpina. LINN. Sp. 2. *Circaea foliis brevioribus dentatis*. HALL. H. 2. *Circaea minima*. COLUMN. Echpr. II. p. 80. In sylvis Hercynicis.

ISOSTEMONES STELLATAE.

Flos, ut in umbelliferis, duobus seminibus connatis impositus, sed corolla monopetala, limbo quadrifido, stamina quatuor, stylus stigmate bifido. Caulis quadrangularis, folia verticillata, vires adstringentes.

ASPERVLA. LINN. 113. *Sherardia*. LINN. 112. Rubeola. T. 50.

Flos nudus tubo cylindraceo.

Asperula seminibus coronatis. HALL. Gott. 186. *Sherardia foliis omnibus verticillatis, floribus terminalibus*. LINN. Sp. 2. HALL. H. 457. Rubeola arvensis

venfis repens coerulea. B. Pin. 334. Prodr. 145.
In agris vbiq. A.

H. *Asperula umbellis pilosis, foliis senis, longis & obtusis.* HALL. H. 457. *Asperula foliis senis, floribus sessilibus aggregatis terminalibus. (arvensis.)* LINN. Sp. 2. *Asperula coerulea arvensis.* B. Pin. 334. In Anglia. Belgio. Helvetia. A.

I. *Asperula foliis octonis, seminibus hispidis, ex summo caule paniculata.* LINN. Sp. 1. *Asperula foliis octonis lanceolatis, florum fasciculis pedunculatis. (odorata.)* LINN. Sp. 1. *Asperula montana odorata.* B. Pin. l. c. In sylvis. P.

H. *Asperula foliis quaternis, ovato-lanceolatis, ramis alternis.* LINN. Cliss. 33. HALL. Helv. l. c. *floribus fasciculatis terminalibus. (Taurina.)* LINN. Sp. 3. *Rubia levis Taurinensium.* LOB. ic. p. 800. In Helvetia.

I. *Asperula foliis quaternis linearibus, floribus paniculatis.* HALL. G. l. c. H. 458. *Asperula cynanchica.* LINN. Sp. 6. *Rubia quadrifolia. laevis, floribus purpurascens.* I. R. H. *Gallium mont. latifolium cruciatum.* COLV. Esph. l. p. 297.

H. *Asperula foliis plerisque binis, inferioribus ellipticis, superioribus linearibus, segmentis floris in apice glandula notatis. Plantula procumbens, diffusa. Flores extus rubent. An Asperula foliis quaternis lanceolato-linearibus, caule erecto, floribus saepe trifidis. Pyrenaica.* LINN. Sp. 5. *Rubia cynanchica saxatilis.* B. Pin. 133, Prodr. 146. Descriptam dedi in Comm. Gott. T. V.

Aspe-

STELLATAE.

231

Asperula foliis linearibus, inferioribus senis, intermediis quaternis, caule flaccido, floribus trifidis.
HALL. G. 186. HALL. Rupp. Jen. p. 8. *tindoria*.
LINN. Sp. 4. *Galium album*. III. TABERN. T. 733.
f. 1. In M. *Altenstallberg*.

CRUCIANELLA. LINN. 118.

Flos tubulosus, calyce trifolio cinctus. Segmenta apice inflexa.

H. *Crucianella erecta, foliis senis linearibus. angustifolia*. LINN. Sp. 1. *Asperula spicata, glumis floralibus maximis*. HALL. G. 187. *Rubeola angustifolia spicata*. B. Prodr. 145. Pin. 334. *Monspeli*.

H. *Crucianella procumbens, foliis quaternis lanceolatis, floribus spicatis. latifolia*. LINN. Sp. 2. *Rubia latifolia spicata*. B. Pin. 334. *Monspeli*.

GALIVM: HALL. 458. LINN. 117. Vailantia. LINN. 1019.

Corolla rotata sine tubo. Semina subrotunda.

Galium caule tereti, fulcrato, foliis octonis, glaucis & obtusis. HALL. H. n. 8. *Mollugo montana latifolia ramosa*. B. Pin. 334. *Mollugo* CLVS. Hist. p. CXLV. In sylvis.

Galium foliis octonis, glabris, acuminatis. Mollugo vulgarior. LOB. ic. 802. *Galium album vulgare*. I. R. H. Ad vias.

Galium caule recto, foliis senis, inferne canaliculatis. Galium mont. altissimum, foliis angustis,
P 4 *albi-*

albicansibus. RVPP. *Jen.* p. 5. In *M. Altenstollberg.*

Galium caule subaspero, foliis quaternis & senis, spinula terminatis. HALL. *H. n.* 1. *vliginosum.* LINN. *Sp.* 2. *Aparine palustris minor Parisensium, flore albo.* I. R. H. 114. In vliginosis.

Galium foliis quaternis obovatis inaequalibus, caulibus diffusis. *palustre.* LINN. *Sp.* 2. HALL *H. n.* 6. Galium *palustre album.* B. *Pin.* 335. In aquosis pratis

Galium foliis senis & novenis, subasperis, spinula terminatis. HALL. *H. n.* 5. Galium *album minus.* VALL. T. 78. Ad vias locis lapidosis.

Galium caule ramosissimo procumbente, foliis quinis obverse ovatis. *saxatile.* LINN. *Sp.* 8. HALL. *H.* 460. Galium *saxatile supinum molliore folio.* JussIEV. *Alt. Par.* 1714. p. 490. Super *Hardegseth* & in sylvis *bercynicis* vbique.

Galium foliis octonis confertis, gracilissimis, spica flava caulem terminante. HALL. *H. n.* 9. LINN. *Sp.* 9. Galium *luteum.* B. *Pin.* 335. BLACKW. T. 435. In arenosis. v. g. *beym Bouender Galgen.*

H. Galium procumbens, foliis senis, gracilissimis, confertis seminibus subhirsutis. HALL. *G.* 188. An *Gal. minutum* LINN. *Sp.* 9. ? In *Sibiria.*

Galium foliis quaternis lanceolatis trinerviis, caule erecto, seminibus hispidis. HALL. *H.* 2. LINN. *Sp.* 16. *Rubia prat. laevis, acuto folio.* B. *Pin.* 333. In *M. Altenstollberg.*

Galium caulis alis, foliorum ora nervoque serratis, fructu levi. HALL. *Aparine semine minore*. VAILL. *Par.* T. 4. f. 4.

Galium caulis alis, foliorumque ora nervoque serrato, fructu subaspero. HALL. G. 188. *Vaillantia* LINN. *Sp.* 2. *Aparine semine laevi*. VAILL. *Par.* T. 4. f. 3. In agris.

Galium caulis alis, foliorumque ora & nervo serratis, fructu hispido. HALL. H. 3. LINN. 19. *Aparine vulgaris*. B. *Pin.* 334. Ad fepes.

Galium foliis quaternis, flosculis in alis confertis. HALL. H. 10. *Vaillantia floribus masculis quadrifidis, pedunculis diphyllis*. LINN. *Sp.* 4. *Cruciata hirsuta*. B. *Pin.* 335. Flores mares & androgyni misti. Ad vias.

H. *Galium* seminibus exasperatis, flosculis lateralibus masculis. HALL. G. 188. *Aparine semine Coriandri saccharati* PARKINS. *Monspeli.*

H. *Galium* flosculis inter folia ovata, amplissima reflexa positis. HALL. G. 187. *Cruciata Orient. glabra humifusa*. I. R. H. *Cor.*

H. *Galium* floribus masculis trifidis, omnibus plantae partibus hispidis. *Vaillantia erecta hispida*. LINN. *Cliff.* 460. LINN. *Sp.* 1. β.

H. *Galium* floribus masculis trifidis, pedunculo communi cristato, foliis ciliatis. *Vaillantia procumbens glabra*. LINN. *Cliff.* 468. *Cruciata muralis minima Romana*. COLVMN. *Ecpbr.* I. T. 297. Vtriusque diversitatem proxime fusius exponam.

RVBIA. LINN. 19.

Calyx minimus. Flos quinquefidus. Baccae duae glabrae.

H. Rubia foliis senis. *tinctorum*. LINN. *Sp.* 1.
HALL. *H.* 462. Rubia *sylvestris aspera*. B. *Pin.* 33.
BLACKW. *T.* 326. In Europa australi.

SPERMACOCE. LINN. III. DILL.

Eltb. 227.

Calyx minimus. Capsulae bicornes connatae.

H. Spermacoce glabra, staminibus inclusis. LINN.
Sp. 1. Spermacoce *verticillis tenuioribus*. DILL.
Eltb. 370. *T.* 227. *f.* 359. In Carolina. A.

ANTHOSPERMUM. LINN. 1029.

Calyx revolutus sine petalo. Stamina receptaculo inserta. In polygamia dioicia.

H. Anthospermum. LINN. *Cliff. T.* 27. *Sp.* 7.
1058. Frutex *Africanus ambram spirans*. WALTH.
H. 24. *T.* 9. In horto planta mascula tantum adest.
In Aethiopia. Fr.

CORNUS. LINN. 139.

Flores umbellati. Bacca officulo biloculari.

1. Cornus arborea, umbellis involucrium aequantibus. *mas.* LINN. *Sp.* 2. Cornus *sylvestris mas.*
B. *Pin.* 447. CAMER. *Epir.* 158. In M. Hengst
prope Lützelengen.

1. Cornus arborea cymis nudis. *sanguinea*. LINN.
Sp. 3. Cornus *umbellis involucrio multoties longioribus.*

ribus. HALL. H. 463. *Comus femina.* B. Pin. 447.
Ad sepes.

ISOSTEMONES CONGLOMERATAE.

Stellatis affines sunt flore quadrifido, quatuor staminibus, & fructu saepe biloculari, qui in aliis sub flore, in aliis intra florem est. Flores semper aut in spicas aut alas congesti.

SANGVISORBA. LINN. 136. Pimpinella. HALL.
469. Poterium. LINN. 948.

Corolla persistens. Fructus bilocularis. Sigmata duo plumosa.

Sanguisorba tetrastemon, staminibus tubo floris aequalibus, spicis ovalibus. Pimpinella. HALL. l. c. n. 1. Sanguisorba officinalis. LINN. Sp. 1. Pimpin. sanguisorba maior. B. Pin. 160. Prope Elbingerde.

h. Sanguisorba tetrastemon, staminibus tubo floris, longioribus, spicis cylindricis. An Pimpinella tetrastemon, spica ovata, foliis ovatis serratis. HALL. G. 197. Pimpinella minore di Canada. ZANNON. T. 138. Flores rubri, spica cylindrica, flores superiores prius florere incipiunt, ut in priori specie. Diversissima a sanguisorba Canad. spicis longissimis.

Sanguisorba polystemon, staminibus longissimis pendulis. Pimpinella. HALL. n. 2. BLACKW. T. 413. Poterium sanguisorba. LINN. Sp. 1. Pimpinella sanguisorba minor laevis. B. Pin. 160. In siccis. P.

h. San-

H. Sanguisorba *polystemon*, spinis ramosis. *Pimpinella ramosa*, *spinosa*. HALL. G. 197. MORIS. III. S. VIII. T. 18. f. 5. *Poterium spinosum*. LINN. Sp. 3. In *Creta*.

PLANTAGO. LINN. 133. Psyllium
TOURN.

Flores spicati, fructus intra florem.

1. *Plantago foliis ovatis glabris, nudo scapo tereti, spica flosculis imbricatis. maior*. LINN. Sp. 1. HALL. H. n. 5. Pl. *latifolia sinuata*. C. B. 1. R. H. *septinervia*. BLACKW. T. 35. In cultis.

β. *Plantago latifolia rosea*, foliis inter flores longioribus enatis.

H. *Plantago foliis latissimis, ovatis dentatis, rigidioribus, spica brevi, densa*. AMMAN. n. 23. HALL. G. 199.

1. *Plantago foliis ovato-lanceolatis pubescentibus, spica cylindrica, scapo tereti. media*. LINN. Sp. 3. HALL. *Helv. n. 4*. Pl. *latifolia incana*. B. Pin. 189. Ad vias.

H. *Plantago foliis lanceolato-ovatis pubescentibus subdenticulatis, spicis laxis pubescentibus, scapo angulato. Virginica*. LINN. Sp. 4. Pl. *media incana*, *Virginiana*, *serratis foliis*. MORIS. III. S. 8. T. 15. f. 8. In *Virginia*. A.

1. *Plantago foliis lanceolatis, spica subovata nuda scapo angulato. lanceolata*. LINN. Sp. 5. Pl. *angustifolia maior*. B. Pin. 189. BLACKW. T. 14. In gramineis.

H. Plan-

H. *Plantago foliis lanceolatis, spica ovata his-
ta, scapo angulato. Lagopus, LINN. Sp. 6. Pl. angu-
stifolia paniculis lagopi. B. Prodr. 98. MORIS.
III. S. 8. T. 16. f. 13. In Europa australi.*

H. *Plantago foliis semicylindraceis integerrimis
basi lanatis, scapo tereti. maritima. LINN. Sp. 10.
Pl. marit. minima, folio gramineo, rigido. I. R. H.
In falsis. v. g. am gesalzenen See.*

H. *Plantago foliis linearibus pinnato-dentatis,
scapo tereti. LINN. Sp. 10. Coronopus sylvestris
birsutor. B. Pin. 190. Eodem loco cum priori.*

H. *Plantago caule ramoso, foliis integerrimis,
spicis foliosis. LINN. Sp. 15. HALL. H. n. 1. Psyl-
lium maius erectum. B. Pin. 191. In Marchia.*

*Plantago (Psyllium) perennis, angustifolia & su-
pina. I. R. H. Psyllium. CAMER. Epit. 811.*

H. *Plantago caule ramoso, foliis dentatis, spicis
aphyllis. Cynopr. LINN. Sp. 16. Psyllium Diosco-
ridis vel Indicum, foliis crenatis. B. Pin. 191.*

BASELLA. LINN. 343.

Corolla sexfida succulenta facta semen fovet.

H. *Basella foliis planis. rubra. LINN. Sp. 1. Fl.
Zeyl. 119. In India.*

CVSCVTA. LINN. 156.

*Planta aphylla, parasitica. Flores in glomeris
congesti.*

*Cuscuta floribus sessilibus, repens filiformis,
LINN. Sp. 1. Cuscuta maior. I. R. H.*

β. Epi-

β. *Epithymum*. f. *Cuscuta minor*. G. B.

ILEX. LINN. 158. *Aquifolium*. T.

Flores quadrisidi ad alas foliorum congesti. Styli quatuor.

Ilex foliis ovatis acutis spinosis. LINN. *Sp.* 1. *Aquifolium*. HALL. *Helv.* 467. BLACKW. *T.* 205. In arenosis *Cellensibus*.

ISOSTEMONES CAMPANACEAE.

Calyx quinquefidus. Corolla plerumque campanulata. Stamina tot, quot segmenta floris. Stylus simplex. Fructus in aliis sub flore, in aliis intra florem, quae autem ob summam analogiam a se invicem divelli nequeunt.

CONVOLVULVS. LINN. 198.

Petalum plicatum, limbo leviter quinquefido. Stigmata duo.

Convolvulus foliis sagittatis utrinque acutis, pedunculis unisporis bisporisque. *arvensis*. LINN. *Sp.* 1. HALL. *H.* 489. n. 2. B. *Pin.* 294. Volubilis *arvensis*, flore roseo. TABERN. 877. In arvis ubique.

Convolvulus foliis sagittatis, postice truncatis, pedunculis unisporis. *sepium*. LINN. *Sp.* 2. *Conv. maior, albus*. Y. K. H. BLACKW. *T.* 38. In sepibus.

Convolvulus foliis sagittatis, postice truncatis, pedunculis remosis bisporis. LINN. *Sp.* 3. *Mat. med.* 82. Convolvulus *Syriacus*. MORAS II, *S. L. T.* 3.

f. 5. *Scammonia Syriaca*. B. Pin. 124. DOD. Purg. 206. In Syria.

H. *Convolvulus* foliis ovato-lanceolatis, duobus foliis flori substratis. HALL. G. 1205. ROYEN. 428. LINN. Sp. 17. *Convolv. Siculus minor, flore parvo auriculato*. BOCC. Sic. 89. T. 48. In Sicilia.

H. *Convolvulus* foliis lanceolato-ovatis glabris, caule declinato, floribus solitariis, tricolor. LINN. Sp. 25. *Conv. peregrinus coeruleus*. B. Pin. 295. In Europa australi.

H. *Convolvulus* foliis lanceolatis obtusis sericeis, pedunculis multifloris. LINN. Sp. 21. *Conv. major rectus, Creticus argenteus*. MORIS. II. S. I. T. 3. f. 1. *Conv. serpens maritimus spicae folio*. LOB. ic. 622. P.

H. *Convolvulus* calycibus tuberculatis pilosis, foliis cordatis. LINN. Sp. 6. *a. coeruleus folio subrotundo*. DILL. Elth. 97. T. 82. f. 94. Stigma trifidum.

H. *Convolvulus* calycibus tuberculatis pilosis, folio anguloso. LINN. Sp. 6. *ζ. η.* DILL. Elth. 96. T. 80. f. 90. 91. Stigma obtusum.

IPOMOEAE. LINN. 199. Quamoclit. T.

Petalum infundibuliforme. Stigma capitatum.

H. *Ipomoea* foliis cordatis acuminatis, basi angulatis, pedunculis multifloris. *coccinea*. LINN. Sp. 4. *Quamoclit Amerit. folio hederae, flore coccineo*. COMMEL. rar. 21. T. 21. A.

H. Ipo-

H. *Ipomoea* foliis pinnatifidis linearibus, floribus solitariis. LINN. *Sp.* 1. *Quamoclit. f. Jasminum americanum.* CAMER. *Hort. f.* 40. CLVS. *Posth.* 9. RON. *Bald. p.* 71. In India.

H. *Ipomoea* foliis trilobis cordatis, pedunculis unisporis. AN LINN. *Sp.* 12. Folia accurate conveniunt, sed pedunculus trifloros dicit.

STRAMONIVM. TOURN. 43. *Datura.*
LINN. 218.

Calyx circabasin rotunde deciduus. Petalum pliatum. Fructus hispidus.

1. *Stramonium (Datura)* pericarpis spinosis erectis ovatis. LINN. *Sp.* 1. HALL. *H.* 489. BLACKW. *T.* 313. *Solanum manicum Dioscoridis.* COLUMN. *Phytobas. p.* 47. *Solanum foetidum, pomo spinoso, oblongo.* C. B. *Pin.* 168. Sponte nunc in hortis & fimeris. β. flore & caule violaceo.

H. *Stramonium* calyce pentagono, fructu pendulo spinoso globoso. HALL. *Gott.* 206. *Datura* LINN. *Spec.* 2. *Stramonium feron.* BOCCON. *Sic.* p. 5.

H. *Stramonium* fructu pendulo, spinoso, calyce caupaniformi. HALL. *Gott. l. c.* *Nux metella.* CAMER. *epit.* 173.

H. *Stramonium* calyce campaniformi, fructu pendulo papillato. HALL. *l. c. c. deier.*

H. *Stramonium Aegyptiacum*, flore pleno, intus albo, extus violaceo. 1. R. H. *Datura Contarena.* P. ALP. *exot.* 167. *Datura Aegyptiaca.* VESL. p. 58.

MIRA-

MIRABILIS. LINN. 215. Jalappa.

TOURN.

Petalum infundibuliforme ex nectario globoso
ortum.

H. Mirabilis foliis corollisque glabris. corollis
folio brevioribus. Mirabilis *Jalappa*. LINN. *Sp.* I.
Admirabilis *Peruviana*. *Stv. s. Hist.* II. p. 87.

H. Mirabilis foliis villosis viscidis, tubo floris cy-
lindrico, villoso, foliis longiore. COMM. *Soc. Gart.*
T. V. ubi novam hanc speciem, iconem addita, de-
scripsi. Eodem tempore eiusdem descriptionem &
iconem edidit Ill. LINNAEVS in *Act. Soc. Holm.*
1755. p. 176. T. VI. eandemque habet in *Cent. I.*
Plant. n. 18.

PLUMBAGO. LINN. 196.

Calyx hispidus. Petalum infundibuliforme. Val-
vulae circa germen staminiferae.

H. Plumbago foliis petiolatis. LINN. *Sp.* 2. *Lych-*
itis indica spicata, ocymastri foliis, fructibus sap-
pacels oblongis. COMMEL. *H. H.* p. 169. T. 5. in
India.

POLEMONIUM. LINN. 196.

Campanulae similis, sed fructus intra florem.

H. Polemonium calycibus corollae tubo longio-
ribus. *coerulescens*. LINN. *Sp.* I. *HALLEY H.* 490.
vulgare coerulescens & album T. R. H. In *Hoffm.*
Helvetia.

CAMPANULA. LINN. 201.

Valvae latiores staminiferae fundum floris claudunt. Stigma trifidum.

1. Campanula caule angulato, foliis petiolatis, calycibus ciliatis, pedunculis trifloris. *Trachelium*. LINN. Sp. 18. HALL. H. n. 1. Camp. *vulgarior urticae folio maior vel asperior*. B. Pin. 94. In sylvis. Folia subcordata. P.

Campanula foliis cordato-lanceolatis, alis unifloris, floribus secundis sparsis. LINN. Sp. 14. HALL. H. 2. Camp. *hort. rapunculi radice*. B. Pin. 94. Ad vias sylvarum.

II. Campanula foliis ovatis glabris subserratis, caule erecto paniculato, ramulis brevibus. LINN. Sp. 8. Camp. *pyramidalis altissima*. VI. R. H. BAVH. Pin. 93.

H. Campanula capsulis quinquelocularibus, rectis calycis sinibus reflexis. LINN. Spec. 22. Campan. *bertensis, folio & flore oblongo*. B. Pin. 94. *Viola Mariana*. DOD. Pempt. 163. β. Varietas flore albo & γ. pleno.

H. Campanula capsulis trilocularibus, rectis calycis sinibus reflexis, floribus maculosis pendulis. *Sibirica*. LINN. Sp. 23. HALL. G. 193. In *Siberia*. P.

1. Campanula floribus sessilibus latioribus foliis insidentibus, capitulo terminali. HALL. H. 4. *glomerata*. LINN. Sp. 19. Camp. *pratensis flore conglomerato*. B. Pin. 94. *Rapunculus umbellatus* &c.

THAL.

THAL. n. VIII n. 2. In locis aridis circa *Harper-
hausen*.

Campanula hispida, racemo ovato-oblongo ter-
minali, caule simplicissimo, foliis lanceolato-linea-
ribus. LINN. Sp. 211. *Camp. foliis Echid. n. Pin.
94. Prodr. 36.* In sylvis montosis versus *Duer-
stet. Montheim. circa Rheinhausen.*

Campanula foliis undulatis, radicalibus lanceo-
lato-ovalibus, panicula coarctata. LINN. Sp. 5.
HALL. H. n. 11. *Rapunculus esculentus. B. Pin. 92.*
In colle arenoso prope *Bibunden.*

Campanula foliis radicalibus obovatis, caulinis lan-
ceolato-linearibus subferratis, floribus latis. *Perfici-
folia.* LINN. Sp. 7. HALL. H. n. 12. *Rapunculus
persicifolius magno flore. B. Pin. 93.* flore coeruleo
& albo. In dumetis. P.

Campanula foliis radicalibus reniformibus, cau-
litis linearibus integris. LINN. Sp. 3.
HALL. Herb. 14. *Camp. minor rotundifolia vulgaris.*
B. Pin. 93. In gramineis. P.

Campanula fructu prismatico longissimo, flore
rotato. HALL. H. n. 17. *Camp. spetulum Veneris.*
LINN. Sp. 27. *Camp. arvensis procumbens, aut cre-
sta.* I. R. H. *Viola pentagonia.* TABERN. 319. In
segetibus M. *Heynberg.* A.

H. Campanula foliis perfoliatis, floribus inferioribus
apetalis. HALL. G. 195. LINN. Sp. 31. *Camp.
pentagonia perfoliata.* MORIS. S. V. T. 2. f. 23. In
Virginia. A.

PHYTEUMA. LINN. 293. *Rapunculus*
TOURN.

Corolla rotata in lacinias lineares profunde fissa, reliqua prioris.

1. *Phyteuma spica oblonga, foliis radicalibus cordatis, superioribus angustis.* LINN. *Sp.* 4. *Rapunculus.* HALL. *H.* 5. *Rapunc. spicatus, flore floescente.* I. R. H. In sylvis. 3. Varietas spica ovali obesa in *M. Bructero.*

1. *Phyteuma capitulo subrotundo, foliis ferratis, radicalibus cordatis.* LINN. *Sp.* 3. *Rapunculus.* HALL. *H.* 3. *Rapuntium mont. corniculatum.* COLV. *Ecphr.* I. T. 224. *Rapunc. spica orbiculari.* B. *Pin.* 92. In pratis prope *Elbingerode.*

JASIONE. LINN. 896.

Flores in capitulum collecti, rapunculi similes. Antherae connatae.

1. *Jasione.* LINN. *Sp.* p. 928. HALL. *H.* 496. *Rapunculus scabiosae capitulo coeruleo.* B. *Pin.* 92. In ficcioribus prope *Bovenden. Rheinbausen. Seeburg.*

LOBELIA. LINN. 897.

Floris tubus ad basin fissus, limbo fere ringente. Antherae connatae.

H. *Lobelia caule erecto, foliis lanceolatis ferratis, spica terminali.* *Cardinalis.* LINN. *Sp.* 10. *Rapuntium coccineum.* MORIS. *Hist.* II. S. V. T. 5. f. 54. RIVIN. *Monop. irreg.* T. 107. Flos fere tripetalos, calycis segmenta tubo aequalia. In *Virginia.* P.

H. LO-

H. *Lobelia* caule erecto, foliis ovato-lanceolatis
 crenatis, floribus lateralibus, calycum sinibus re-
 flexis. LINN. Sp. 11. *Rapuntium Amer. flore dilu-
 te coerulesc.* I. R. H. 163. In *Virginia*.

H. *Lobelia* caule erecto, foliis ovatis subserratis
 pedunculo longioribus, capsulis inflatis. *inflata*.
 LINN. Sp. 12. *Hort. Vps.* 276. A.

H. *Lobelia* caulibus procumbentibus, foliis lan-
 ceolatis serratis, pedunculis lateralibus. LINN. Sp.
 17. *Campanula Afr. Erini facte, caulibus procum-
 bentibus.* HERM. *Lugdb. p.* 109. T. 52. In
Africa. A.

TRACHELIUM. LINN. 204.

Corolla infundibuliformis, Stylus corolla multo
 longior. Flores corymbosi.

H. *Trachelium* (foliis ovatis serratis, corymbis
 compositis.) LINN. Sp. p. 171. ROY. *Lugdb.*
 248. *Trachelium azureum umbelliferum.* I. R. H.
 RON. *Bald. p.* 44. In *Italia*.

SAMOLVS. LINN. 205.

Corolla exigua. Squamæ stamina muniētes.

H. *Samolus Valerandi.* LINN. Sp. p. 171. H.
Vps. 42. *Samolus Afr. folio rotundiore.* WALTH.
 H. 162. T. 23. In maritimis. A.

ISOSTEMONES CVCVRBITACEAE.

Omnes fere in *Monoccia*. Mas. Calyx quinque-
 partitus; petalum campanulatum quinquepartitum

ex calyce ortum. Stamina filamenta tria, antherae quinque flexuosae, *Famina*. Flos insidet fructui ornatissimo, calyx & petalum, ut in *ura*; stylus simplex stigmate trifido. In omnibus caulis scandens.

MOMORDICA. LINN. 967. *Elatium*.

H. *Momordica* pomis angulatis tuberculatis, foliis glabris patepti-palmatis. *Balsamina*. LINN. *Sp.* 1. *Mom. vulgaris*. T. R. H. *Balsamina* *forundifolia* *repens*. B. Pin. 386. Conf. charact. floris in *Comm. Soc. Gart. T. V.*

H. *Momordica* pomis angulatis tuberculatis, foliis villosis longitudinaliter palmatis. *Charantia*. LINN. *Sp.* 2. *Agave* *MORCAN* *Hesp.* 244. *Balsamina cucumerina indica*, fructu maturo flavescens. COMM. H. 1. p. 103. T. 54.

H. *Momordica* pomis oblongis sulcatis, catenularis, foliis incisis. *Luffa*. LINN. *Sp.* 3. *Cucumis Aegyptius reticulatus* f. *Luffa Arabum*. VESLING. *Aegypt.* 199. T. 58. 59. MORIS. I. p. 35. S. 7. T. 7. f. 1. 2. In *Ceylona*.

H. *Momordica* pomis hispitis, cirrhis nullis. LINN. *Sp.* 6. *Elatium* *Offic.* *Cucumis sylvestris* *aspinus*. B. Pin. 314. In *Europa* australi.

CUCURBITA. LINN. 968.

Cucurbita foliis subangulatis tomentosis, basi subtus biglandulosis, pomis lignosis. LINN. *Sp.* 1. *Cucurbita lagenaria*, flore albo, MORIS. *Hist.* 2. S. 1. T. 5. f. 3.

H. Cu-

H. *Cucurbita* foliis lobatis, pomis laevibus, seminum margine tumido. *Pepo*. LINN. *Sp.* 2. *Cucurbita maior Pepo oblongus*. B. *Pin.* 311.

H. *Cucurbita* foliis lobatis, pomis nodoso-verrucosis. LINN. *Sp.* 3. *Melopepo verrucosus*. I. B. H. 106.

H. *Cucurbita* foliis lobatis, caule erecto, pomis depresso-nodosis. LINN. *Sp.* 4. *Cucurbita clypeiformis*. I. B. II. p. 224.

CUCUMIS. LINN. 969. Melo.

H. *Cucumis* foliis multifidis, pomis globosis glabris. LINN. *Sp.* 1. *Colocynthis* B. *Pin.* 313. Lpb. ic. 644.

H. *Cucumis* foliorum angulis rectis, pomis oblongis scabris. LINN. *Sp.* 6. *Cucumis sativus vulgaris*. B. *Pin.* 310.

H. *Cucumis* foliis palmato-finnatis, pomis subovatis echinatis. LINN. *Sp.* 2. *Cucumis sylv. Amer. anguriae folio, fructu ovi figura, spinosis tuberculis momordicae instar muricato*. PLUK. *Phyt.* 170. f. 3.

H. *Cucumis* foliorum angulis rotundatis, pomis torulosis. LINN. *Sp.* 4. *Melo vulgaris*. B. *Pin.* 310. BLACKW. *T.* 329.

H. β. *Melo minimus*, pomis non maior, sphaericus. HALL. *Gott.* 208. An *Cucumis hirsutus pomis subglobosis*. LINN. *Sp.* 5. *Melo variegatus, aurantis figura, odoratissimus*. DILL. *Elsh. T.* 177. f. 218.?

ex calyce oritur. Stamina filamento breviter, antherae quinque flexuosae. *Famina*. Flos insidet fructui ornatissimo. calyx & patulum, ut in *ura*. Stylus simplex stigmate trifido. In omnibus caulibus scandens.

MOMORDICA. LINN. 967. Elaterium.

H. Momordica pomis angulatis tuberculatis, foliis glabris patentibus palmatis. *Balsamina. LINN. Sp. 1.* Mom. vulgaris. T. R. H. Balsamina rotundifolia repens. B. Pin. 366. Conf. charact. floris in *Comm. Soc. Carr. T. V.*

H. Momordica pomis angulatis tuberculatis, foliis villosis longitudinaliter palmatis. *Charantia. LINN. Sp. 2.* *Charantia* NOBILIS. *INsp. 244.* Balsamina cucumerina indica fructu maiore flavescens. *COMM. H. 1. p. 103. T. 54.*

H. Momordica pomis oblongis sulcatis catenulatis, foliis incisis. *Luffa. LINN. Sp. 3.* Cucumis Aegyptius reticulatus f. *Luffa Arabum. VESLING. Aegypt. 199. T. 58. 59. MORIS. 1. p. 35. S. 1. T. 7. f. 1. 2. In Ceylona.*

H. Momordica pomis hispida, cirrhis nullis. *LINN. Sp. 6. Elaterium offic.* Cucumis sylvestris asinus. B. Pin. 314. In Europa australi.

CUCURBITA. LINN. 968.

Cucurbita foliis subangulatis tomentosis, basi subtus biglandulosis, pomis lignosis. *LINN. Sp. 1.* Cucurbita lagenaria, flore albo, *MORIS. Hist. 2. S. 1. T. 5. f. 3.*

H. Cu-

CUCURBITACEAE.

247

H. *Cucurbita* foliis lobatis, pomis laevibus, seminum margine tumido. *Pepo*. LINN. *Sp.* 2. *Cucurbita maior Pepo oblongus*. B. *Pin.* 311.

H. *Cucurbita* foliis lobatis, pomis nodoso-verrucosis. LINN. *Sp.* 3. *Melopepo verrucosus*. I. B. H. 106.

H. *Cucurbita* foliis lobatis, caule erecto, pomis depresso-nodosis. LINN. *Sp.* 4. *Cucurbita clypeiformis*. I. B. II. p. 224.

CUCUMIS. LINN. 969. Melo.

H. *Cucumis* foliis multifidis, pomis globosis glabris. LINN. *Sp.* 1. *Colocynthis* B. *Pin.* 313. L. P. B. ic. 644.

H. *Cucumis* foliorum angulis rectis, pomis oblongis scabris. LINN. *Sp.* 6. *Cucumis sativus vulgaris*. B. *Pin.* 310.

H. *Cucumis* foliis palmato-5-angulatis, pomis subovatis echinatis. LINN. *Sp.* 2. *Cucumis sylv. Amer. anguriae folio, fructu ovi figura, spinosis tuberculis momordicae instar muricato*. PLUK. *Phyt.* 170. f. 3.

H. *Cucumis* foliorum angulis rotundatis, pomis torulosis. LINN. *Sp.* 4. *Melo vulgaris*. B. *Pin.* 310. BLACKW. *T.* 329.

H. β. *Melo minimus*, pomo non maior, sphaericus. HALL. *Gott.* 268. An *Cucumis hirsutus pomis subglobosis*. LINN. *Sp.* 5. *Melo variegatus, aurantii figura, odoratissimus*. DILL. *Elth. T.* 177. f. 218.?

BRYONIA. LINN. 970. Melothria.

LINN. 48.

Bacca ovalis, laevis. Sexus varius.

1. Bryonia foliis palmatis, vtrinque calloso-scabris. LINN. Sp. 1. Bryonia foliis quinquangulis scabris. HALL. H. 506. Br. alba, baccis nigris. B. Pin. 297. BLACKW. T. 37. Flores in dioica. Ad sepes prope portam Wehdunam.

H. Bryonia foliis palmatis quinquepartitis vtrinque laevibus, laciniis pinnatifidis. LINN. Sp. 5. Bryonia Afr. laciniata, tuberosa radice, floribus herbaceis. HERM. Parad. 107. T. 108.

H. Bryonia foliis palmatis scabris, laciniis lanceolatis serratis, lateralibus minimis. LINN. Sp. 4. Fl. Zeyl. n. 355. Bryonia Zeylonica fructu rubro, striis albis variegato. Pica est in HERM. Lugdb. T. 95. Hort. Malab. T. VIII. T. 19.

H. Bryonia floribus androgynis. HALL. G. 108. Melothria. LINN. Sp. p. 35. Omnes flores androgyni. In America.

SYCIOS. LINN. 971.

Bacca spinis obsita, unilocularis.

H. Sycios foliis angulatis. LINN. Sp. 1. Bryonioides flore & fructu minore. DILL. Elth. T. 51. f. 59. In Canada.

PASSIFLORA. LINN. 910. Granadilla.

Calycis folia & perala quinque. Triplex corona setarum circa stylum, qui stamina quinque & tres stylos profert.

H. *Passiflora* foliis palmatis integerrimis, involu-
cris cordatis tryphyllis. *coerulea*. LINN. *Sp.* 22. H.
Cliff. 432. HALL. *G.* 210.

H. *Passiflora* foliis crenatis tripartito divisis.
HALL. *l. c.* An *Passiflora foliis trilobis serratis*.
(*incarnata*.) LINN. *Sp.* 21.

H. *Passiflora* foliis trilobis cordatis aequalibus
obtusis glabris integerrimis. *lutea*. LINN. *Sp.* 15.
Passiflora folio tricuspidate, flore parvo flavescente.
I. R. H. HALL. *l. c.*

H. *Passiflora* foliis trilobis cordatis pilosis, invo-
lucris multifido-capillaribus. *foetida*. LINN. *Sp.* 19.
HALL. *l. c.* *Clematis Indica birsuta, foetida*. PLUM.
Amer. 71. T. 86.

ISOSTEMONES ROTACEAE.

Campanaceis valde similes calyce, staminibus & tu-
bis. Flos in plurimis infundibuliformis. Fructus po-
lypermus intra florem.

GENTIANA. HALL. 473. LINN. 285. *Swerdia*
LINN. 284.

Corolla varia. Fructus oblongus, bivalvis, disty-
lus. Folia alterna, simplicia.

Gentiana corollis brevissime quadrifidis, imber-
bibus verticillatis, sessilibus. LINN. *Sp.* 20. HALL.
Sp. 3. *Gentiana Cruciata*. I. R. H. T. 40. C. B. *Pin.*
188. *Im Göttinger-Wald*. In pratis prope *Herper-*
hausen.

1. *Gentiana corollis quadrifidis, corollis ciliatis, ciliata.* LINN. *Sp.* 19. HALL. *H.* n. 1. *Gent. coerulea oris pilosis.* B. *Pin.* t. c. *Gent. coerulea, sumbricata, autumnalis, angustifolia.* COLUMN. *Esqbr.* *Tab.* 221. In sylvis. v. g. *Göttinger Wald.*

2. *Gentiana foliis amplexicaulibus, corollis quinquefidis hypocrateriformibus, fauce barbatis.* LINN. *Sp.* 17. HALL. *H.* 2. *Gent. autumnalis ramosa.* B. *Pin.* 188. In sylvis *M. Heymberg.*

3. *H. Gentiana uniflora, corolla quinquefida campanulata, caulem excedente.* LINN. *Sp.* 8. HALL. *H.* 9. *Gent. alp. magno flore.* C. B. *Gentianella Helvetica,* LOBEL. *ic.* 310. In montibus *Helvetiae.*

4. *Gentiana foliis linearibus, floribus in alis sessilibus. Pneumonanthe.* LINN. *Sp.* 5. HALL. *H.* 10. LOBEL. *ic.* 309. *Gent. angustifolia autumnalis.* 1, R. H. In paludosis ducatus *Cellensis.*

5. *Gentiana foliis ternerviis, ovatis, floribus umbellatis, tubo strictissimo.* LINN. *Sp.* 14. HALL. *H.* 4. *Centaureum minus.* B. *Pin.* 178. BLACKW. *T.* 452. In sylvis.

6. *Centaureum minus palustre, ramosissimum.* VAILL. *T.* 6. f. 1. In paludosis.

7. *Gentiana caule brachiato, floribus flavis quadrifidis.* HALL. *Gott.* 201. *filiformis.* LINN. *Sp.* 2. *Centaureum palustre, luteum, minimum.* VAILL. *T.* 6. f. 3. Circa *Engesen* versus *Cellam.*

LYSIMACHIA. LINN. 188.

Flos rotatus. Capsula unilocularis decemvalvis.

Lyfi-

Lyfimachia foliis ovato-lanceolatis, racemis terminalibus. vulgaris. LINN. *Sp.* 1. HALL. *H.* n. 1. BLACKW. *T.* 278. *Lyfim. lutea maior, quae Dioscoridis.* B. *Pin.* 245. I. R. H. In paludibus circa *Diemen & Landolfshausen.* Ora segmentorum calycis miniata.

H. *Lyfimachia petiolis ciliatis, foliis cernuis. ciliata.* LINN. *Sp.* 7. *Lyfimachia Canadensis, folio Joleppae.* WALTH. *Hort.* 32, *T.* 12. In *Virginia.*

H. *Lyfimachia racemis simplicibus terminalibus, petalis obtusis, stipminibus corolla brevioribus. Ephemcrum.* LINN. *Sp.* 2. *Lyfim. spicata purpurea minor.* BUXB. *Cent.* 1. p. 22. *T.* 33. *Ephemcrum Matthioli.* B. *Pin.* 244. DOD. *Pempt.* 203. In *Oriente.*

H. *Lyfimachia foliis subquaternis, pedunculis verticillatis unifloris. punctata.* LINN. *Sp.* 6. *Lyfim. lutea minor, foliis nigris punctis notatis.* B. *Pin.* 245. *Lyfim. lutea* II. CLYS. *Hist.* II. p. 52. f. 2. In *Marchia.*

H. *Lyfimachia calycibus corollam superantibus, caule erecto ramosissimo.* LINN. *Sp.* 8. H. *Vps.* 37. *Lyfim. minimum stellatum.* B. *Pin.* 214. *Prodr.* 107. In *Gallia.*

Lyfimachia foliis ovato-acutis, floribus solitariis, caule procumbente. nemorum. LINN. *Sp.* 9. HALL. *H.* 3. *Anagallis lutea nemorum.* B. *Pin.* 252. In sylvis circa *Mosadon & Hordegfen.*

Lyfimachia foliis subcordatis, floribus solitariis, caule repente. LINN. *Sp.* 10. HALL. *H.* 2. *Nummularia s. centimorbia.* I. B. III. p. 370.

ANA.

ANAGALLIS. LINN. 189.

Priori similis. Capsula horizontaliter dehiscens.

1 Anagallis foliis indivisis, flore phoeniceo. HALL. H. 1. LINN. Sp. 1. BAVH. Pin. 252. Anag. terrestris mas. BLACKW. T. 43.

2 Anagallis flore coeruleo. HALL. H. 2. LINN. A. c. terrestris femina. T. 274. Vtraque super Papiermühle.

3 Anagallis foliis cordatis amplexicaulibus, caulibus compressis. LINN. Sp. 3. præter magnitudinem omnium partium mihi quidem diversa non videtur.

TRIENTALIS. LINN. 419.

Omnium partium numerus septenarius.

Trientalis. LINN. Sp. 1. Fl. Lapp. 139. Alfinanthemum. HALL. Iter. Hercyn. §. 22. In rupibus prope Waken & fylvis densioribus. Planta septentrionalis.

GLAVX. LINN. 257.

Calyx nullus. Petalum campanulatum.

1 Glaux maritima. LINN. Sp. p. 207. B. Pin. 215. LOBEL. ic. p. 415. In prato sale turgido prope Harste.

MENYANTHES. LINN. 185.

Segmenta floris fimbriata.

1 Menyanthes foliis ternatis. LINN. Sp. 3. HALL. Helv. 487. Trifolium palustre. BLACKW. T. 476. fibrinum. TABERN. 521. & offic. In paludibus. v. g. circa Diemern, Seeburg, Harste.

HOT-

HOTTONIA. LINN. 186.

Corollae infundibuliformis tubo stamina imposita.

Hottonia palustris. LINN. *Sp.* 145. HALL. *H.* 487. *Millefolium aquaticum*. B. *Pin.* 141. *Stratiotes vulgaris*. VAILL. *Act. Par.* 1719. p. 27. In vdis *Cellenfis*bus.

CYCLAMEN. LINN. 184.

Segmenta floris sursum flexa. Stamina vt in priori.

H. *Cyclamen corolla retroflexa*. LINN. *Sp.* 1. HALL. *H. n. 1.* *Artanita cyclamen* BLACKW. *T.* 147. LEB. *sc.* 604.

PHLOX. LINN. 197. *Lychnidea* DILL. Elth. 166.

Corolla infundibuliformis. Stamina inaequalia.

H. *Phlox foliis lato-lanceolatis*, inferioribus alternis, caule ramoso. *disarticata*. LINN. *Sp.* 5. *Lychnidea Virginiana*, *alsinet aquaticae*, *foliis*. PLUK. *Mant.* 121. In *Virginia*.

PRIMULA. LINN. 180.

Flores plures involucri excepti. Calyx quinqueangularis.

Primula foliis rugosis, hirsutis, scapo unifloro. HALL. *H.* 1. *Primula veris*, BLACKW. *T.* 52. singulari flore. I. R. H.

β. calyce colorato in segmenta petaloidea fissus.

Pri-

1. *Primula foliis rugosis hirsutis, spica inclinata, flore pallido inodoro.* HALL. *Hely. n. 2.* *Primula veris pallido flore elatior.* CLVS. *Hist. I. p. 391.*
1. R. H. LINN. *Sp. I. 3.* In pratis & sylvis.

1. *Primula foliis rugosis, hirsutis, spica inclinata, flore flavo odoro.* HALL. *H. I.* *Primula odorata, flore luteo simplici.* 1. R. H. Paralytis. BLACKW. *T. 226.* In sylvis. Omnino a priori diversa.

Primula foliis serratis carnosis glabris. LINN. *Sp. 5.* HALL. *H. 5.* *Auricula v. f.* CLVS. *Hist. 302.*

3. Varietates hortenses variorum colorum.

ANDROSACE. LINN. 179.

Flores ex involucrio. Petalum tubo ventricoso.

H. *Androsace perianthis maximis.* LINN. *Sp. I.* *Androsace vulgaris latifolia.* 1. R. H. *Andros. atropa.* CLVS. *Hist. II. p. 134.*

Androsace foliis lanceolatis dentatis glabris, perianthis angulatis corolla brevioribus. LINN. *Sp. 2.* *Andros. Coronopi folio glabra, flosculis ex verticilli foliosi umbilico plurimis albis.* AMMANN. *Rust. n. 22.* In Sibiria.

COFFEA. LINN. 209.

Corolla infundibuliformis. Stigmata duo. In bacca semina duo.

H. *Coffea.* LINN. *Sp. 192.* BOISSIEV. *Alt. Par. 1743. p. 388. T. 7.* TREW. *Commer. Nor. 1745. heb. 52.* BLACKW. *T. 337.* BOIS ALPIN. *Agropy. T. 36.* In Arabia.

LYCIVM. LINN. 232.

Priori similis, sed bacca bilocularis, polysperma.

H. *Lycium foliis cuneiformibus. Europaeum.* LINN. *Sp.* 3. *Jasminoides aculeatum, salicis folio, flore parvo ex albo purpurascente.* MICHEL. *gen.* 224. *T.* 105. *f.* 1. In Europa australi.

H. *Lycium foliis linearibus. Afrum.* LINN. *Sp.* 1. *Jasminoides aculeatum humile, balsmi minoris folio, flore violaceo.* MICHEL. *gen.* 224. *T.* 105. *f.* 2. H In Africa.

ISOSTEMONES ASPERIFOLIAE.

Calyx quinquefidus. Corolla limbo quinquefido. Stamina plerumque in tubo floris latentia, minima; tubum saepe claudunt pall aut glandulae aut plicae. In fundo tubi semina quatuor nuda, in quorum medio exurgit tuba unica simplex. Folia alterna, aspera aut hirsuta.

ECHIVM. LINN. 175. HALL. H. 513.

Corolla campaniformis limbo inaequali.

Echium caule simplici erecto, foliis caulinis lanceolatis, hirsutis, floribus spicatis lateralibus. *vulgare.* LINN. *Sp.* 3. HALL. *H. n.* 1. BLACKW. *P.* 199. CLVS. CLXIII. Vbique ad vias.

H. Echium calycibus frutescentibus distantibus, caule procumbente. *Creticum.* LINN. *Sp.* 4. Ech. *Cret. rubrum.* B. *Pin.* 254. CLVS. *Hist.* 2. CLXV.

CERINTHE. LINN. 171. HALL. H. 515.

Corolla regularis rubulato ventricosa, fauce nuda pervia. Capsulae feminales binae, dispermae.

H. Cerinthe foliis amplexicaulis, densibus floribus revolutis brevissimis. HALL. H. 1. *maior*. LINN. Sp.

1. CLVS. CLXVII. α. flore rubro. purpurascente.

H. β. flavo flore asperior. B. Pin. 258. In *Helvetia, Italia, &c.* A.

H. Cerinthe corollis fere semiquinquefidis acutis, clausis. HALL. H. 2. *minor*. LINN. Sp. 3. CLVS. CLXVIII. B. Pin. 268. Glasum montanum. DALECH. Prope *Jenam*.

SYMPHYTUM. LINN. 170. HALL. H. 514.

Corolla priori similis, sed faux palis clausa.

1 Symphytum foliis ovato-lanceolatis, decurrentibus, LINN. Sp. 1. HALL. H. 1. Symph. *Consolida maior*. BLACKW. T. 252. flore pallido, femina. V. bique in pratis.

1 β. flore purpureo *mas*. In *Helvetia*.

PULMONARIA. LINN. 169.

Calyx pentagonus. Limbus floris erecto-patens. In fauce ungues breves pilosi.

1 Pulmonaria foliis radicalibus ovato-cordatis scabris. *officinalis*. LINN. Sp. 2. HALL. H. 1. *maculosa*. BLACKW. T. 276.

1 β. *non maculosa*. In sylvis.

LITHOSPERMUM. LINN. 166. HALL.

H. 517.

Calycis segmenta subulata. Faux plicis munita.
Semina nitentia.

H. Lithospermum seminibus laevibus, floribus in
alis sessilibus parvis, foliis nervosis. HALL. H. 2.
LINN. Sp. 1. *maius erectum*. B. Pin. 158. 1. R. H.
Milium folis. 1. B. In *Helvetia*.

Lithospermum seminibus rugosis, corollis vix ca-
lycem superantibus. LINN. Sp. 2. HALL. H. 3.
BLACKW. 436. *arvense radice rubra*. C. B. Ubique
ad vias.

Lithospermum corollis calycem multisoties super-
antibus, caulibus procumbentibus radicatis. LINN.
Sp. 4. HALL. H. 1. Lithosp. *repens maius*. CLVS.
CLXIII. In praecipitio montis super *Luticlengen*.
In sylva prope *Grossen-Schneen*. P.

HELIOTROPIUM. LINN. 164. HALL.

Limbus planus. Tubus plicis concoloribus
clausus.

H. Heliotropium foliis ovatis acutis, spicis soli-
tariis, fructibus bifidis. LINN. Sp. 1. *americanum*
coeruleum, *hormini folio*. PLYK. Phyt. 245. f. 4.
HERM. Lugdb. 307. In *India*. P.

H. Heliotropium foliis ovatis integerrimis rugo-
sis, spicis coniugatis. LINN. Sp. 2. HALL. H. 520.
Hel. *maius Dioscoridis*. B. Pin. 253. CLVS. XLVI.
In *Europa australi*. A.

H. *Heliotropium*. foliis lanceolatis linearibus glabris, spicis coniugatis. LINN. *Sp.* 4. *americanum procumbens, facie lini umbilicati*. HERM. *Parad.* T. 143. In *America*.

MYOSOTIS. LINN. 165. *Scorpiurus*

HALL. 519.

Limbus plantis. Faux clausa glandulis coloratis.

Myosotis feminibus nudis, foliis hirsutis, radice annua, flore minori. LINN. *Sp.* 1. α . *Scorpiurus*. HALL. 1. *Echium scorpioides arvense*. C. B. flore coeruleo in humidis, albo in macilentis.

Myosotis foliis subhirsutis, radice perenni, flore maiori. LINN. *Sp.* 1. β . *Scorpiurus*. HALL. 1. *Echium scorpioides palustre*. C. B. in palustribus. β . Varietas hirsutior in ficcis.

H. Myosotis feminibus aculeatis glochidibus, foliis ovato-oblongis, ramis divaricatis. *Virginiana*. LINN. *Sp.* 2. Semina angulata. Fructus nutans. In *Virginia*.

Myosotis feminibus aculeatis glochidibus, foliis lanceolatis pilosis. *Lappula*. LINN. *Sp.* 3. *Cynoglossum foliis lingulatis, floris minimi tubo longiore*. HALL. H. 1. *Cynogl. minus*. B. Pin. 257. *Cynoglossa montana, minor, altera serotina*. COLVMN. *Ecpbr.* p. 180. Circa *Pless*. In muris.

CYNOGLOSSVM. LINN. 168.

Pali villosi tubum claudunt. Semina compressa, cabra.

Cyno-

Cynoglossum flaminibus corolla brevioribus, foliis lato-lanceolatis tomentosis sessilibus. officinale. LINN. Sp. 1. HALL. H. 2. BLACKW. T. 249. Cyn. vulg. B. Pin. 257. Ubique ad vias.

Cynoglossum flaminibus corollam aequantibus, foliis asperis. LINN. Sp. 4. Cynogl. mont. flore rubra, folio virente. I. R. H. Cyn. mont. maximum frigidarum regionum. COLYMN. Ecphr. I. T. 1704. In rupibus prope Herzberg; super Hefeld.

H. *Cynoglossum corollis calyce duplo longioribus, foliis lanceolatis sericeis.* LINN. Sp. 3. Cyn. Creticum argenteo angusto folio. B. Pin. 257. Cyn. Creticum I. CLVS. CLXII. Corollae albae, varis sanguineis. In Creta.

H. *Cynoglossum foliis lineari-lanceolatis glabris limifolium.* LINN. Sp. 5. Cynogl. minus album, limifolium foliis glaucis, semine umbilicato. MOR. III, S. XI. T. 30. f. 11. Omphalodes Lusitana elatior, cynoglossi folio. I. R. H. 140. In Lusitania.

H. *Cynoglossum repens, foliis radicalibus cordatis.* LINN. Sp. 6. Omphalodes verna symphyti folio. Hort. G. 223. Symphytum repens, Borriginis facie. B. Pin. 209. In Lusitania.

ANCHUSA. LINN. 167. Buglossum.

HALL. & TOURN.

Calyx teres tubo cylindrico floris aequalis.

Anchusa foliis lanceolatis, spicis imbricatis secundis. officinalis. LINN. Sp. 1. Buglossum floribus longe spicatis. HALL. H. 1. Bugl. angustifolium minus.

*ius. B. Pin. BLACKW. T. 500. Circa Haem-
veram.*

LYCOPSIS. LINN. 174. HALL. H. 523.

Tubus floris incurvus ventricosus. Semina sub-
longa.

† Lycopsis foliis lingulatis, sinuatis, hispidis, ca-
lycibus florescentibus erectis. *arvensis*. LINN. Sp. 1.
HALL. H. 523. Echioides RIV. T. 7. Buglossum
syto. minus. C. B. In agris inter segetes. P.

H. Lycopsis foliis integerrimis, calycibus fructe-
scentibus inflatis pendulis. LINN. Sp. 1. Lyc. *flore*
pullo calycibus amplis. HALL. G. 224. Echioides
flore pullo. RIV. T. XI. In *Thuringia*. A.

H. Lycopsis foliis repandis dentatis callosis, cau-
le decumbente, corollis cernuis. *variegata*. LINN.
Sp. 2. Lyc. *foliis hispidis verrucosis, flore variega-
to*. HALL. G. l. c. Buglossoides *Cretica*. RIV. T. 9.
In *Creta*. A.

ASPERVGO. LINN. 173.

Calyx quindecimfidus, laciniis quinque maiori-
bus, figura bivalvi semina comprehendit.

Asperugo. LINN. p. 138. HALL. H. 522. Bu-
glossum *sylvestre, caulibus procumbentibus*. B. Pin.
C. B. In vallo urbis ad portam *Weendanam*. A.

BORRAGO. LINN. 172.

Calyx & flos sine tubo stellatus, expansus.

H. Borrago foliis omnibus alternis, calycibus pa-
tentibus, *officinalis*. LINN. Sp. 1. HALL. H. 524.
Borra-

Borrago. BLACKW. T. 36. *floribus coeruleis* & β . *albis*. Sponte in hortis. Praeter palos elastra, stamina vrgentia. A.

H. Borrago radice perenni, flore reflexo. Differt a vulgari defectu vnguinum & longiori tubo floris.

H. Borrago foliis ramificationum oppositis, calycinis foliis sagittatis. LINN. Sp. 2. Cynoglossoides folio caulem amplexante. Ait. Par. 1725. f. 10. A.

H. Borrago calycibus tubo corollae brevioribus, foliis cordatis. LINN. Spec. 4. Borrago Constantinopolitana, flore reflexo coeruleo, calyce vesicario. TOURN. It. p. 523. Circa Constantinopolin. P.

ISOSTEMONES SOLANACEAE.

Calyx quinquefidus. Corolla limbo quinquepartito, regulari aut irregulari. Stylus simplex. Fructus ut plurimum bilocularis fuscus aut succulentus, feminibus receptaculo communi affixis. Odor gravis. Vires anodynae, narcoticae.

SOLANVM. LINN. 224.

Calyx & corolla rotata. Antherae conniventes. Fr. bacca.

Solanum caule inermi herbaceo, foliis ovatis, dentato-angulatis, umbellis nutantibus. *nigrum*. LINN. Sp. 12. HALL. H. 2. *bortense*. BLACKW. T. 167. *officinarium*. B. Pin. 166. Ubique in cultis & hortis. A.

β. *Solanum Guineense*, fructu magno instar ca-
rasi. DILL. *Elth. T.* 274. f. 354.

γ. *Solanum* ramis teretibus, foliis & floribus
villosis, baccis luteis. LINN. *Sp.* 12. γ. DILL. *Elth.*
274. f. 353. a vulgari, constanti partium villositate
diversum videtur.

δ. *Solanum* caule inermi herbaceo, foliis pinna-
tis integerrimis. *tuberosum*. LINN. *Sp.* 8. *Solan. tu-
berosum esculentum*. B. *Prodr. T.* 89. *Papas Peru-
vianorum*. *Tartassela*. Gottingae copiosissime co-
luntur. A.

ε. *Solanum* caule inermi herbaceo, foliis pinna-
tis incis, racemis simplicibus. *Lycopersicum*. LINN.
Spec. 9. *Poma amoris fructu luteo*. CAMER. *Epit.*
821. In *America*. A.

ζ. *Solanum* caule inermi frutescente flexuoso,
foliis superioribus hastatis, racemis cymosis. LINN.
Sp. 5. *Solan. scandens lignosum, f. Dulcanara*. I. R.
H. BLACKW. *T.* 34. In paludosis ad sepes. P.

η. *Solanum* caule frutescente subinermi, foliis
cuneiformibus sinuato-repandis. LINN. *Sp.* 7. *Solan.
Bonariense arborescens, papas floribus*. DILL. *Elth.*
T. 274. f. 351. Folia & flores maximi. In agro
Bonariensi. Fr.

θ. *Solanum* caule inermi fruticoso, foliis lanceo-
latis repandis, umbellis sessilibus. *Pseudo-capsicum*.
LINN. *Sp.* 3. *Solanum fruticosum bacciferum, Ame-
ricanum Plinii*. I. R. H. Fr.

ι. *Solanum* caule inermi fruticoso, foliis ovatis
integerrimis, floribus ad basin ramulorum confertis.

LINN.

LINN. *Sp.* 2. *Solan. Afr. tignosum*, folio atroviridi angusto oblongo obtuso. BOERH. II. p. 68. In Guinea. Fr.

H. *Solanum* foliis ovato-lanceolatis, fructu calloso, striato. HALL. G. 213. c. descr. An *Solanum* LINN. *Sp.* 4?

H. *Solanum* caule infermi herbaceoq, foliis ovatis tomentosis integris, calycibus spinosis. LINN. *Sp.* 13. *Melongena fructu oblongo luteo*. Mala insana, DOD. *Pempt.* 458. β. fructu rubro longo.

H. *Solanum* caule aculeis recurvis, foliis sinuatis subtus tomentosis utrinque aculeatis, pedunculis (& calycibus) aculeatis. LINN. *Sp.* 19. *Fla. Zeyl.* 95. * *Solanum pomiferum frutescens, Africanum, spinosum, nigricans, borraginis flore, foliis profunde laciniatis*. HERM. *Lugdb.* 573. T. 575. MOR. III. S. XIII. T. I. f. 15. PLUK. T. 226. f. 5.

CAPSICVM. LINN. 225.

Calyx erectus. Flos rotatus plicatus. Bacca magna exsucca.

H. *Capsicum* caule herbaceo, *annuum*. LINN. *Sp.* 1. *Piper Indicum vulgarissimum*. B. *Pin.* 132. Varietates ex fructu plurimae.

α. *Capsicum siliquis longis propendentibus, purpureis*.

β. *Caps. fructu flavo pyramidalis, oblongo*. I. R. H.

γ. *Caps. fructu latiore & rotundiore*. I. R. H.

δ. *Caps. fructu parvo, rotundo, acerrimo*. I. R. H.

ε. Capf. *Americ. fructu luteo, minimo.*

η. Capf. *Americ. fructu parvo, conico, erecto.*

θ. Capf. *fructu cordiformi erecto, & flore, ac in reliquis, maiori.*

PHYSALIS. LINN. 233. Alkekengi. T.

HALL. 506.

Calyx inflatus, coloratus, clausus fructum effecum comprehendit. Stamina libera.

H. *Physalis caule fruticoso, foliis ovatis integerrimis; floribus confertis.* LINN. *Cliff.* 62. *somnifera.* LINN. *Sp.* 1. *Alkekengi fruticosum, floribus verticillatis.* HALL. *G. Solanum somniferum.* CLVS. *H. p.* 85. In *Syria. Creta.*

1) *Physalis foliis integerrimis cordatis geminis, floribus solitariis.* LINN. *Sp.* 5. *Mat. med.* 93. *Alkekengi. f. Halicacabum.* BLACKW. *T.* 161. *officinarum.* 1. R. H. *Solanum vesicarium.* C. B. *Pin.* 166. In *M. Pless.*

H. *Physalis ramosissima, ramis angulosis glabris, foliis dentato-ferratis, angulata.* LINN. *Sp.* 6. *Alkekengi Indicum glabrum chenopodii folio.* DILL. *Elch. T.* 12. *f.* 12. *Solanum vesicarium Indicum.* COMM. *H. L. p.* 70. *T.* 17. A.

H. *Physalis ramosissima, foliis villosis-viscosis, ramis pubescentibus, pedunculis nutantibus, pubescens.* LINN. *Sp.* 7. *Alkekengi Virginianum, fructu luteo.* FEVILL. *Peruv.* III. *p.* 5. *T.* 1. *Solanum vesicarium Virgin. procumbens annuum, folio lanuginoso.* MORIS. III. *p.* 527. *S.* 13. *T.* 3. *f.* 24. *Corollae maculae fuscae.* In *India.* A.

ATRO-

ATROPA. LINN. 222. Belladonna. T.

HALL. 506.

Calyx & corolla tubuloso-campaniformis.

Atropa caule herbaceo; foliis ovatis integris. LINN. *Sp.* 1. Belladonna. HALL. *H.* 507. Solanum *melano-cerasus*. B. *Pin.* 166. *lethale* CLVS. II, LXXXVI. In sylvis copiose. A.

H. *Atropa* foliis sinuato-angulatis, calycibus clausis acutangulis. *physalodes*. LINN. *Sp.* 2. cum descr. Alkekengi amplo flore violaceo. FEVILL. *Peruv.* 724. T. 16. A.

NICOTIANA. LINN. 220.

Corolla infundibuliformis calyce longior, capsula ovata.

H. *Nicotiana* foliis lanceolatis. *Tabacum*. LINN. *Sp.* 1. *Nicot. maior. latifolia*. B. *Pin.* 169. *Tabacum. Petum*. copiose colitur. A.

H. *Nicotiana* foliis ovatis. *rustica*. LINN. *Sp.* 1. *Nicotiana minor*. B. *Pin.* 170. In *America*. A.

H. *Nicotiana* foliis cordatis, floribus paniculatis, tubis clavatis. LINN. *Sp.* 3. *Nicotiana minor, folio cordiformi, tubo floris praelongo*. FEVILL. *Peruv.* I. p. 717. T. 10. HALL. *Gutt.* 219. In *Peru*.

VERBASCUM. LINN. 217.

HALL. *H.* 509.

Corolla rotata, limbo irregulari. In plerisque stamina inclinata & villosa. Fr. fuscus.

I. Verbascum foliis decurrentibus vtrinque tomentosis, floribus densissime spicatis sessilibus. HALL. H. 1. *Thapsus*. LINN. Sp. 1. Verbas. *mar. latifolium, luteum*. B. Pin. 239. Vbique ad vias. A.

β. Verbascum *flore maiore, stigmate bicolli*.

II. Verbascum foliis sessilibus vtrinque viridibus, non decurrentibus, fasciculis florum remotis. Verbas. *foliis viridibus annuum, floribus luteis*. HERM. H. Lugdb. 618. HALL. H. 2 16. simile *Verbasco mari*, foliis autem & spica rariori diversa. A.

I. Verbascum foliis serratis cuneiformibus oblongis inferne villosis, pedunculis ramosis. HALL. H. 2. *Lychnitis*. LINN. Sp. 2. Verbas. *flore albo parvo*. G. B. 240. Verb. *candidum femina*. Rarius in ruderalis. A.

I. Verbascum foliis cordato-oblongis petiolatis, pedunculis ramosis, staminum barba purpurascens. HALL. H. 3. LINN. Sp. 3. Verb. *nigrum, flore ex luteo purpurascens*. I. R. H. Ad viam versus *Papiermühle*. In quatuor speciebus modo dictis praeter reliquas partes calyx & fructus lana obducti sunt.

H. Verbascum foliis sessilibus longe acuminatis, serratis, floribus in raram spicam dispositis, solitariis, staminum barba purpurascens. Planta vix satis nota. Summitates plantae vndique pilis glanduliferis obsitae. Flores fere sessiles, rari, tribus bracteis succincti. Fructus subrotundus calyci aequalis. Folia hirsuta non tomentosa.

H. *Verbascum* foliis amplexicaulibus oblongis glabris, floribus solitariis longius pedunculatis. *Blattaria*, LINN. *Sp.* 4. *Blattaria lutea*, folio oblongo laciniato. G. B. 240. α. flore albo extus purpureo; β. flore luteo. Folia radicalia sinuata, superiora serrata. Summitates pilis glanduliferis obsitae. In Germania australi. A.

H. *Verbascum* foliis ovatis nudis crenatis, caule subnudo racemoso. *phoeniceum*. LINN. *Sp.* 7. *Verb. foliis glabris indivisis*. HALL. G. *Blattaria purpurea*. B. *Pin.* 241. In Hungaria, Bohemia. P.

H. *Verbascum* tetrastemon. HALL. G. 217. *Celsia foliis duplicato-pinnatis*. LINN. *Sp.* 621. *Verbascum orientale*, *sophiae folio*. TOURN. *Cor.* 8. In Oriente, Habitus totus *Verbasci*, sola stamina recedunt. A.

HYOSCYAMVS. LINN. 219.

Corolla limbo irregulari. Fructus operculo testus.

Hyoscyamus foliis amplexicaulibus. *niger*. LINN. *Sp.* 1. *Mat. med.* 86. HALL. H. 512. *Hyosc. vulgaris vel niger*. B. *Pin.* 169. Ubique ad vias & in ruderalis. A.

H. *Hyoscyamus* foliis crispis circumferratis petiolatis, floribus pedunculatis. HALL. G. 218. LINN. *Sp.* 3. *Hyosc. Cretic. luteus minor*. I. R. H. G. B. *Pin.* *Hyosc. aureus*. ALPIN. *Exot.* p. 98. In Creta. *Monspeli.* A.

H. *Hyoscyamus* foliis petiolatis oblongis ampliter dentatis, floribus pedunculatis. *Hyosc. minor albo similis*, *umbilico floris atropurpureo*. TOURN. *Cor.*

MANETT. *Virid.* 518. Habitus totus albi, nisi quod flores pedunculati sint, in fundo violacei. A.

H. Hyoscyamus foliis petiolatis, floribus sessilibus. LINN. *Sp.* 2. Hyosc. *albus vulgaris.* CLVS. CXVIII. In *Europa* australi. A.

H. Hyoscyamus foliis lanceolatis subdentatis, calycibus spinosis. *pufillus.* LINN. *Sp.* 4. HALL. G. 218. In *Persia.* A.

ISOSTEMONES DICARPAE.

S. CONTORTAE.

Calyx quinquepartitus. Corolla quinquefida. Segmenta floris oblique dextrorsum flexa. In plurimis circa stamina nectaria cava, antherae conniventes aut connatae, & stigmata duo corpusculo cavo tecta. Folliculi duo polyspermi, seminibus pappo involutis.

NERIVM. LINN. 262.

Corollae tubus terminatus corona lacera.

H. Nerium foliis lineari-lanceolatis ternis, *Oleander.* LINN. *Sp.* 1. Rhododendrum. DOD. *Pempt.* 851. In *Palaeestina.*

β. Nerium *Indicum, flore pleno.* HERM. H. *Lugdb.* p. 549. Fr.

VINCA. LINN. 261. Pervinca

HALL. H. 526.

Corolla hypocrateriformis. Stylus stigmate duplici, inferiori orbiculato, superiori capitato.

Vinca

Vinca caulibus procumbentibus, foliis lanceolato-ovatis. LINN. *Sp. I. Pervinca vulgaris angustifolia*. F. R. H. *Clematis Daphnoides minor*. C. B. Pin. In sylvis. v. g. super *Herperhausen*. Fr.

H. *Vinca* caulibus erectis, foliis ovatis. *maior*. LINN. *Spec. 2. Pervinca latifolia flore coeruleo*. GARD. T. 81. In *Gallia Narbonensi*. Fr.

PERIPLOCA. LINN. 267.

Corolla rotata. Nectarium monophyllum ligulas emittens. Stigmata tecta.

H. *Periploca* foliis oblongis, floribus interne hirsutis. LINN. *Sp. I. HALL. G. 215. c. descr. Apocynum II. angustifolium*. CLVS. *Hist. Lp. 123. In Syria*. P.

ASCLEPIAS. LINN. 270.

Flores umbellati. Nectaria cava cum corniculis circa machinam staminum operculatam.

H. *Asclepias* foliis ovalibus subtus tomentosis, caule simplicissimo, umbellis nutantibus. *Syriaca*. LINN. *Sp. 4. Apocynum minus Syriacum rectum*. CORNVT. *Canad. T. 93. Apocyn. Syriacum*. CLVS. *Hist. II. LXXXVII. P.*

H. *Asclepias* foliis lanceolatis, caule superne diviso, umbellis terminalibus congestis. *incarnata*. LINN. *Sp. 8. Apocynum minus rectum Canadense*. CORNVT. *Canad. 9. T. 93. In Canada*. P.

H. *Asclepias* foliis lanceolatis petiolatis glabris, caule simplici, umbellis erectis solitariis. *Curassavica*.

vinc. LINN. Sp. 9. *Apocynum radice fibrosa, petalis coccineis, corollulis croceis.* DILL. *Eleph.* T. 30. f. 33. *Apocynum Curassavicum, fibrosa radice, floribus aurantiis, chamaeneris foliis latioribus.* HERM. *Par.* T. 36. HALL. G. 226. P.

h. *Asclepias caule erecto fruticoso, foliis revolutis lanceolatis, umbellis nutantibus, conceptaculis inflatis pilosis, fruticosa.* LINN. Sp. 14. *Apocynum erectum Afr. villosa fructu, fultis folio.* HERM. *Par.* 23. T. 24. COMMELE. H. f. 25. In Syria. Fr.

h. *Asclepias foliis ovato-lanceolatis, glabris, floribus racemosis, albis.* HALL. H. 525. LINN. Sp. 15. *Asclepias alba flore B. Pin. 363. Vincetoxicum.* DODON. *Pempt.* 407. In collibus lapideis. v. g. M. *Plefs. Hengst.* P.

APOCYNVM. LINN. 269.

Nectaria simplicia. Antherae connuentes.

h. *Apocynum caule rectiusculo herbaceo, foliis ovatis utrinque glabris, cymis terminalibus.* LINN. *Spec.* 1. *Apocynum Canad. foliis androsæmi majoris.* BOCC. *Sic.* 35. T. 16. f. 3. MORIS. III. p. 609. S. 15. T. 3. f. 16. In Virginia. Fr.

STAPELIA. LINN. 271. HALL. G. 227.

Corolla rotata. Nectarium stellatum, lacerum. Machinam stamineam & stigmata stellula tegit.

h. *Stapelia caulis aculeatis, flore levi.* HALL. G. 227. c. descr. *Stapelia denticulis ramorum pro-*
ten-

terribus. (variegata.) LINN. Sp. 1. *Apocynum humile aizoides, filiquis erectis, africanum.* HERM. Lugdb. T. 53. Fr.

H. *Stapelia* caulibus quadrangulis, flore pubescente. HALL. C. 228. *Stapelia dentatella ramorum erectis. (birsuta.)* LINN. Sp. 2. *Asclepias Afr. aizoides, flore pulchra subriata.* COMMEL. var. T. 19. Fr.

ISOSTEMONES, FLORE DIFFORMI.

Petala quatuor aut quinque irregularia; Epimedio excepto, cui flos regularis. In omnibus nectaria. Stamina antheris conniventes. Fructus est siliqua modo longior, modo brevior.

MELANTHUS. LINN. 712.
Flos papilionaceo similis, in cuius fundo fovea nectarifera. Stamina quatuor.

H. *Melanthus* stipulis solitariis petiolo adnatis. *major.* LINN. Sp. 1. *Melanthus Africanus.* HERM. Lugdb. 414 T. 415. In *Aethiopia.* Fr.

H. *Melanthus* stipulis geminis distinctis. *minor.* LINN. Sp. 2. *Melanthus Africanus foetidus minor.* COMMEL. var. 4. T. 4. Fr.

HYPECOUM. LINN. 157.

Petala quatuor inaequalia trifida. Siliqua articulata.

H. *Hypecoum* filiquis erectis torulosis. LINN. Sp. 3. *Hypecoum tenuifolium, filiquis erectis terribus.*

bus. AMMANN. *Ruth.* 58. T. 9. HALL. *Gott.* 230.
In *Sibiria.* A.

VIOLA. LINN. 898.

Petala quinque, infimum calcare nectarifero. Siliqua ovata.

f. *Viola acaulis*, foliis cordatis, stolonibus reptantibus. *odorata.* LINN. *Sp.* 5. HALL. *H.* 1. *Viola martia.* BLACKW. T. 55. *Viola martia purpurea*, flore simplici odora. B. Pin. 119. T. R. H.

f. *Viola caulibus adscendentibus floriferis*, foliis oblongo cordatis, *canina.* LINN. *Sp.* 9. *Viola montana inodora sylvestris.* B. Pin. 199. In sylvis.

h. *Viola foliis reniformibus*, caulibus foliosis floriferis. HALL. *H.* 5. *Viola caule bifloro*, foliis reniformibus serratis. (*biflora.*) LINN. *Sp.* 16. *alp. rotundifolia minor lutea.* PLVKN. T. 233. f. 1. In rupibus *Helvetiae.*

h. *Viola caule triquetro*, foliis reniformi-cordatis, floribus caulinis apetalis. *mirabilis.* LINN. *Sp.* 15. *Viola montana latifolia*, flores e radice, semina in cacumine ferens. DILL. *Elth.* T. 303. f. 390. In *Marchia.*

h. *Viola stipulis crenatis*, foliis ovato-lanceolatis. HALL. G. 23 I. c. descr. *Viola palustris*, foliis *Trachelii glabris.* RVPP. p. 289.

f. *Viola caule erecto*, stipulis laciniatis, reliquis longe ellipticis. HALL. *H.* 10. *montana.* LINN. *Sp.* 10. *Viola martia arborescens purpurea.* B. Pin. 199. In sylvis *hercynicis.*

Viola

Viola annua, caule procumbente, stipulis lacinatis, flore calycem excedente, tricolore. HALL. H. 11. *Viola tricolor*. BLACKW. T. 46. *γ. trinitatis*, TABERN. 306. In arenosis. v. g. *Celleri* Ducatu, circa Noribergam. A.

Viola annua, caule procumbente, stipulis lacinatis, flore calyci aequali, bicolore. HALL. H. 12. *γ. bicolor arvensis*. I. R. H. *Viola caule tetra diffuso, foliis oblongis incisi, stipulis dentatis*. LINN. Sp. 11. utramque iungit, notas autem diversitatis proxime exponam. In agris & ficcis. A.

IMPATIENS. LINN. 899. HALL. 501.

Corolla hexapetala, petalum supremum fornicatum, infimum calcaratum. Capsula elastice diffiliens.

H. *Impatiens pedunculis confertis, unifloris*. LINN. Sp. 5. *Balsamina femina*. B. Pin. 306. In India. A.

Impatiens pedunculis solitariis multifloris. HALL. H. 505. LINN. Sp. 7. *Balsamina lutea* f. *noti me tangere*. B. Pin. I. R. H. In ipsa urbe bey der grusfen Müble. In sylvis. A.

EPIMEDIUM. LINN. 138.

Nectaria quatuor cyathiformia, totidem petalis incumbentia.

H. *Epimedium alpinum*. LINN. Sp. 117. Hort. Vps. 29. DOD. Pempt. 599. In alpinis Italiae. P.

MEIOSTEMONES.

His stamina duo, quotquot segmenta floris sint,
Corolla infundibuliformis. Fr. bacca intra florem.

NYCTANTHES. LINN. 16.

Corollae limbus octopartitus.

H. Nyctanthes foliis inferioribus cordatis obtusis,
superioribus ovatis acutis. LINN. Sp. 2. Sambac
Jasmin Arabum. ALP. *Aegypt.* p. 73. CLVS. *cut.*
post. III. T. 3. Fr.

JASMINVM. LINN. 17.

Corollae limbus quinquefidus.

H. Jasminum foliis oppositis pinnatis. *officinale*,
LINN. Sp. 1. *Jasm. vulgare, flore albo.* I. R. H.
B. Pin. Fr.

H. Jasminum foliis alternis ternatis simplicibus
que, ramis angulatis. *fruticans*. LINN. Sp. 3.
Jasm. luteum vulgo dictum bacciferum. B. Pin.
398. Fr.

H. Jasminum foliis alternis ternatis pinnatisque
ramis angulatis. *humile*. LINN. Sp. 4. *Jasm. humile*
luteum. B. Pin. 397. Fr.

H. Jasminum foliis alternis ternatis pinnatisque
ramis teretibus. *odoratissima*. LINN. Sp. 5. *Jasm.*
Ind. flavum odoratissimum. FERRAR.

SYRINGA. LINN. 22. Lilac. T.

Corolla limbus quinquefidus. Capsula bilocu-
laris.

Syrin-

MEIOSTEMONES

212

Syringa folia ovata, cordatis, vulgaris. LINN. Sp. 1. *Syringa coerulescens flore.* CLVS. Hist. p. 56, β. flore albo. Ad sepes quali spontanea. Fr.
H. *Syringa foliis lanceolatis integris dissectisque.* Persica. LINN. Sp. 2. *Agemlilac Persarum.* CORNVT. 189. In Asia minori. ATTZOL.

OLEA. LINN. 20.

Corollae limbus quadrifidus. In drupa ovata nuda.
H. *Olea foliis lanceolatis.* LINN. Sp. 1. *Olea.* CLVS. Hist. p. 56. BLACKW. T. 213. *Sylvestris, folio duro subtus incano.* B. Pin. 472. In Europa australi. Fr.

LIGVSTRUM. LINN. 18.

Corollae limbus quadrifidus. Bases bilobularis.
Ligustrum. LINN. Sp. 1. HALL. H. 527. BLACKW. T. 142. *germanicum.* B. Pin. 473. Ad sepes. Fr.
Variae meiostemones ob analogiam relatae sunt inter didynamias.

DIDYNAMIAE.

Omniis his communia sunt calyx monophyllus apice incisus, corolla monopetala irregularis bilabata. Stamina quatuor duo maiora, duo minora, rarius alterum par deficit. Stylus simplex. Fructus in priori classe capsularis, in altera femina quatuor nuda.

DIDYNAMIAE ANGIOSPERMAE

Calyx quadri- aut quinquepartitus. Corolla vario modo difformis & bilabiata. Capsula unilocularis aut bilocularis.

MONTIA. LINN. 96.

Corolla quinquefida, stamina tria.

Montia. LINN. Sp. p. 87. HALL. H. 608. Montia aquatica minor. MICHEL. gen. 18. T. 13. f. 2. Alsinoides annua verna. VAILL. Paris. T. 3. f. 4. In pratis vdis super Hardeggen.

β. Varietas minor MICHEL. T. 13. f. 2. Circa Herzberg.

LIMOSELLA, LINN. 694.

Limosella. LINN. Sp. 631. HALL. Helv. 609. LIND. Alf. 156. T. 5. Plantaginella HALL. Jm. 23. T. 6. f. 2. palustris. B. Pin. 190. In inundatis limosis. v.g. vor dem Wehnder Thor gleich zwischen den Garten; an dem Lichtenmeer; bey der alten Leinenbrücke. Corollae limbus subaequalis. Plantula per flagella repens cespitosa, foliis longe petiolatis ovalibus.

OROBANCHE. LINN. 697.

Corolla ringens. Habitus aphyllus.

Orobanche caule simplici pubescente, bracteis longioribus, staminibus inclusis. LINN. Sp. 2. HALL. H. 611. n. 3. Orob. I. CLVS. I. p. 270. RIV. Irrig. T. 88. In M. Altenstollberg.

SQVA-

SQUAMARIA. HALL. Skt. Lachrya.

Squamaria. HALL. J. v. RIV. Hreg. T. 88. Lathraea caule simplicissimo, corollis pendulis, labio inferioris trifido. LINN. Sp. 4. Orobanche radice dentata major. ENCH. Calyx & corolla quadrifida. Pl. aphylla radice squamata.

LANTANA. LINN. 683.

H. Lantana foliis oppositis, caule inermi ramo-
so, floribus capitato-umbellatis. Camara. LINN. Sp.
3. Viburnum Amer. verticis foliis, lamii odore,
floribus miniatis. COMMEL. H. I. p. 151. T. 78.
In America calidior. P.

VITEX. LINN. 708.

H. Vitex foliis digitatis, spicis verticillatis. Agnus
castus. LINN. Sp. 1. RIV. T. 117. Vitex foliis
angustioribus cannabis modo dispositis. B. Pin. 475.
In Europa australi. Fr.

VERONICA. LINN. 26.

Corolla rotata quadrifida, stamina duo.

H. Veronica spicis terminalibus, foliis quaternis
quinisve. LINN. Sp. 1. Veron. Virgin. procerior,
foliis ternis quaternis & quinis caulem amplexans.
lin., spicis florum candidissimis. PLUK. T. 70. f. 2. P.

H. Veronica foliis ternis, ovato-lanceolatis, ser-
ratis, floribus spicatis. HALL. G. 234. An Veron. fo-
liis ternis aequaliter serratis, spicis terminalibus.
LINN. Sp. 2. Veron. spicata recta, profunde ser-
rata. I. B. III. p. 284. In Europa australi. P.

H. *Veronica foliis tenuis, ramulis dentatis, folia rariori.* HALL. G. l. c. *Veron. spicis terminalibus, foliis tenuis, inaequaliter serratis, (maritima.)* LINN. Sp. 3. In maritimis Europae. P. 1000. 1100. 1200.

H. *Veronica spicis terminalibus, foliis oppositis crenatis obtusis, caule erecto compositis.* LINN. Sp. 5.

5. *Veronica spicata lanuginosa & mollis, floribus coeruleis.* AMMANN. Ruth. 30. In Peruvia, P.

Veronica caule simplici, nudo, spicato, foliis oppositis scabris, ovatis, dentatis. HALL. G. l. c. LINN. Sp. 6. *Veronica spicata minor.* B. Pin. 247. Ad arcem Regensberg.

Veronica spicis lateralibus, pedunculatis, foliis oppositis scabris ovatis, caule procumbente. HALL. H. 7. *officinalis.* LINN. Sp. 8. *Veron. mas supina & oblongissima.* H. Pin. 246. BLANK. P. 143. In sylvis P. VII. A. VII. 1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

Veronica foliis ovatis cordatis, superioribus angustioribus ellipticis, ex alis racemosa. HALL. H. 6. *Veron. supina, facie Teucrii pratensis.* I. R. H. Ad vias. v. g. verius *Boraginaceae*. P. 1000. 1100. 1200.

Veronica foliis cordatis, superioribus maioribus, ex alis racemosa. HALL. H. 5. *Veron. racemosa terminalibus, foliis ovatis, angustioribus, caule debili.* (Chamaedryas.) LINN. Sp. 4. *Veron. minor, foliis imis rotundioribus.* I. R. H. In pratis.

Veronica foliis cordatis crenatis, longe minoribus, ex alis racemosa. HALL. H. 4. *Veron. latifolia.* LINN. Sp. 19. *Veron. pra. minima maxima.* v. g. In 34. In Helvetia.

Veron.

Veronica caule debili, foliis ovatis rugosis crenatis petiolatis, racemis lateralibus paucifloris, calyce hirsuto. LINN. *Cent. I. nov. spec. n. 1.* Veron. *procumbens.* RIV. T. 92. HALL. *Gott.* 235. Ad rivum supra *Geismar* in sylva.

Veronica racemis lateralibus, foliis ovatis planis, caule repente. *Beccabunga.* LINN. *Sp.* 14. HALL. *H.* 3. *Anagallis aquatica maior, folio subrotundo.* C. B. BLACKW. T. 48. In fossis. P.

Veronica racemis lateralibus, foliis lanceolatis serratis, caule erecto. LINN. *Sp.* 15. HALL. *H.* 2. Veron. *aquat. maior, folio oblongo.* B. *Pin.* 252. In fossis. P.

Veronica racemis lateralibus, pedicellis pendulis, foliis linearibus integerrimis. LINN. *Sp.* 16. HALL. *H.* 1. *Anagallis aquatica angustifolia scutellata.* Ad lacum prope *Seeburg* & alibi.

Veronica foliis subrotundis, crenatis, hirsutis, pedunculis solitariis calyce brevioribus. HALL. *H.* 15. *arvensis.* LINN. *Sp.* 21. Veron. *stosculis cauliculis adhaerentibus.* B. *Pin.* 250. In pratis & hortis.

Veronica foliis glabris, ovatis, floribus rariis, spicatis, sessilibus. HALL. *H.* 14. Veron. *racemo terminati subspicato, foliis ovatis glabris, crenatis.* LINN. *Sp.* 13. Veron. *pratensis Scorpillifolia.* C. B. In gramineis.

Veronica floribus solitariis, foliis cordatis incis pedunculo brevioribus, agrestis. LINN. *Sp.* 20. HALL. *H.* 18. Veron. *chamaedryos folio.* RIV. *monop. irreg.* T. 96. In cultis & agris.

Veronica foliis cordatis quinquelobis, calycis segmentis sagittatis. HALL. *H.* 17. LINN. *Sp.* 22. *Veron. Hederae folio.* RIV. *T.* 96. In agris & hortis.

β. *Veronica Chia verna, Cymbalariae folio, flore albo.* HALL. *G.* 237. *BUXB. Cent. I. T.* 40.

Veronica foliis trifidis & quinquefidis. HALL. *H.* 18. *Veron. floribus solitariis, foliis digitato-partitis pedunculo brevioribus.* *triphyllor.* LINN. *Sp.* 23. *Ver. rutae folio.* RIV. *T.* 95. In agris.

CAPRARIA. LINN. 686.

H. Capraria biflora. LINN. *Sp.* 628. foliis alternis, corollis quinquefidis. CLIFF. 320. *Curassavica.* HERM. *Parad. T.* 110. *Gratiolae affinis frutescens Americana, foliis Agerati s. Veronicae erectae maioris.* COMMEL. *H. L. p.* 79. *T.* 40. In America meridionali, Fr.

DODARTIA. LINN. 698.

H. Dodartia foliis linearibus integerrimis glabris, orientalis. LINN. *Sp.* 1. *Dodartia orient. flore purpureascente.* TOURN. *itin. III. T.* 208. In Oriente.

MIMULUS. LINN. 701.

H. Mimulus. LINN. *Sp.* 634. *H. Vpf.* 176. *T.* 2. *Lythmachia gutericulata.* GRONOV. *Virg.* 69. *Digitalis perfoliata glabra, flore violaceo minore.* MORIS. *II. S.* 5. *T.* 8. *f.* 6. In Canada.

ANGIOSPERMAE.

231

BROWALLIA. LINN. 691.

H. Browallia. LINN. *Sp.* 631. *H. Cliff.* 318. *T.* 17. *H. Vps.* 179. In *America australi.*

RVELLIA. LINN. 702.

H. Ruellia foliis petiolatis, floribus verticillatis subsessilibus. LINN. *Sp.* 1. *Vps.* 178. Ruellia *strepens*, capitulis camosis. DILL. *Elth.* 300. *T.* 249. *f.* 321. HALL. *G.* 304. In *Virginia.*

H. Ruellia foliis petiolatis, pedunculis longis subdivisis nudis. LINN. *Sp.* 2. *Vps.* 179. * Ruellia *capulis teretibus.* DILL. *Elth.* 328. *T.* 248. *f.* 320. HALL. *G. l. c.* In *Barbados.*

BIGNONIA. LINN. 677.

H. Bignonia foliis simplicibus cordatis, caule erecto, floribus diandris. *Catalpa.* LINN. *Sp.* 1. Bignonia *flore sordide albo, intus maculis purpureis & luteis adperso.* &c. CATESB. *Cor.* 1. *T.* 49. In *Carolina.* Fr.

DIGITALIS. LINN. 676.

Digitalis calycinis foliolis ovatis acutis; corollis obtusis. LINN. *Sp.* 1. HALL. *H.* 1. BLACKW. *T.* 16. Digit. *purpurea.* DOD. *Pempt.* 168. LOBEL. *ic.* 572. In sylvis *bercynicis* & sylva *Solingen.* *β.* flore albo.

Digitalis foliolis calycinis linearibus acutis, flore obtuso venoso, maculoso. HALL. *H.* 2. Digit. *lutea magno flore.* TABERN. *p.* 567. C. B. *Pin.* 1. R. H. In sylva *bercynica* circa *Elbingerode* & *Herzberg.*

H. *Digitalis* foliis calycinis lanceolatis, corollis acutis, labio superiore bifido. LINN. *Sp.* 2. *Digitalis lutea minore flore.* MORIS. II. p. 479. S. 5. T. 8. f. 5. RIV. T. 104. *parva.* LOBEL. *ic.* p. 573. In *Gallia.* B.

H. *Digitalis* calycinis foliolis ovatis obtusis, corollae labio inferiore longitudine floris. LINN. *Sp.* 3. *Digit. angustifolia, flore ferrugineo.* B. *Pin.* 244. HALL. G. T. 309. B.

CHELONE. LINN. 666.

H. *Chelone* foliis lanceolatis serratis, summis oppositis. *glabra.* LINN. *Sp.* 1. *Chelone Acadienfis flore albo.* TOURN. *Art. Par.* 1706. In *Canada.* Stamina fecunda quatuor, quintam abortivum.

GRATIOLA. LINN. 27.

H. *Gratiola* floribus pedunculatis, foliis lanceolatis serratis. *officinalis.* LINN. *Sp.* 1. p. 17. *Gratiola.* DOD. *Pempt.* 361. 362. BLACKW. T. 411. HALL. H. 618. 619. In paludosis *Europae* australioris. Filamenta duo antherifera, tria abortiva.

ANTIRRHINUM. LINN. 668.

H. *Antirrhinum* foliis oppositis cordatis crenatis. *Asarina.* LINN. *Sp.* 1. *Asarina* L. *Hedera saxatilis.* LOB. *ic.* 601. P.

Antirrhinum foliis cordatis quinquelobis alternis. LINN. *Sp.* 2. HALL. H. 1. *Cymbalaria.* C. B. CAMER. *Epit.* 860. RIV. T. 85. In rupibus prope *Münden.*

Antirrhinum foliis hastatis imis coniugatis, superioribus alternis, caule flaccido. LINN. *Sp.* 3. HALL.

Elatine f. H. 3. Elatine folio acuminato. C. B. 253. Elatine f. Veronica femina. BLACKW. T. 170. In Sylva, Lich bey Lengler. A.

Antirrhinum foliis ovatis, imis coningatis, superioribus alternis, caule flaccido. LINN. Sp. 4.

Elatine f. H. 3. Elatine folio subrotundo. C. B. Brydus Papiermühle. A.

H. Antirrhinum foliis ternis ovatis. triphyllum. LINN. Sp. 6.

Linaria Valentina Hiss. clus. l. p. 920. Linaria coerulea apula. COLUMN. Ephem. p. 78 c. ic. Variat coeruleo & luteo flore. In Hispania. A.

H. Antirrhinum foliis quaternis lanceolatis, caule erecto ramoso, floribus pedunculatis. LINN. Sp. 7. Linaria Americana maxima, purpureo flore. HERM. Lugdb. p. 377. RIV. T. 81. In America.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

H. Antirrhinum foliis quaternis linearilanceolatis, nectariis subulatis recurvis. LINN. Sp. 9. Linaria purpurea major odorata. C. B. l. r. h. Linaria flore purpureo. RIV. T. 84. In Gallia.

3. *Linaria ardens*, foliis carnosis, sessilibus minimis flavis. C. B. h. c. HALL. G. 307.

H. Antirrhinum foliis quaternis subulatis, caulis erectis. *mollis*. LINN. Cent. 1. n. 44. *Linaria foliis confertis linearibus carnosis*. HALL. G. 306. *Linaria maritima foliis succulentis*. RYXB. Cent. 4. p. 23. T. 37. A.

H. Antirrhinum foliis quinis linearibus multicaulis. LINN. Sp. 15. *Linaria Sicula multicaulis, mollis folio*. BOCC. Sic. 38. T. 19. f. 1. HALL. G. h. c. In Sicilia. A.

H. Antirrhinum foliis lanceolatis alternis, caule fruticoso. *Dalmaticum*. LINN. Sp. 18. *Linaria maxima folio Lappi*. I. B. III. p. 498. *Linaria fruticosa de Ragusa, odorata*. ZANON. T. 106. HALL. G. 307.

Antirrhinum foliis lanceolato-linearibus confertis ascendentibus, spicis terminalibus sessilibus, floribus imbricatis. *Linaria*. LINN. Sp. 11. HALL. H. 2. 6. *Linaria lutea vulgaris*. B. Pin. 212. BLACKW. T. 115. In gramineis & arvis.

H. Antirrhinum foliis lineari-lanceolatis alternis, floribus racemosis, calycibus corollis longioribus, caule erecto. *Chalopense*. LINN. Sp. 23. *Linaria annua angustifolia, sessilibus albis longius caudatis*. TRIVMF. Obs. T. 87. f. 2. In Italia. A.

H. Antirrhinum foliis lanceolatis obtusis viscidis alternis, caule ramosissimo diffuso. LINN. Sp. 24. HALL. Helv. 7. Antirrhinum *arvense minus*. B. Pin. 312. *Linaria arvensis minima*. RIV. T. 84. In siccis. A.

H. An-

H. *Antirrhinum* corollis ecaudatis spicatis, calycis foliolis brevibus ovatis. *glabrum*. LINN. *Sp.* 233. HALL. *H.* 9. *Antirrhinum* DOD. *Pempt.* 182. 202? 404. RIV. *T.* 81. A.

Antirrhinum corollis ecaudatis, sparsis, sessilibus, calyce brevioribus. *Orontium*. LINN. *Sp.* 26. HALL. *H.* 8. p. 615. *Gent.* 308. *Antirrhinum sylvestre*, *Polygama*; DOD. *Pempt.* p. 182. In agris.

ACANTHVS. LINN. 711.

H. *Acanthus* foliis sinuatis inermibus. LINN. *Sp.* 1. p. 939. *Mat. med.* 318. *Acanthus mollis Virginii*. B. *Pin.* 383. RIV. *T.* 86.

JUSTICIA. LINN. 26. *Adhatoda*. T.

H. *Justicia* arborea, foliis lanceolato-ovatis, bracteis ovatis persistentibus, corollarum galea contraria. *Adhatoda*. LINN. *Sp.* 1. *Adhatoda Zeylanensis*. HERM. *Lagdb.* *T.* 643. *Ecbolium*. RIV. *T.* 87. Fr.

H. *Justicia* fruticosa, foliis lanceolatis integerrimis, pedunculis trifloris encipitibus, bracteis calyce brevioribus. *hyssopifolia*. LINN. *Sp.* 4. In insula fortunatis. Fr.

SCROPHULARIA. LINN. 674.

HALL. 618.

Scrophularia caule obtusangulo, foliis cordatis, racemo terminali, calyce tubo floris aequali, galea emarginata. *Scrophularia nodosa foetida*. B. *Pin.* 235. LINN. *Sp.* 2. HALL. *H.* 2. RIV. *T.* 106. CAMER. *Epit.* 866. In fossis. P.

Stro-

Scrophularia caule membranis angulato, foliis cordatis, tubo floris ventricoso, e calyce longe eminentes. *Scrophularia aquatica maior*. CL. & BLACKW. T. 86. LINN. Sp. 3. HALL. H. L. De diversitate utriusque speciei plura dixi in *Comment. Gott. Tom. V. P.*

n. *Scrophularia* foliis cordato-oblongis, basi appendiculatis, racemis terminalibus, auriculata. LINN. Spec. 4. *Betonica aquatica septentrionalium*. LÖB. ic. 133. P.

n. *Scrophularia* foliis cordatis duplicato-serratis, racemo composito. LINN. Sp. 5. *Scrophularia Sverodoniae folio*. MORIS. S. V. T. 35. PLVKN. T. 59 f. 5. *Scroph. melissae folio*. L. H. 166. In *Lusitania*. P.

n. *Scrophularia* foliis cordatis superioribus alternis, pedunculis axillaribus bifloris. *peregrina*. LINN. Sp. 10. *Scrophul. peregrina*. CAMER. Hort. 157. T. 43. An *Scroph. folio rubro*. L. B. III. p. 422. A.

n. *Scrophularia* foliis cordatis acuminatis, floribus globosis, squamis sub galea dentatis. HALL. H. p. 619. HALL. Gott. 311. *vernalis*. LINN. Sp. 7. *Scroph. flore luteo*. B. Prodr. T. 112. RIV. T. 106. *Lamium Pannonicum* II. CLVS. Hist. II. XXXVIII. In *Italia, Helvetia*. P.

n. *Scrophularia* foliis difformibus, pedunculis axillaribus aggregatis. LINN. Sp. 8. *Scrophularia Sambuci foliis, flore rubro luteo vario pulchro*. PLVKN. T. 313. f. 6. MONTING. T. 238. HALL.

G. 311. *Sideritis Sambuci folio*. P. ALB. *Exot.* p. 202. Flores in hoc genere maximæ. In *Hispania*. P.

H. *Scrophularia* foliis pinnatis, calycis segmentis ovatis, flore ventricolo, squamula transversim latiori. *Scrophularia Cretica* L. f. *foliis filicis modo laciniatis*. CLVS. CCIX. An *Scroph. glauca folio*, in *amplas lacinias diviso*. TOURN. *Voy. Levant.* p. 84. Euse hanc speciem descripsi, & a recentioribus neglectam restitui in *Comment. Goss. Tom. V.* Tuba recta, stigmate hamato, recurvo.

H. *Scrophularia* foliis pinnatis, calycis segmentis rotundatis ala alba cinctis, tubo floris recto, squamula lineari. *Scrophularia Ruta canina dicta*. B. Pin. 236. HALL. *H. n. 4.* LINN. *Sp.* 9. *Ruta canina*. CLVS. CCIX. Tuba propendens reflexa, stigmate obliquo. In *Helvetia*.

PEDICULARIS. HALL. 620. LINN. 664.

Pedicularis caule ramoso erecto, calycibus bifidis crenatis, corollis labio obliquis. HALL. *H.* 621. *patens*. LINN. *Sp.* 1. *Pedicularis pratensis rubra etatior*. RAIL. *Hist.* 700. RIV. *T.* 91. Ad lacum prope *Seeburg*.

Pedicularis caule ramoso procumbente, calycibus quinquefidis crenatis, floribus solitariis labio cordato. HALL. *H.* 622. *sylvatica*. LINN. *Sp.* 1. *pratensis purpurea*. B. Pin. 163. In sylvis paludosis.

ALECTO-

ALECTOROLOPHVS. HALL. H. 624.

Rhinaanthus LINN. 658.

Alectorolophus corollarum labio superiore compresso, calycibus glabris. HALL. H. 624. 1. Rhinaanthus. LINN. Sp. 3. Pedicularis *pratensis lutea* f. *crista Galli*. B. Pin. 163. In pratis.

Alectorolophus corollarum labio superiore compresso, calycibus hirsutis. HALL. H. 624. 2. Pedicularis lutea. TABERN. 791.

MELAMPYRVUM. LINN. 660.

Melampyrum spica densa longiori, foliis inter flores laciniatis, adscendentibus. HALL. H. 2. Melamp. *spicis conicis laxis laceris. (arvense.)* LINN. Sp. 2. Melamp. *purpurascente coma*. B. Pin. 234. In segetibus.

Melampyrum spica densa brevi, foliis inter flores cristatis, reflexis. HALL. H. 625. 1. Melamp. *spicis quadrangulatis compactis, obtusis, imbricatis. (aristatum.)* LINN. Sp. 1. RIV. T. 80. *luteum angustifolium*. C. B. In M. Altenstollberg.

Melampyrum spica laxissima, foliis inter flores laciniatis, patentibus. HALL. H. 3. Melamp. *bracteis cordato lanceolatis, summis coloratis floribus, calycibus lanatis. (nemorosum.)* LINN. Sp. 3. Melamp. *coma coerulea*. C. B. Ad oras nemorum.

Melampyrum floribus sparsis sessilibus, binatis, foliis superioribus incis. HALL. H. 4. Melamp. *florum coniugationibus remotis, crotolis clausis. (pratense.)* LINN. Sp. 4. Mel. *luteum latifolium*. C. B. In sylvis circa *grossen Schween & Hardeggen*.

Melam-

GYMNOSPERMAE.

222

Melampyrum, floribus sparsis, sessilibus, binatis, hiantibus, foliis omnibus indivisis. HALL. *H. 5.*
Melamp. corollis hiantibus sylvaticum. In sylvis *Hercynicis.*

ODONTITES. HALL. 627. Euphrasia.

LINN. Sp.

Odontites foliis inter flores ferratis. HALL. *H. 1.*
Euphrasia foliis linearibus, omnibus ferratis. LINN.
Sp. 4. Euphr. pratensis rubra. C. B. In pratis & agris humentibus.

EVPHRASIA. HALL. 628. LINN.

659. Sp.

Euphrasia foliis ovatis, acutè dentatis. HALL. *ff.*
n. 1. officinalis. LINN. *Sp. 2. officinarum.* C. B.
 In pratis.

β. flore minori. In sylvis.

DIDYNAMIAE GYMNOSPERMAE.

S. VERTICILLATAE.

Caulis quadrangularis. Folia opposita. Flores in verticillos dispositi. Calyx monophyllus, quinquefidus. Corolla irregularis, maximum bilabiatam ringens, labio superiori bifido, inferiori tripartita. Semine quatuor duo maiora, duo minora. Stylus simplex, stigmate bifido. In fundo calycis semina quatuor nuda. Omnibus vires aromaticae, roboretantes.

T

I. Flore

I. Flore galea destituta.

BVGVL. HALL. 633. *Aiuga*.
LINN. 624.

Labium inferius tripartitum.

Bugula foliis ovatis, inter flores stipatis, flagellis reptans. HALL. H. p. 634. *Aiuga stolonibus reptatricibus*. LINN. Sp. 4. *Bugula*. DOD. Pempt. 135. In sylvis.

Bugula foliis angulose dentatis, caule simplici. HALL. H. 635. *Aiuga tetragono-pyramidalis*. LINN. Sp. 3. *Bugula montana*. RIV. T. 75. In sylvis M. Heymberg & alibi.

II. *Bugula* (*Aiuga*) floribus resupinatis. LINN. Sp. 1. *Bugula Orient. villosa, flore intersecoerulea, alba macula notata*. DILL. Elth. T. 43. f. 61. HALL. Gott. 316.

TEVCRIVM. LINN. 625. *Chamaedrys*.
HALL. 630.

Labium inferius quinquepartitum.

Teucrium foliis multifidis, floribus verticillatis utrinque ternis. LINN. Sp. 3. *Chamaedrys foliis pinnatis, pinnis obtusis, trifidis*. HALL. H. 633. *Iva moschata, folio multifido*. RIV. T. 14. In M. arcis Pless.

II. *Teucrium* foliis trifidis linearibus integerrimis, floribus sessilibus axillaribus. *Chamaepitys*. LINN. Sp. 5. *Bugula foliis trifidis, floribus per alas sparsis*. HALL. H. *Chamaepitys lutea vulgaris*. si Pin. 249. Sponte in Thuringia. A.

II. *Teu-*

H. Teucrium foliis integerrimis oblance-ovatis petiolatis, supra villosis, subius tomentosis incanis.

Teucrium latifolium. LINN. *Sp.* 10. *H. Vfs.* 159.

G. I. *Teucrium fruticans* *baccicum ampliare folia*. DILL. *Elth.* 197. *T.* 284. f. 367. Folia superne ex atroviridia, lana telae araneae instar obducta, sit ne obtusiori; inferne cum ramis tomento niyeo tecta. Fr.

H. *Teucrium* foliis integerrimis ovatis utrinque acutis, recentis secundis villosis. *Marum*. LINN. *Sp.* 12. *Marum Syriacum*. BLACKW. *T.* 47. Fr.

H. *Teucrium* foliis ovatis acutis subdenticulatis, floribus solitariis. *Marum Hispan. nigrum* *Piperella dicta*. BOCC. *Mass.* II. *T.* 117. BARREL. *var.* *T.* 694. Prioris similis, sed folia differunt, & odor ille penetrans & tortis desideratur. Periole foliis breviores.

Teucrium foliis cordatis serratis petiolatis, racemis lateralibus secundis, caule erecto. LINN. *Sp.* 17. *Scorodonia*. RIV. *T.* 12. In sylvis prope *Münden*; in nemore *Solingen*. P.

H. *Teucrium* foliis cordatis crenatis petiolatis, spicis longissimis, densissimis. HALL. *Comm. Gott.* 1752. cum *ic. H. Gott.* 315. In *Hyrcania*. R.

H. *Teucrium* foliis cordatis undulatis obtuse serratis, floralibus integerrimis concavis, caule fruticoso. *flavum*. LINN. *Sp.* 20. *Teucrium vulgare fruticans* L. CLVS. *Hist.* I. p. 348. *Teucrium* c. 2. RIV. *T.* 10. In *Europa* australi.

2. *Teucrium* foliis oblongis sessilibus dentato-serratis, foliis geminis axillaribus, caule diffuso. LINN. *Sp.* 8. *Chamaedrys foliis ellipticis, serratis, multibus, verticillis inter folia paucifloris.* HALL. *H. p.* 631. *Scordium* C. B. BLACKW. *T.* 475. In patudosis. P.

H. *Teucrium* foliis ovatis inciso-crenatis petiolatis, floribus subverticillatis. LINN. *Sp.* 19. *Chamaedrys procumbens, foliis solidatis, ovatis, rotunde crenatis.* HALL. *H. p.* 630. *Chamaedrys repens.* C. B. RIV. *T.* 10. In Thuringia. P. METGM.

H. *Teucrium* corymbo terminali, foliis lanceolatis integerrimis. LINN. *Sp.* 21. *Chamaedrys foliis angustis integris, ellipticis subtus incanis.* HALL. *H. p.* 632. *Polium matris Cordi* Hist. 104. b. *Anga folio integra.* RIV. *T.* 15. In Thuringia. P.

Galea trifida.

SCUTELLARIA. LINN. 653. *Callida.*

Calyx cristatus. Tubus floris incurvus.

Scutellaria floribus cordato-lanceolatis crenatis, floribus axillaribus *galeatutata.* LINN. *Sp.* 4. *Callida.* HALL. *p.* 635. *Callida palustris vulgarior, flore coeruleo.* I. R. H. In aquosis & ad fossas

H. *Scutellaria* caule ramosissimo, foliis ovato-lanceolatis, racemis axillaribus multifloris. *Callida latifolia, flore minore.* RIV. *T.* 77. HALL. *G.* 317. *Racemi florum inter folia latent. Folia floralia reliquis similia, floribus maiora. Flosculi parvi et minimis huius generis coerulei.* An *Scutellaria palust.*

Just. repens: Virginiana maior, flore minore, MORIS.
III. p. 416. LINN. Sp. 3.?

H. *Scutellaria foliis subcordatis serratis, spicis interruptis, peregrina, LINN. Spec. 8. H. Vpf. 1784.*
M. L. *Ossida COLYMNAE Ephn. I. p. 187. T. 189.*
HALL. G. 1. t. *Lanum, astragaloides. CORNUT.*
Canad. T. 129. Bracteae calycibus minores, ovatae, obtuse serratae, glabrae.

H. *Scutellaria foliis cordatis, crenatis, floribus spicatis secundis, bracteis flore multo minoribus, Bracteae calycis longitudine, ovatae, integerrimae, pilosae. Calyx fuscus, insigniter cristatus. Flores omnes erecti, secundi, purpurei, labio inferiori albo. An. Cassida Orientalis, altissima, virens folio.*
I. R. H. *Coroll. HALL. G. 318. An Scutellaria foliis subovatis crenatis, florum racemis nudiusculis.*
LINN. Sp. 11.?

H. *Scutellaria foliis cordatis serratis, verticillis diffusis, bracteis superantibus. Caules diffusi ramosi. Bracteae ovatae, integerrimae, floribus duplo maiores. Calyx parvus, fuscus, pilosus. Floris exigui galea carnei coloris, hirsuta, barba albida.*

H. *Scutellaria foliis subovatis, serratis, villosis, spicis imbricatis, bracteis flores aequantibus. Bracteae inferiores foliis similes, superiores floris longitudine obovatae integerrimae pilosae. Calyx pilosus, insigniter cristatus. Corollae villosae ochroleucaae. An Scutellaria foliis cordatis obtusis, obtusisque serratis, spicis fastosis. LINN. Sp. 12. Cassida. Cres. fruticosa, folia Catarinae, flore albo.*
TOURN. Coroll.

H. *Scutellaria* foliis ovatis crenatis, floribus capitato-spicatis bracteis superantibus. Spicae breviter densae. Bracteae ovato-acutae ellipticae imbricatae; corollae unciales patulae, tubo ochroleuco, limbo violaceo. An *Scutellaria* LINN. Sp. 9. *Tedcrum alpinum* inodorum, magna flore. B. Prodr. 116:1.

H. *Scutellaria* (*Cassida*) spicis foliosis. GMELIN. HALL. Gott. 318. c. defet.

H. *Scutellaria* foliis incisis subtus tomentosis. *O. sternalis*. LINN. Sp. 1. *Cassida Orient.* flos *Cassidator*, flore luteo. TOURN. Itin. III. T. 306. COMMEL. rar. T. 30. XIV. T. 77.

III. *Galea erecta*, integra, aut emarginata.

SALVIA. LINN.

Calyx bilabiatus $\frac{1}{2}$. Filamenta duq brevissima, quibus in medio fere articulatione affixa sunt duo alia longa, quorum extremitati superiori anthera inferiori glandula aut callosa aut parum farinosa insidet. In omnibus rudimenta duorum staminum sterilium.

I. *Salviae*. TOURN. Calycis & corollae fauces hiantes. Galea recta. Filamenta antherifera fere semicircularia, inferius glandula pulverulenta aucta.

III. *Salvia* foliis lanceolato-ovatis integris crenatis, floribus secundis subdidynamis tubo calyce superante. *Salvia officinalis*. LINN. Sp. 4. *Salvia major. minor. a. non auriculata.* B. P. L. 111. T. 72. *a. auriculata*, fere und alterne foliis ad basin folii accessorio. C. B. I. B. III. p. 105. III. *Europea natural.*

c. n. *Salvia* foliis pinnatis, pinnis ovatis serratis, calycibus amplis inflatis, tubum floris aequantibus. *Salvia pinnata*. LINN. *Sp.* 26. *Salvia Orientalis latifolia bifurcissima, viscosa, pinnata, flore & calyce purpureis, odorata*. BOERH. *Lugdb.* II. T. 167. In Oriente. Fr.

II. *Sclareae*. Galea magna falcata. Filamenta filamentum falcata, glandula callosa.

H. *Salvia* foliis rugosis oblongo-cordatis serratis, bracteis concavis acuminatis coloratis, calyce longioribus. *Sclarea*. LINN. *Sp.* 33. H. *Vpf.* 10. n. 4. *Horminum Sclarea dictum*. C. B. *Pin.* 235. In Syria.

H. *Salvia* foliis cordato-sagittatis serratis acutis glutinosa. LINN. *Sp.* 19. H. *Vpf.* 11. HALL. H. 639. *Horminum luteum glutinosum*. B. *Pin.* 238. *Horminum sylvestre* II. GLYS. XXIX. In Helvetia.

H. *Salvia* foliis oblongis dentato-angularibus hirsutis, verticillis lanatis, corollarum labio faccato. LINN. *Sp.* 23. *Aethiopsis foliis sinuosis*. C. B. 241. A.

H. *Salvia* foliis rugosis pinnatifidis lanatis verrucosis. *cratophylla*. LINN. *Sp.* 24. *Horminum rugoso verrucosoque folio, cornu cervi exprimente*. PLUKN. *Phyt.* T. 194. f. 5. HERM. *Parad.* T. 186. HALL. *Gott.* 323. A.

H. *Salvia* foliis hastato-triangularibus oblongis crenatis obtusis. *Canariensis*. LINN. *Sp.* 20. *Horminum Canariense, hastato folio, caule tomentoso*. PLUKN. *Phyt.* T. 301. f. 2. MORIS. III. 394. S. XL. T. 13. f. 17. HALL. G. 321. In Canariis. Fr.

Salvia foliis cordato-oblongis crenatis, verticillis subnudis diffusis, calyce tubo floribus brevior. Salvia *pratensis*. LINN. Sp. 11. Sclarea *prat.* foliis serratis, flore coeruleo. I. R. H. *Horminum sylv.* AUC. CAMER. Epit. 629. Prope Rudorf. P.

H. Salvia foliis cordatis acute crenatis, verticillis subnudis remotissimis, calyce aristato. LINN. Sp. 17. Sclarea *Indica*, floribus variegatis. I. R. H. HALL. G. 320.

H. Salvia foliis ovatis utrinque acuminatis serratis. *Mexicana*. LINN. Sp. 15. Sclarea *Mexic.* altissima; facie *Heltotropis*. DYLL. Elib. T. 254. f. 136. HALL. G. 320. Jungia. HEISTERI Praefat. ad Epist. *Burkhardianam*. F. L. p. 92.

H. Salvia foliis cordatis inaequaliter basi eximie caule nudo, spica florente pendula. HALL. Comm. Gott. T. I. T. XI. LINN. Sp. 27. Var. flore albo. In *Russia*.

III. *Hormina*. Galea brevis, parum erecta. Stamina recta.

H. Salvia foliis ovatis, crenatis, bracteis cordato-acuminatis sensim decreescentibus, viridibus, in spicam tetragonam digestis. Salvia *foliis oblongis crenatis*, corollarum galea semiorbiculata, calycibus fructiferis reflexis. LINN. Sp. 7. H. Vps. 11, n. 8. *Horminum coma viridi*. I. R. H. Calyces fructiferi reflexi, labii superioris dentibus divergentibus. A.

H. Salvia foliis obtusis, crenatis, bracteis summis sterilibus maioribus coloratis. LINN. Sp. 8. *Horminum*

innatum *colorata*, *rubra*. HALL. G. 326. I. R. H.
Calyx priori similis, ovis autem reflectitur. A. 300

H. *Salvia* foliis cordato-sagittatis dentatis, verticillis multifloris, stylo labio inferiori incumbente.
Salvia verticillata. LINN. Sp. 18. H. Vpf. n. 7.
Horminum sylvestre latifolium verticillatum. C. B.
 HALL. Gorr. 322. In Austria, Helvetia. A.

H. *Salvia* foliis cordato-lanceolatis, duplicato-crenatis, undulatis, galea punctata, staminibus galea multo brevioribus. *Hormini sylvestris quinti altera species.* CLVS. XXXI. *Horminum sylvestre salvisolium minus vel maculatum.* C. B. *Sclarea folio salvide maior vel maculata.* I. R. H. BOERH. Lugd. II. 165. Hanc cum sequente iungit LINNAEVS Sp. 8. nomine *Salviae foliis cordato-lanceolatis, bracteis coloratis flore brevioribus.* P.

Salvia foliis cordato-lanceolatis, simpliciter ferratis, staminibus galeam aequantibus. *Hormini sylv. quinti species prior.* CLVS. l. c. *Horminum sylv. f. salvisolium minus.* C. B. RYPP. Fl. Jen. 221. MORISON. III. p. 394. *Sclarea folio Salviae minor f. glabra.* I. R. H. BOERH. l. c.

H. *Salvia* foliis ovato-ferratis, hirsutis galea obtusa, stylo galeae longitudine. *Salvia verbenaca.* LINN. Spec. 13. *Hormini sylv. quarti quinta species.* CLVS. XXXIII. *Horm. sylv. lavendulae flore.* C. B. 320. I. R. H. *Horm. sylv. intiso folio, parvo flore, italicum.* BARKER. 220. *Horm. Verbenae laciniis angustifolium.* TRIVMF. Obs. T. 66. HALL. G. 322.

H. *Horminum minus, folio sinuato, flore amaranthino coeruleo.* DIEL. HALL. G. l. c. A.

H. *Salvia foliis pinnato-dentatis, asperis, hirsutis, verticillis pilosis, floribus apetalis. Planta a priori omnino diversa, foliis exsuccis, hirtis, rugosioribus, terrae appressis, caule simplicissimo piloso, bracteis & calycibus undique hirsutissimis. In calyce iuniori quidem corolla minima emarcida reperitur, quae vix unquam autem adolescit, semina tamen fecunda sequuntur.* A.

H. *Salvia foliis sinuato-serratis glabris, calyce piloso viscido, corolla vix calycem superante. Horminum Virginianum Blattariae folio.* HALL. G. 224. *Calyx latus post florescentiam amplo rictu, hians violaceus. Corolla violacea, galea recta brevissima. An Salvia foliis radicalibus lyratis dentatis, corollarum galea brevissima. (lyrata.)* LINN. Sp. 3. *Salvia corollarum labio superiore brevior, fauce patente.* ERONOV. Virg. 82. A.

H. *Salvia foliis ovato-oblongis simpliciter serratis, calyce hirsuto longius aristato, corolla brevissima. An Horminum folio betonicae, flore coeruleo.* BOERH. l. p. 165. *Calyx angustior viridis, neque diffusus. Corolla coerulea, tenuior.*

H. *Salvia foliis ovatis serratis, spicis tetragonis imbricatis, calycibus tridentatis, corollarum galea brevissima. Salvia foliis ovatis, basibus mucronato-tuberosis, spicis imbricatis, calycibus trifidis. Hispanica.* LINN. Sp. 16, c. desc.

ROSMARINUS. LINN. 34.

H. Rosmarinus officinalis. LINN. Sp. p. 23. Mat. med. 16. Rosmarinus hort. angustiore folio. C. B. In Europa australi. Stamina duo, sed simplicia, neque bifurcata. H.

MONARDA. LINN. 34.

Galea linearis, longissima. Stamina duo.

H. Monarda capitulis terminalibus, caule obtusangulo. fistulosa. LINN. Sp. 1. Origanum fistulosum Canadense. CORNUT. Canad. T. 14. RIV. T. 61. In Canada. P.

H. Monarda floribus verticillatis, corollis punctatis. LINN. Sp. 4. Clinopodium Virgin. angustifolium; quodis verticillo duodecim foliis rubantibus cincto. PLVKN. Alm. III. T. 24. f. In Virginia. P.

ZIZIPHORA. LINN. 33.

H. Ziziphora capitulis terminalibus, foliis ovatis, capitata. LINN. Sp. 1. Clinopodium fistulosum pumilum Indiae occidentalis, summo caule floridum. PLVKN. Alm. T. 164. f. 4. In Syria. Huic generi calyx tubulosus, corollae limbus parvus.

H. Ziziphora floribus lateralibus, foliis lanceolatis, tenuior. LINN. Sp. 12. HALL. G. 324. Achnos Syriacum, folio mucronato, capsulis bifurcis. MORIS. III. p. 304. S. XI. T. 19. f. 3. 4.

OCLIVM. LINN. 61.

Calyx superne latus, planus. Flores inversi.

H. Ocl-

H. Ocimum foliis ovatis, glabris, calycibus ciliatis. *Basilicum*. LINN. Sp. 2. Ocimum *styracis* f. *melissae* odore. BOERH. Ocimum *caryophyllatum*, *maximum* & *maius*. C. B.

β. *latifolium maculatum vel crispum*. C. B.

γ. *viride foliis bullatis*. C. B. A.

H. Ocimum foliis ovatis integerrimis. *minimum*. LINN. Sp. 3. Ocimum *minimum*. C. B. MORIS. III. p. 407. S. XI. T. 10. f. 17. A.

BRYNELLA. LINN. 654.

H. Brunella foliis imis integris, superioribus laciniatis. HALL. H. 637. Brunella *foliis ovato-oblongis petiolatis*. LINN. Sp. 1. γ. *laciniata*. Brunella *fol. laciniato*. C. B. 261. RIV. T. 30. In *Helvetia*. P.

Brunella foliis dentatis, spica foliis minore. HALL. H. 636. Brunella LINN. Sp. 1. Var. α. Brunella *maior, foliis non dissectis*. C. B. In *prat.* P.

Brunella foliis vix dentatis, spica foliis maiore. HALL. H. 637. LINN. Sp. pr. Var. β. Brunella *flor. maiore*. RIV. T. 29. Circa *Gesfhereder Feld*. P.

LEONURVS. LINN. 641. Cardiac. TOURN.

Calyx pentagonus, rubrofusus. Gales *cancra* integra. Antherae punctatae.

Leonurus foliis caulinis lanceolatis trilobis. *Cardiac*. LINN. Sp. 1. HALL. H. p. 639. Cardiac 1. R. H. & aut. BLACKW. T. 171. In *ruderalis*.

β. Car-

B. Cardiaea *Oriens*, *caulescens*, folio *vanunculi* *pratensis*, flore *purpureo*, calyce *moli*. 1. R. H. *Hieracii* calycis & foliorum *differe* videtur.

H. *Leonurus* foliis *tripartitis* *laciniatis*, calycibus *villosis*. *Tartaricus*. LINN. *Sp.* 3. *Cardiaea* foliis *ternatis* & *profundius* *intesis*, *glabra*. AMMANN. *Ruth.* 49. 50.

H. *Leonurus* foliis *tripartitis* *multifidis* *linearibus* *obtusiusculis*. *Sibiricus*. LINN. *Sp.* 4. *Phlomis* foliis *multifidis*. HALL. *G.* 329. *Ballote* *inodora*, foliis *Coronopi*. AMM. *Ruth.* 48. T. 81. In *Siberia*.

H. *Leonurus* foliis *palmatis* *dentatis*, calycibus & *caule* *lanatis*. *Cardiaea* *Chinensis*, albis calycibus, *aculeatis*, *semine* *nigro*. *Comm. Soc. Sci. Gott. T. III.* *Ballote* foliis *palmatis* *dentatis*, *caule* *lanato*. LINN. *Sp.* 2.

• *Leonurus* (*Cardiaea*) foliis *cordiformibus*. HALL. *H.* 3. *Galeopsis*. LINN. *Sp.* 4. *Lamium* *oblongo folio*, flore *luteo*. G. B.

PHLOMIS. LINN. 642.

Involucrum *setaceum* *verticillis* *densis* *subiectum*. *Priori* *affinis*.

H. *Phlomis* foliis *subrotundis* *tomentosis* *crenatis*, *involucris* *lanceolatis*. *fruticosa*. LINN. *Sp.* 47. *Phlomis* *fruticosa*, *Salviae* *folio* *longiore*. 1. R. H. *Phlomis* *latifolia* *capitata* *lutea* *grandiflora*. DILL. *Elsh. T.* 237. f. 306. In *Hispania*. Fr.

H. *Phlomis* *involucris* *hispidis* *subulatis*, foliis *cordatis* *scabris*, *caule* *herbaceo*. *suberosa*. LINN. *Sp.*

Sp. 6. *Phlomis foliis condiformibus galba lactea*. HALL. *Comm. Gott. T. I. T. 12. Enum. Gott. 328. Phlomis foliis parvis subrotundibus, villosissimis, berrimi foliis*. AMMANN. *Roth. n. 28. P.*
 H. *Phlomis foliis lanceolatis serratis, involucris linearibus nudis, calycibus decagonis, decemdentatis*. LINN. *Sp. 12. Leonurus capitis bonae Spei*. BREYN. *Cent. T. 86. Leonurus Afr. frutescens angustifolia, flore longissimo phoeniceo, Leonurus dista*. MORIS. III. S. XI. T. 19. f. 17.

LAMIUM. LINN. 636.

Lamium foliis distitis cordatis acutis. HALL. H. n. 1. *Lamium foliis cordatis acuminatis serratis petiolatis*. LINN. *Sp. 1. Lamium album. β. purpureum non foetens, folio oblongo*. O. B. *Archangelica alba*. BLACKW. T. 33. Ad sepes. Huic generi calyx aristatus, fauces floris spinula terminatur.

Lamium foliis petiolatis, cordatis, obtusis, in summo caule congestis. HALL. H. 2. LINN. *Sp. 2. Lamium purpureum foetidum, folio subrotundo*. O. B. *Vrtica iners altera*. DOBON. *Pemps. 153. In hortis*.

Lamium foliis superioribus amplexicaulibus obtusis incisis. HALL. H. 2. LINN. *Sp. 3. Lamium minus, folio caulem ambiente*. O. B. 231. In hortis.

GALEOPSIS. LINN. 637.

Galeopsis foliis & caule fulcrato, hispidis, flore calyce duplo longiore. HALL. H. n. 1. *Galeopsis internodiis superius incrassatis, verticillis summis sub-*

subcontinguis. LINN. *Sp.* 2. var. *α*. *Cannabis spuria*. RIV. T. 31. In sylvis caeduis & ruderibus.

1. flore albo.

γ. flore magnovario colore. *Cannabis spuria, flore majore*. RIV. T. 32. In dumetis um den Landolfsbäuser Teich.

Galeopsis foliis ovato-lanceolatis, mollissimis & tomentosis. HALL. H. 3. *Galeopsis pr. spae, var. β*.

1. var. flore ochroleuco, barba violacea.

2. flore purpureo.

Galeopsis foliis longe ellipticis subhirsutis. HALL. H. 4. *Galeopsis internodiis caulinis aequalibus, verticillis omnibus remotis*. LINN. *Sp.* 1. *Ladanum segetum*. RIV. T. 24. In segetibus.

STACHYS. LINN. 638.

Stachys verticillis sexfloris, foliis cordatis petiolatis acuminatis, sylvatica. LINN. *Sp.* 1. HALL. H. 1. *Stachys sylvatica*. RIV. T. 26, *Galeopsis*. BLACKW. T. 84. In sylvis & ad sepes.

1. *Stachys verticillis sexfloris, foliis lineari-lanceolatis semiamplexicaulibus, palustris*. LINN. *Sp.* 3. *Stachys verticillis spicatis, foliis subhirsutis ellipticis, ad basin emarginatis*. HALL. H. 4. *Stachys palustris foetida*. C.B. In agris humectis.

H. *Stachys* foliis hirsutis spicam distinguendis, imis cordiformibus. HALL. H. 2. *Stachys verticillis multifloris, tomentosis, foliis lanceolato-ovatis basi emarginatis*. LINN. *Cliff.* 301. *alpina*. *Sp.* 3. *Stachys*

chrys folia obscure vivente, flore ferruginea. MOENCH.
II. p. 382. T. X, f. 11. In montosis Helvetiae.

H. *Stachys caule lanato foliis tomentosis, obis, oblongo-cordatis, verticillis multifloris, densissimis.*
HALL. H. 3. *germanica.* LINN. Sp. 4. *Stachys maior Germanica.* C. B. I. R. H. *montana.* RIV. T. 2.

BETONICA. LINN. 631.

Betonica foliis, inter verticillos in vertice congestis, brevissimis & integris. HALL. H. 3. *officinalis.* LINN. Sp. 1. *Betonica* BLACKW. T. 46. *vulgaris purpurea.* C. B. I. R. H. In sylvis. Ch. gen. *Galea plana erecta.*

Betonica foliis petiolatis, crenatis, subglabris, bracteis inter verticillos dissitis longioribus, ellipticis, serratis. HALL. H. 1. *Betonica verticillata, calycibus spinosis.* LINN. Sp. 4. *Var. α . Sideritis arvensis, latifolia, glabra.* C. B. Inter fegetes.

Betonica foliis sessilibus, crenatis, hirsutis, bracteis inter verticillos dissitis brevibus, aristatis, integris. HALL. H. 2. LINN. Sp. pr. var. β . *Sideritis vulgaris hirsuta.* C. B. *vulgaris* I. CLVS. H. XXXIX. *Virus quae diversitatem specificam fuisse dixi in com. Gott. T. V.*

SIDERITIS. LINN. 632. Cunila

HALL. 334.

Corolla subaequalis, calyce parum longior, vix singens, stamina intra tubum floris.

Sideritis (Stachys) foliis oblongis acutis, oppositis, serratis: floralibus cordatis acutis, interger-

Sideritis. ROY. *Lugdb.* 318. HALL. G. 327. *Sideritis perfoliata*. LINN. *Sp.* 4. *Sideritis orient. phlo-*
midis folio. 1. R. H. *Corollae albae venis rufis*.

H. *Sideritis fruticosa tomentosa, foliis cordatis,*
pedunculis ante florescentiam nutantibus. Canariensis. LINN. *Sp.* 1. *Stachys Canariensis frutescens;*
Verbasci folio. COMMEL. H. II. p. 197. T. 99.
HALL. G. 327. In *Canariis*. Fr.

H. *Sideritis fruticosa, tomentoso-lanata, foliis*
lineari-lanceolatis. Syriaca. LINN. *Sp.* 3. *Cunila*
foliis ovatis tomentosis. HALL. G. 335. *Sideritis*
Cretica tomentosa, candidissima, flore luteo. 1. R. H.
In *Creta*. Fr.

H. *Sideritis foliis lanceolatis subdentatis, bracteis*
cordatis dentato-spinosis, calycibus aequalibus,
birsuta. LINN. *Sp.* 8. *Sideritis foliis ellipticis, cre-*
natis, distinctis verticillis. HALL. *Helv.* p. 647.
Sideritis Monspeliaca scordioides. LOBEL. *Adv.* 424.
Vide characterem in *Obs. botan. epist. ad Ill. WERL-*
ROFIVM. In *Gallia Narbonensi*. P.

H. *Sideritis herbacea decumbens, calycibus spi-*
nosus, labio superiori trifido. montana. LINN. *Sp.* 5.
Cunila calycibus patulis aristatis. HALL. G. 335.
Sideritis montana, parvo flore nigro purpureo, ca-
pite medio croceo. COLVMN. *Ephr.* I. T. 196. In
Italia. A.

H. *Sideritis (Cunila) foliis ovatis dentatis, ver-*
ticillis multifloris, calycibus sessilibus spinosis.
HALL. G. 334. *Leonurus*. LINN. *Sp.* 2. *Marrubia-*
strum Cardiacae folio viridi. ROCC. *Mus.* II. T.
98. Circa *Wittbergam*. A.

Sideritis foliis cordatis oblongis crenatis, verticillis sexfloris, staminibus tubo florispaulo maioribus. *Cunila foliis ovatis, dentatis, hispidis*. HALL. G. 334. *Glechoma cretensis*. LINN. Sp. 2. *Stachys minima*. RIV. T. 27. In collibus inter segetes.

IV. Galea profunde fissa.

BALLOTE. LINN. 639.

Ballote foliis cordatis indivisis serratis. *nigra*. LINN. Sp. 1. Ballote HALL. H. 648. BLACKW. T. 136. Marrubiastrum. TABERN. 506. RIV. T. 65. Ad sepes.

MARRUBIUM. LINN. 640.

I. Calycibus quinquedentatis.

H. Marrubium foliis cuneiformibus, verticillis involucro destitutis. LINN. Sp. 1. Marrubium foliis cordatis, retusis, serratis. HALL. G. 336. Alysson Galeni. CLVS. II. XXXV. In Hispania.

H. Marrubium foliis ovato-lanceolatis serratis, calycum denticulis setaceis. *peregrinum*. LINN. Sp. 2. Marrub. foliis ellipticis serratis venosis, calycibus quinquedentatis. HALL. G. 336. Marrubium Creticum. TABERN. 539. In pago Erdeborn.

H. Marrubium foliis oblongis obtusis, mollibus orenatis, calycum denticulis subulatis. Marrubium Hisp. supinum; foliis sericeis, argenteis. DILL. Elsb. F. 215. nostram optime exprimit. A priori differt,

sele, foliis de caule lanatis, verticillis denso-
ribus. L. 6007. *Steno hyssopifolius*.

Marrubium dentibus calycinis fetaceis vinctatis.
LINN. *Sp. 1517*. Marrubium *album vulgare*. CIV. S. II.
XXXIV. Q. B. L. R. H. In ruderatis. v. g. ad arcem
Pless.

G. Marrubium *album villosum*. G. B. *Prodr.* 140.
Pin. 239. R. H. 192. egregia varietas caule de
inferna parte foliorum denso tomento obductis,
odore suavi aromatico.

H. Marrubium foliis cordatis, serratis, tomen-
tosis. HALL. G. 336. Marr. *calycum limbus paten-
tibus, densiculis acutis*. LINN. *Sp.* 7. Marr. *al-
bum rotundifolium Hispanicum*. HERM. *Par.* T. 201.
In Hispania. P.

H. Marrubium foliis ovatis tomentosis integer-
rimis. HALL. G. 4. c. Marrub. *calycum limbus pla-
nis villosis, foliis cordatis, caule fruticosa*. LINN.
Sp. 8. Pseudodictamnus. DON. *Pempt.* 281. In
Creta. Fr.

NEPETA. LINN. 629. Cataria HALL.
649. TOURN.

Nepeta floribus interrupte spicatis, verticillis
subpedicellatis, foliis petiolatis cordatis dentato
serratis. LINN. *Sp.* 1. Cataria 1201. H. 2. Herba
Cataria MATH. 719. Nepeta. L. *Mentha cataria*.
BLACKW. T. 455. In ruderatis.

H. Nepeta floribus paniculatis, foliis cordatis
petiolatis obsolete crenatis. *Pannonica*. LINN. *Sp.* 2.

000000

V 2

Cata-

Cataria angustifolia *minor*. 1. R. H. *Calamintha montana praealta*, *Pulegii* odore. BOCC. Mus. II. ic. 38. In *Pannonia*.

H. *Nepeta* (*Cataria*) foliis ovatis, serratis, hirsutis. HALL. G. 337. *Cataria peregrina latifolia*. LOBEL. 511.

H. *Nepeta* foliis petiolatis cordato-oblongis dentatis, verticillis pedunculatis distite spicatis. LINN. Sp. 3. HALL. H. 2. *Cataria Hispanica B. tonicae folio angustiori*, flore coeruleo. 1. R. H. In *Helvetia*. P.

H. *Nepeta* floribus spicatis sessilibus, bracteis ovatis coloratis verticillos excipientibus, foliis sessilibus. LINN. Sp. 8. *Nepeta rubro-rosea, spicata, purpurea, Hispanica*. BOCC. Mus. T. 36. In *Hispania*. P.

H. *Nepeta* caule acuto quadrangulo, spicis densissimis nudis. HALL. G. 338. *Hyssopus* LINN. Sp. 3. *Betonica maxima, Scrophulariae folio, floribus incarnatis*. HERM. Par. p. 106. c. 1c. Flores iuniores ochroleuci, aetate rubescunt. In *Virginia*.

H. *Nepeta* (*Cataria*) floribus inversis. HALL. G. 338. *Comment. Gott.* 1752. *Hyssopus corollis transversalibus*. LINN. Sp. 2. In *Sibiria & China*.

HYSSOPUS. LINN. 628. HALL.

H. *Hyssopus* spicis secundis, officinalis. LINN. Sp. 1. *Hyssopus officinarum coerulea f. spicata*. C. 1. 217. BLACKW. T. 296. β. flore albo. In rupibus *Satz des Helden* quasi sponte.

DRACO-

DRACOCEPHALVM. LINN. 648.

Moldavica HALL.

H. *Dracocephalum* floribus spicatis, foliis lanceolatis ferratis. LINN. *Sp.* 1. BREYN. *ic.* 33. *T.* 27. RIV. *T.* 25. *Pseudodigitalis Persicae foliis.* BOCC. *Sic.* 12. *T.* 6. *f.* 3. In *Virginia*.

H. *Dracocephalum* floribus spicatis, foliis compositis. LINN. *Sp.* 2. *Moldavica foliis ternatis.* HALL. *G.* 331. *Cedronella Canariensis, viscosa, foliis plerumque ex eodem pedunculo ternis.* COM-MEL. *Amst.* II. *T.* 41. *Dracocephalo affinis, Americana trifoliata, Terebinthi odore.* VOLKAM. *Norib.* *T.* 145. In *Canariis*.

H. *Dracocephalum* foliis ex lanceolato-linearibus rarius dentatis spinulosis. *Comm. Gott.* III. *p.* 436. *Dracoceph. floribus oppositis, bracteis lanceolatis integerrimis, foliis lanceolatis, mucronatis, dentatis.* LINN. *Cent.* II. *Sp.* nov. *n.* 160. In *Sibiria*.

H. *Dracocephalum* floribus spicatis, foliis bracteisque lanceolatis indivisis muticis. LINN. *Sp.* 5. *H. Vpf.* *p.* 156. *Russchiana glabra foliis integris.* AMMANN. *Ruth.* 50. In *Sibiria*. P.

H. *Dracocephalum* floribus verticillatis, bracteis lanceolatis, serraturis capillaceis. LINN. *Sp.* 7. *Moldavica foliis ellipticis, profunde crenatis.* HALL. *G.* 331. *Melissa Moldavica.* CAMER. *Epit.* 576. *Melissophyllum Turcicum.* LOB. *adv.* 220. *Melissa Turcica offic.* In *Moldavia*. A.

H. *Dracocephalum* floribus verticillatis, bracteis oblongis, serraturis spinosis, foliis subtomentosis
V 3 *canescens.*

canescens, LINN. Sp. 18. H. V. 166. 7. Moldavia orientalis betonicae folia, flore magno violaceo, COMMEL. rar. T. 28. TOURN. Cor. p. 11.

β. Eadem flore magno albedo. TOURN. f. p. 22. Sideritis annua flore luteo, utriculis & foliis longioribus. MORIS. III. p. 389. S. XL T. 8. f. 18. In Oriente.

γ. Dracocephalum floribus verticillatis, bracteis orbiculatis, serrato-ciliatis, peltatum, LINN. Sp. 9. H. V. 166. * Moldavia foliis ovatis, crenatis, floralibus orbiculatis creberrime ciliatis, peramplis. HALL. G. 332. Moldavia orient. salicis folia, flore coeruleo. I. R. H. Cor. A.

δ. Dracocephalum floribus verticillatis, bracteis oblongis, ovatis integerrimis, corollis manifestis axillaribus. LINN. Sp. 10. H. V. 167. 4. 6. Moldavia foliis glabris ovatis, floralibus lanceolatis, floribus calyce duplo longioribus. HALL. G. 333. Moldavia Betonicae folio, floribus minoribus coeruleis pendulis. AMM. Ruth. 44. In Sibiria.

ε. Dracocephalum floribus verticillatis, bracteis oblongis integerrimis, corollis ovatis calyce minoribus. LINN. Sp. 11. H. V. 167. 4. 7. Moldavia foliis ovatis, floribus calycem vix superantibus. HALL. G. 332. Moldavia Betonicae folio, floribus minimis pallide coeruleis. AMM. Ruth. 46.

ζ. Dracocephalum (Moldavia) foliis cordatis triangularibus, serratis corymbis geminis pedunculatis axillaribus. HALL. G. 333. Comment. Gort. 1752. Nepeta Sibirica. LINN. Sp. 11. Gortia mon-

montana foliis veronicae praetensis Gent. III.
p. 27. T. 50. f. 1. In Sibiria.

Andillensis PRASIVM LINN. 655.

H. Prasium foliis ovato-oblongis serratis. *maius*.
LINN. Sp. p. 601. Teucrium fruticosum, amplo
albo flore, italicum. BARREL. ic. 895. Lamium
arborescens perenne Ericicum. ZANON. Hist. 1. 12. T.
26. In Italia. II.

MOLVCELLA LINN.

H. Molucella calycibus quinquedentatis, denti-
culis aequalibus. LINN. Sp. p. 1. Molucca Vae-
voda. DOD. Pappr. 99. Molucca Moluccana odorata.
DOD. Pl. Syn. 2. 8.

H. Molucella calycibus septemdentatis. *spinosa*.
LINN. Sp. 2. Melissa Moluccana spinosa. DOD. l. c.
HALL. G. 338. A. H. M. L. C. 1070.

MELITIS. LINN. 650.

H. Melissa thalictophyllum. LINN. Sp. 597. Me-
lissa. HALL. H. 630. Melissa Pichsi. CAMER.
Hort. p. 99. n. 30. MELISSA. LINN. Sp. 515. Lamium
Pannonicum I. CLVS. XXXVII. In theonotis Germa-
niae. Helvetiae. R.

HORMINYM LINN. 649.

H. Horminum foliis cuneiformi-oblongis, caule
bifido. *virgatum*. LINN. Sp. 2. Melissa atropu-
rens, bugulae folios. DILL. Ehb. T. 175. f. 216.
in Virginia. Nobis semper depectulata, sedina ca-
men petist. Aeneas in quoque eluso.

libit

MELISSA. LINN. 647. *Calamintha*.
HALL.

H. Melissa floribus ex alis inferioribus subsessilibus.
LINN. *Vps.* 163. *Mat. med.* 294. *officinalis*. LINN.
Sp. 1. Melissa BLACKW. *T.* 27. *Bortensis*. B. *Pin.*
229. P.

H. Melissa foliis ovatis obtusis, caule procum-
bente. HALL. *H. n.* 2. *Calamintha pulegii odore*.
B. *Pin.* 228. *Calamintha mont. praealta, pulegii*
odore. BOCC. *Mus.* II. p. 45. *T.* 40. In *Anglia*.
Belgio. P.

H. Melissa racemis terminalibus, pedunculis sa-
lutaris brevissimis. *Crepida*, LINN. *Sp.* 5. *Calamin-*
tha incana, ocyimi foliis. C. B. 569. *Calamintha*
Pulegii odore, flore minori, erecta, Virginiana.
HALL. G. 339.

CLINOPODIUM. HALL. 653.

LINN. 644.

Clinopodium verticillis densissimis, diffusis, fo-
lia tomentosis ovatis acuminatis. HALL. *H.* 653. n. 1.
LINN. *Sp.* 1. Clinopodium *Origano simile*. C. B. 224.
C. B. *Circa Lüneb. lungen.*

Clinopodium verticillis paucifloris, ab imo ad
summum dispersis. HALL. *H.* 654. n. 2. *Thymus*
LINN. *Sp.* 4. *Acinno* RIV. 47. Clinopodium *or-*
vense, Ocyimi facie. C. B. 225. *vulgare*. L. 506. In
collibus lapidosis. v. g. circa *Plafs.*

H. Clinopodium subrotundo acuminatis nervis
subferratis, caule procumbente, floribus ver-
ticil-

scillatis. Clinopodium perenne, Patgii odore, Majorana folio. BOCCON. HALL. *Ger.* 341.

THYMUS. LINN. 646. *Serpillum.*
HALL. 654.

H. *Thymus erectus*, foliis revolutis, ovatis, floribus verticillato spicatis. LINN. *Sp.* 2. *Thymus vulgaris*. RIV. *T.* 41. & *officinarum*. Fr.

Thymus floribus capitatis, foliis planis obtusis basi ciliatis, staminibus rubum floris excedentibus. *Serpillum* HALL. H. 1. *Thymus* LINN. L. 1. *Serpillum vulgare majus*, In apricis glareosis. Fr.

Thymus floribus capitatis, foliis planis obtusis basi ciliatis, staminibus in tubo floris latentibus. *Serpillum* HALL. H. 2. *Thymus* LINN. *Sp.* 2. *Serpillum vulgare minus*. Similibus locis. Flores duplo minores ac in priori. Fr.

B. *Serpillum foliis Citri odore.*

H. *Thymus* foliis ellipticis & caule hirsutus. HALL. H. 2. *Serpillum saxatile hirsutum, thymifolium, exanum flore rubello, & flore albo.* VAILL. *T.* XXXI. 1. 40. *Serpillum Pannonicum* III. GLYS. *Pann.* 2. 656. In sabuletis Cellensibus. Fr.

SATUREIA. LINN. 626.

H. *Satureia pedunculis bifloris. hortensis.* LINN. *Sp.* 7. *Max. med.* 284. *Satureia* RIV. *T.* 44. *Satureia*. 1. 2. III. p. 272. In Italia, Gallia. A.

H. *Satureia pedunculis lateralibus solitariis, trifloris, foliis carinatis, punctatis, ciliatis, mucronatis.*
V 5

LAVANDULA. LINN. 630.

H. *Lavandula foliis lanceolatis integris, spicis nudis. Spica.* LINN. *Sp.* 1. *α.* *Lavandula angustifolia.* C. B. 226. BLACKW. T. 294.

β. *Lavandula latifolia.* C. B. BLACKW. T. 295. In *Europa* australi. Fr.

H. *Lavandula foliis duplicato-pinnatifidis.* LINN. *Sp.* 2. *Lav. folio dissecto.* C. B. *Pta. l. c.* *Lavand. multifido folio.* CLVS. *Hist. l.* 345. In *Hispania*. Fr.

H. *Lavandula foliis serratis, spicis foliosis.* HALL. G. 342. LINN. *Sp.* 3. *Stoechas folio serrate.* C. B. *Stoechas. II. crispifolia.* CLVS. H. l. 346. *Stoechas serratifolia.* RIV. T. 55. In *Hispania*.

H. *Lavandula foliis lanceolato-linearibus, spica comosa.* LINN. *Sp.* 4. HALL. G. 1. c. *Stoechas arabica.* RIV. T. 55. *purpurea.* C. B. l. c. CLVS. L. 344.

ORIGANVM. LINN. 645.

Origanum spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. vulgare. LINN. *Sp.* 7. HALL. H. 656. *Origanum sylv. f. vulgare.* TABERN. 346. BLACKW. T. 286. *Stramonium ex tubo floris eminent.* Ad sepes & in collibus.

H. *Origanum spicis longis pedunculatis aggregatis, bracteis longitudine calycum.* LINN. *Sp.* 6. *Origanum spumibum in tubo floris larentibus.* HALL. G. 345. *Orig. Virens, officinum.* TABERN. 345. LOB. ic. 492. In *Europa* australi.

DICTAM-

DICTAMNVS. TOERN.

H. *Dictamnus Creticus*. HALL. G. 343. LOBEL. 46. 502. B. Pin. 222. *Origanum foliis inferioribus tomentosis, spicis nutantibus*. LINN. Sp. 2. *Bracteae in spica laxa imbricata magnae coloratae. Calycis labium inferius vix ullum, superius ovatum integrum*. Fr.

MAIORANA. TOERN.

H. *Maiorana*. HALL. G. 343. BLACKW. T. 319. RIV. T. 58. *vulgaris annua. β. perennis. Origanum foliis ovalibus obtusis, spicis subrotundis compactis pubescentibus*. LINN. Sp. 10. *Bracteae decedentes in spicis densis imbricatis tetragonis, calyx bifolius*.

MENTHA. LINN. 633.

H. *Mentha floribus spicatis, foliis cordatis dentatis undulatis sessilibus*. LINN. Sp. 1. *Mentha crispa Danica*. MORIS. III. S. XL. T. 6. f. 5. *Mentha crispa*. RIV. 47.

Mentha foliis elliptico-lanceolatis, verticillis in spicam nudam continuatis. HALL. H. n. 2. *Mentha spicata*. LINN. Sp. 2. *β. longifolia. Mentha sylvestris longiore folio*. C. B. *Menthastrum*. LOBEL. 46. 509. BLACKW. T. 292. In aquis vbiq.

H. *Mentha foliis ovatis tomentosis, verticillis spicatis, staminibus tubo floris aequalibus. Mentha*. HALL. H. 1. *Mentha sylv. rotundifolia*. L. B. H. LINN. Sp. 2. In fossis *Alsatiae*.

H. Men-

H. *Mentha foliis ovatis inferae tomentosis, verticillis spicatis, staminibus tubum floris superantibus. Menthastrum Anglicanum.* RIV. T. 51. plantam nostram egregie exprimit. *Flosculi carnei. Odeur plantae aromaticus, gratus.*

Mentha foliis ovato-lanceolatis, verticillis paucis, supremo maximo. HALL. H. 658. *Mentha floribus capitatis, foliis ovatis serratis petiolatis, staminibus corolla longioribus.* LINN. Sp. 3. *Mentha rotundifolia palustris.* C. B. In fossis.

Mentha floribus verticillatis, foliis ovatis acutis serratis, staminibus corollam aequantibus. LINN. Sp. 6. HALL. H. *Mentha verticillata arvensis.* BLACKW. T. 290. In agris.

Mentha floribus verticillatis, foliis ovatis acutis serratis, staminibus tubo floris brevioribus. gentilis. LINN. Sp. 5. *Mentha palustris verticillata.* DILL. Gieff.

H. *Mentha floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus.* LINN. Sp. 8. *Pulegium foliis subrotundis, verticillis nudis.* HALL. H. *Pulegium* BLACKW. T. 302. In *Helvetiae* inundatis. P.

H. *Mentha floribus verticillatis, foliis linearibus. Cervina.* LINN. Sp. 9. *Pulegium angustifolium.* MORIS. III. S. XI. T. 7. f. 7. RIV. Monop. 22. In *Europa australi.*

LYCOPVS. LINN. 31.

Lycopus foliis sinuato-serratis. LINN. Sp. 1. HALL. H. 660. *Pseudomarrubium palustre.* RIV. T.

T. 21. *Marrubium palustre glabrum*. G. B. Ad fossas.

β. *Marrubium palustre hirsutum, tenuius latiusculum.*

AMETHYSTINA. HALL. LINN. 32.

H. *Amethystina* Comment. Gott. T. I. Tab. X. p. 209. Act. Vps. 1742. p. 51. f. 1. *Amethystea*. LINN. Sp. p. 21. H. Vps. 9. Amoen. Acad. 386. *. *Amethystina montana erecta, foliis exiguis digitatis trifidis serratis, stoculis cum coma e coerulesco-janthinis.* AMM. Ruth. p. 54. n. 70. In Sibiria. A.

VERBENA. LINN. 30.

Verbena tetrandra, spicis filiformibus paniculatis foliis multifido-laciniatis, caule solitario. officinalis. LINN. Sp. 13. *Verbena foliis laciniatis, superioribus tripartitis, spicis angustis longissimis.* HALL. Helv. 661. *Verbena communis coerulesco flore.* C. B. 1. R. H. BLACKW. T. 41. Ubique ad vias. A.

H. *Verbena tetrandra, spicis filiformibus, foliis multifido-laciniatis, caulibus numerosis. spuria.* LINN. Sp. 12. H. Vps. 8. *Verbena urticae foliis Canadensis, foliis incisis, flore maiore.* R. B. H. HALL. G. 345. In Canada. P.

H. *Verbena spicis filiformibus paniculatis, foliis ovato-lanceolatis serratis petiolatis. urticifolia.* LINN. Sp. 11. HALL. G. 346. *Verbena recta Canadensis, urticae folio.* MORIS. HE. p. 418. & XL T. 25. f. 3. ZANON. 227. T. 175.

H. Ver-

Verbena foliis ovatis serratis sessilibus, spicis filiformibus paniculatis floribus quinque. *Verbena Caroliniana, Melissa folio.* DILL. Elth. T. 301. HALL. G. l. cir.² 357. WALT. 217.

H. *Verbena spicis longis acuminatis, foliis serratis inferioribus hatis, superioribus lineari-lanceolatis.* *hastata.* LINN. Sp. 16.² *Verbena Amer. altissima, spica multiplici, verticillae foliis angustis, floribus corollatis.* HERM. Parad. T. 242. HALL. G. l. c. P.

H. *Verbena spicis fasciculatis, foliis lanceolatis amplexicaulibus.* *Bonariensis.* LINN. Sp. 9. *Verbena Bonariensis altissima, Lavendulae Canariensis folio, spica multiplici.* DILL. T. 300. HALL. G. 347.

H. *Verbena spicis capitate conicis, foliis serratis, caule repente.* LINN. Sp. 8. *Verbena foliis verticaliter ovatis, spicis foliatis ovatis.* ROYEN. 327. *Verbena nodiflora.* C. B. Pin.^o 269. Prodr. 125. *In Virginia.*

TETRADYNAMIAE SILIQVOSAE.

Classi huic naturali calyx est quadrifolius, pene semper caducus, petala quatuor ex longiori vñgue in bracteam reflexa, more Caryophyllaeo. Stamina sex, quatuor maiora, duo minora, ad horum basi glandulae virides saepe adfident. Fructus plerumque bilocularis est aut brevis, subrotundus, oblongus, *siliolata* dictus, & stylo stigmate obtuso instructus; aut est longior, teres, *siliqua* dictus, stigma

te fere sessili. Vires omnibus sunt resolversc,
antiscorbuticae, sapor acer, natura alcalescens.

ISATIS. LINN. 738.

Silicula oblonga anceps unilocularis.

H. Isatis foliis radicalibus crenatis, caulinis sagittatis, siliculis oblongis. *tinctoria*. LINN. Sp. 1. Isatis *sylvestris*. C. B. BLACKW. 246. Circa Jenam. *β. sativa*. Vid. icon. SCHREBERI in proprio libello.

CRAMBE. LINN. 739.

Stamina apice bifido. Capsula rotunda.

H. Crambe foliis subcordatis, cauleque scabris. *hispanica*. LINN. Sp. 3. Rapastrum *maximum rotundifolium monospermum*. CORNYT. *Canad. T.* 148. MORIS. II. S. III. T. 13. f. 1. A.

BUNIAS. LINN. 737. Erucago.

TOURN. 103.

Silicula irregularis, tetragona aut gibba.

H. Bunias siliculis tetragonis; angulis bicipitatis. *Erucago*. LINN. Sp. 2. foliis *antrorsum sinuatis*. Cliff. 340. *Erucago Monspel. filiqua quadrangula echinata*. B. Prodr. T. 41. A.

H. Bunias siliculis ovatis gibbis verrucosis. LINN. Sp. 3. foliis *retrosum sinuatis*. Cliff. H. Vps. 186. *Crambe orient. dentis levis folio, erucaginis facie*. L. R. H. Cor. 14. P.

H. Bunias siliculis ovatis laevibus ancipitibus. *Cakile*. LINN. Spec. 4. *Crambe foliis succulentis lineis*.

linearibus pinnato-dentatis. SAUVAG. *Monsp.* 280.
Eruca marit. Italica, filiqua basiae cuspidi simili.
 B. Pin. 99. *Cakile Scrapionis.* LOB. ic. 223. Ad
 littora maris. A.

ANASTATICA. LINN. 715.

Silicula figura vngulae bovis, stylo medio
 longiori.

H. Anastatica. LINN. *Sp.* 641. H. *Vps.* 183.
Rosa Hierobuntica. CAMER. H. T. 41. C. B. Ad
 littora maris rubri. A.

VELLA. LINN. 714.

Silicula globosa, stylo ovato capsula longiori.

H. Vella foliis pinnatifidis, filiculis pendulis. *an-*
nuua. LINN. *Sp.* 2. Nasturtium *sylo. Valentianum.*
 GLVS. CXXX. In *Hispania.* A.

MYAGRUM. LINN. 713.

Silicula difformis, bi- aut trilocularis, loculo vai-
 co seminifero, reliquis abortivis.

H. Myagrum filiculis biarticulatis dispermis, foliis
 extrorsum sinuatis denticulatis. LINN. *Sp.* 1. *HP.*
Vps. 282. *. Myagrum *filiquae primo nodo frictio-*
ri, altera globoso striata. HALL. H. 536. Gott. 239.
Rapistrum monospermum. B. *Rodr.* T. 37. *ALAB.*
Alsat. T. 260. A.

H. Myagrum filiculis cordatis subsessilibus, foliis
 amplexicaulibus. *perfoliatum.* LINN. *Sp.* 5. H. *Vps.*
 182. *. Myagrum *filiculis utrinque gibbosis mono-*
sper-

Spermis. HALL. G. 1. c. *Myagrū monospermum*, B. Prodr. p. 52. Inter fegetes *Helictiae*. A.

H. *Myagrū siliculis globosis compressis punctato-rugosis.* LINN. Sp. 7. HALL. H. 537. *Myagro similis siliqua rotunda.* B. Prodr. T. 52. Circa *Berolinum*.

ALYSSVM. HALL. 537. LINN. 722.

Silicula subrotunda, stylo longiori. Habitus incanus.

Alysson foliis scabris sagittatis, leviter dentatis. HALL. G. 537. *Myagrū* LINN. Sp. 6. *Alysson fegetum, foliis auriculatis, acutis.* 1. R. H. *Myagrū* f. *Camelina*. Inter linum abunde. A.

G. *Myagrū (Alyssum) foliis auriculatis, glabris profunde dentatis.* HALL. H. l. c. n. 2. Ad fossas.

H. *Alyssum caule erecto, foliis lanceolatis incanis integerrimis, floribus corymbosis. incanum.* LINN. Sp. 5. *Alyssum fruticosum incanum* 1. R. H. *Thlaspi incanum Meublinianse.* L. O. B. ic. 216. CAVA CXXXII. *Petala bifida.* In arenosis, apricis. v. g. circa *Noribergam, Berolinum, Cellam.*

H. *Alyssum caule erecto herbaceo, siliculis sessilibus ovalibus compresso-planis, petalis acuminatis. alpestrum.* LINN. Sp. 7. *Leucorū alyssoides stippeatum maius.* B. Pin. 201. In *Europa australi.*

H. *Alyssum caule herbaceo, foliis lanceolatis dentatis, siliculis inflatis.* LINN. Sp. 8. *Alyssum fistulosum inflatis, foliis tomentosis.* HALL. G. 241. *Alyssoid-*

lyfoides incanum, foliis sinuatis. 1. R. H. *Eruca peregrina.* CLV. CXXXIV. *Petala bifida.*

Alyssum foliis imbricatis, lanceolato-ovatis integerrimis, siliculis tetraspermis, calyce persistente. Clypeola LINN. *Sp.* 2. HALL. *H.* 1. *Alyssum minimum.* CLV. CXXXIII. *Alyssum incanum luteum, Serpylli folio matur.* 1. R. H. A.

H. *Alyssum foliis imbricatis lanceolato-ovatis, punctatis, siliculis tetraspermis, calyce caduco.* HALL. *H.* 2. *Got. 240. Jonthlaspi luteo flore, incanum, discoides, umbellatum, montanum.* COLUMN. *Ecpbr. T.* 280. *Omnia maiora, stylus longior. In arenosis ad Rhenum.* P.

CLYPEOLA. LINN. 729.

Silicula orbiculata, plana, hilo modice excisa.

H. *Clypeola siliculis unilocularibus, monospermis.* LINN. *Sp.* 1. *Thlaspi clypeatum, serpylli folio.* LOB. *ic.* 214. *Jonthlaspi minimum spicatum luteum.* COLUMN. *Ecpbr. T.* 284.

DRABA. LINN. 717.

Silicula elliptica compressa, stylo minimo.

Draba caulibus nudis, foliis incisiss. LINN. *Sp.* 2. HALL. *H.* 538. *Alysson vulgare, polygont folio, caula nudo.* 1. R. H. SEGV. *Veron. I. T.* 4. f. 3. *Alfene minima.* TABERN. 708. *Verno tempore in macilentis.* A.

LVNARIA. LINN. 725.

Silicula elliptica, plana maxima. Semina reniformia.

H. *Lunaria filiculis oblongis. rediviva.* LINN. Sp.

1. *Lunaria foliis cordatis, perennis, siliqua longiore.* HALL. H. 540. *Lunaria maior siliqua longiore.* I. R. H. *Lunaria odorata.* CLVS. 297. In sylva *bercynica.* P.

H. *Lunaria filiculis subrotundis. annua.* LINN. Sp.

2. *Lunaria maior, siliqua rotundiore.* I. R. H. Num a priori specie satis diversa? Vita annua & fructus suadent.

BISCUTELLA. LINN. 724.

Siliculae duae plantae rotundae sepro affixae.

H. *Biscutella calycibus nectario vtrinque gibbis, filiculis in stylum coeuntibus. auriculata.* LINN.

Sp. 1. *Biscutella foliis rariter dentatis, siliqua aspera.* HALL. H. 2. *Thlaspidium biseutatum villosum, flore calcare donato.* B. Prodr. 49. *Leucoium montanum flore pedato.* COLVMN. Escpr. II. T. 61. A.

H. *Biscutella filiculis orbiculato-didymis a stylo divergentibus. didyma.* LINN. Sp. 7. *Biscutella folio aspero dentato, siliqua levi.* HALL. H. 1. *Jon-draba alyssoides apula spicata.* COLVMN. Escpr. I. T. 285. *Thlaspi clypeatum.* CLVS. CXXXIII. A.

COCHLEARIA. LINN. 720.

Silicula subcordata, scabra, loculis tetraspermis.

H. *Cochlearia foliis radicalibus subrotundis, caulinis oblongis subsinuatis. officinalis.* LINN. Sp. 1. *Cochlearia folio subrotunde.* C. B. A.

β. Var. *foliis integris.* CAMER. Epit. 271.

γ. *foliis*

β. *foliis serratis*. *Cochlearia Britannica*. LOBEL.
1594. Ad littora maris septentrionalis.

h. *Cochlearia foliis caulinis cordato-sagittatis
amplexicaulis glabris*. LINN. *Sp.* 7. *Lepidium
glaberrimum*. C. B. MORIS. H. S. III. T. 21. f. 3. A.

*Cochlearia foliis radicalibus lanceolatis crenatis,
caulinis incis. Armoracia*. LINN. *Sp.* 6. *Cochlearia
folio cubitali*. T. R. H. *Raphanus rusticus*. C. B.
In prato prope Hyperbansen. P.

CORONOPVS. HALL. H. 543.

*Silicula transversa lata, verrucosa, loculis mono-
spermis*.

α. *Coronopus Rubellus*. BLACKW. T. 120. *Nastur-
tium Salustre, capsulis cristatis*. T. R. H. *Cochlearia
foliis pinnatifidis*. LINN. *Sp.* 6. *Hinter der Mäsch-
mühle am Weg gegen den Berg hin, & ad fontes
salcos*. A.

IBERIS. LINN. 721.

Petala exteriora maiora.

h. *Iberis frutescens, foliis cuneiformibus inte-
gerrimis, obtusis, semperflorens*. LINN. *Sp.* 1. *Thlaspi
Persicum*. RIV. T. 110. *Thlaspi fruticosum Persicum
folio Cheiri*. ZANON. T. 165. Fr.

h. *Iberis frutescens, foliis linearibus acutis inte-
gerrimis, sempervivens*. LINN. *Sp.* 2. *Thlaspi fru-
ticosum sempervirens, flore alba Cretici*. BARREL.
734. In Creta.

h. *Iberis foliis lanceolatis acutis subdentatis, flo-
ribus racemosis*. LINN. *Sp.* 7. *Iberis foliis sursum*

Intescentibus dentatis. HALL. H. 543. *Thlaspi umbellatum arvense amaram.* L. R. H. TABERN. 462. In Helvetia.

H. Iberis foliis lanceolatis, acuminatis, inferioribus serratis, superioribus integerrimis, umbellata. LINN. Spec. 5. *Thlaspi umbellatum Genticum.* J. B. H. A.

H. Iberis foliis linearibus superne dilatatis, serratis. LINN. Sp. 8. *Thlaspi pernon IV. odorato folio.* CLVS. CXXXII.

Iberis foliis sinuatis, caule nudo simplici. nudicaulis. LINN. Sp. 9. HALL. H. 1. *Bursa pastoris minor, foliis incis.* B. Pin. 108. *Nasturtium pratense.* TABERN. 451. In rupibus sub Hecken. A.

NASTURTIVM. HALL. 543. *Thlaspi.* LINN. 719.

Silicula subrotunda, apice hilo exsculpta; stylo hylo aequalis aut brevior.

Nasturtium (Thlaspi) foliis oblongis dentatis glabris, filiculis orbiculatis. HALL. H. 5. LINN. Sp. 2. *Thlaspi arvense, filiquis latis.* G. B. I. R. H. In agris. A.

Nasturtium (Thlaspi) foliis imis ovatis, reliquis amplexicaulibus, cordatis, serratis, filiculis subrotundis. HALL. H. 4. LINN. Sp. 5. *Thlaspi persoliatum minus.* G. B. I. R. H. *Thlaspi III. pumilum.* CLVS. CXXXI. In vallo urbis & agris. A.

Nasturtium (Thlaspi) filiculis subrotundis, foliis sagittatis, dentatis incanis. HALL. H. 3. LINN. Sp. 7. *Thlaspi II.* TABERN. 458. In muris.

β. varis.

... *β. varietas foliis glabris vix serratis. Thlaspi vulgare.* BLACKW. T. 107. In agris. A.

H. Nasturtium (*Thlaspi*) siliculis subrotundis pilosis, foliis caulinis sagittatis hirsutis. LINN. Sp. 6. *Thlaspi villosum capsulis hirsutis.* B. Prodr. T. 222. In Italia. B.

H. Nasturtium (*Thlaspi*) siliculis obcordatis, foliis radicalibus cuneiformibus integerrimis, summis amplexicaulibus sagittatis. LINN. Sp. 8. *Iberis foliis imis cordatis, superioribus amplexicaulibus.* HALL. G. 243. *Lepidium Helv. 1. Thlaspi mont. II. Badense.* CLVS. CXXXI. *Thlaspi mont. bursae pastoris folio.* COLUMN. Ecphr. T. 276. *Flores exteriores leviter inaequales.*

H. Nasturtium (*Lepidium*) siliculis ellipticis, foliis pulposis subrotundis, antheris lateralibus. HALL. G. 241. *Thlaspi* LINN. Sp. 4. *Lithonthlaspi quartum carnosum rotundo folio.* COLUMN. Ecphr. I. T. 277. f. 2. A.

H. Nasturtium (*Lepidium*) foliis tetradynamis, foliis oblongis multifidis. LINN. Sp. 6. *Nasturtium bortenfe.* BLACKW. T. 23. *β. varietas foliis crispis.* B. Prodr. T. 43. A.

Nasturtium (*Thlaspi*) foliis radicalibus, pinnatifidis, siliculis verticaliter cordatis. LINN. Sp. 9. HALL. H. n. 6. *Bursa pastoris.* BLACKW. T. 5. V-bique in gramineis. A.

Nasturtium (*Lepidium*) floribus diandris foliis imis multifidis, superioribus linearibus integerrimis. HALL, H. 1, LINN. Sp. 10. *Nasturtium sylvestre*

Oxyridis folio. C. B. Iberis. BLACKW. T. 312. In muris v. g. Auf der Albaner Kirchhof. Murr. A.

H. Nasturtium (*Lepidium*) floribus dianthis minutissimis, foliis linearibus pinnatis. LINN. Sp. 11. Iberis. humilior, annua, ramifera, Virginiana. MORIS. II. S. III. T. 21. f. 2. A.

LEPIDIVM. LINN.

Silicula subcordata, apice non emarginata; stylus capitatus.

H. Lepidium foliis caulinis pinnatis, multifido ramiferis cordatis amplexicaulibus integris. LINN. Sp. 1. HALL. Gott. 245. Thlaspi venum Dioscoridis. ZANON. T. 129. Nasturtium Orient. foliis inferioribus Millefolii, superioribus Persiliatæ. I. R. H. A.

H. Lepidium foliis pulposis ovato lanceolatis serratis. LINN. Sp. 8. Lepidium latifolium. C. B. I. R. H. Circa Spiram; in Anglia. P. A.

H. Lepidium foliis lanceolatis amplexicaulibus dentatis. LINN. Sp. 14. Draba umbellata. C. B. Draba I. vulgaris. CLVS. II. CXXIII. In Germania australi. P.

II. Fructu longiori s. siliqua.

SISYMBRIUM. HALL. 547. LINN. 728.

Erysimum. LINN. 729.

Calyx deorsum non gibbus. Stylus brevissimus.

Sisymbrium siliquis declinatis, foliis pinnatis, foliolis subcordatis. LINN. Sp. 1. Sisymbrium foliis pinnatis, angulosis, floribus super folia vix elatis.

HALL.

HALL. H. *Nasturtium aquaticum*. C. B. BLACKW. T. 260. Ad fontes. P.

Sisymbrium siliquis declinatis, foliis pinnatis, foliolis lanceolatis serratis. LINN. Sp. 2. HALL. H. *Sisymbrium patuliflorum repens, Nasturtii folio*. I. R. R. *Erica sylv. luteo flore*. C. B. Ad vias & fossas.

Sisymbrium siliquis declinatis oblongo-ovatis, foliis pinnatifidis serratis. LINN. Sp. 3. HALL. H. 3. *Raphanus aquaticus, foliis in profundas lacinias divisis*. B. Prodr. 7. 38.

8. *Sisymbrium* foliis imis integris ovatis, serratis, superioribus pinnatis. HALL. H. 2.

9. *Sisymbrium* foliis simplicibus dentatis serratis. HALL. H. 1. *Raphanus aquat. alter*. B. Prodr. 37. Ad rivos & fossas. P.

H. *Sisymbrium* siliquis axillaribus sessilibus subulatis aggregatis, foliis repando dentatis. LINN. Sp. 5. HALL. H. 10. *Erysimum polyceratum, s. corniculatum*. B. Pin. 101.

Sisymbrium petalis calyce minoribus, foliis decomposito-pinnatis. LINN. Sp. 11. *Sisymbrium pinnis foliorum, pinnatis, pinnulis subhirsutis linearibus*. HALL. H. 7. *Nasturtium sylv. renursum divinum*. C. B. *Sophia Chirurgorum*. BLACKW. T. 240. Ad muros vias. A.

H. *Sisymbrium* foliis pinnatis, extrema pinna triangulari dentata, petiolis longis hirsutis. HALL. G. 248. *Sisymbrium foliis pinnato-bastatis, acutibirtis, caule retrorsum hispido*. LINN. Cent. Pl. I. n. 51. *Erysimum mont. Irionis folio, macroleptacera*.

craton. COLVYN. *Ecphr.* I. T. 268. In totis fo-
mineis *Eislebiae*.

H. *Sisymbrium* foliis pinnato-hastatis flaccidis,
foliis sublinearibus integerrimis, pedunculis lan-
atissimum. LINN. *Sp.* 13. *Erysimum Monspellu-*
num, *Sinapis folio*, WALTH. *Hort.* 55. T. 22.
Rapistrum sylvestre Irionis folio. COLVYN. *Ecphr.*
I. T. 268. In *Gallia*. A.

H. *Sisymbrium* foliis pinnato-hastatis tomentosis,
caule levi. LINN. *Cent. Pl.* II. n. 173. *Erysimum*
Orient. sonchi folio, *flore sulphurea*, *siliquis longis-*
simis. BOERH. I. p. 14. HALL. G. 248. A.

H. *Sisymbrium* foliis pinnatis, extrema pinna am-
plissima, siliquis praelongis. HALL. G. 247. *Hespe-*
ris foliis dentato-pinnatifidis, *caule laevi*. LINN.
Sp. 6. *Hesperis flore albo minimo*, *siliqua longa*,
folio profunde dentato. DILL. *Elth.* T. 148. f. 177.
In *Sicilia*.

Sisymbrium foliis lyratis, pinna extrema amplis-
sima, subrotunda. HALL. H. 6. *Erysimum* LINN. 2.
Barbarea femina. TABERN. 452. Ad fossas. A.

Sisymbrium caule ramosissimo, foliis inferioribus
angulatis sinuatis, superioribus dentatis, siliquis nodo-
sis. *Sisymbrium foliis imis Barbareae*, *superioribus*
integris dentatis. HALL. *It. Hercyn.* c. 16. *Opusc.*
bot. p. 100, n. XL. Ad rivos *hercynicos*.

Sisymbrium (*Erysimum*) siliquis cauli adpressis.
HALL. H. 9. LINN. *Sp.* 1. *Erysimum* BLACKW.
T. 28. *vulgare*. C. B. I. R. H. Ad vias. A.

Sisymbrium (Erysimum) foliis cordatis. HALL. H. 15. LINN. Sp. 3. Alliaria. C. B. BLACKW. T. 372. Ubique verno tempore ad fepes. A.

Sisymbrium (Erysimum) foliis integris lanceolatis subasperis. *Cheiranthoides*. HALL. H. 8. LINN. Sp. 4. Turritis *Leucoi folio*. I. R. H. Camelina, *Myagrum alterum Tblaspi effigie*. LOB. ic. 225. Ad fossas. A.

ERVCA.

Calyx neque gibbus, neque patulus. Stylus priori longior, stigmate capitato.

H. Eruca (*Sinapi*) foliis pinnatis, ferratis, extrema maxima, filiquis longissimis. HALL. G. 249. *Sisymbrium foliis lyratis denticulatis, filiquis sessilibus*. LINN. Sp. 6. Eruca *sylv. minor lutea, bursae pastoris folio, Italica*. BARREL. ic. 1016. A.

H. Eruca (*Sisymbrium*) foliis glaucis, pinnatis, pinnis longis, rariter ferratis. HALL. H. 11. *Brassica foliis dentato pinnatifidis, caule hispido, filiquis laevibus*. LINN. Sp. 7. Eruca *sylvestris*. BLACKW. T. 268. Eruca *rennifolia, perennis, flore luteo*. I. R. H. In Palatinatu, Helvetia. A.

H. Eruca (*Brassica*) foliis lyratis, caule hirsuto, foliis glabris. LINN. Sp. 8. Sinapi *filiquis tetragonis, floribus venosis*. HALL. G. 250. Eruca HELV. 593. BLACKW. T. 242. Eruca *latifolia alba sati-va*. C. B. I. R. H. flore albo & luteo, utroque venoso. In Valtellia. A.

H. Eruca (*Sinapi*) filiquis laevibus aequalibus, foliis lyratis oblongis glabris, caule scabro. LINN.

Cent.

Cent. Pl. H. n. 174. Eruca (Dioscori), flos albo, italica. BARREL. ic. 132. Caulis retrofusus scaber; pedunculis similes patentibus. Calyx 6. ungues petalorum alborum exsiccatis videtur evadunt; stigma bicolle.

SINAPIS. LINN. 735.

Calyx patens. Glandulae virides quatuor. Stylis anceps, filiquae aequalis.

1. *Sinapis filiquis multangulis torosis turgidis, rostro longioribus. arvensis. LINN. Sp. 1. HALL. H. 3. Sinapi arvensis praecox, semine nigra. R. H. In agris & hortis. A.*

2. *H. Sinapis filiquis retrorum hispidis, apice latetragonis compressis. LINN. Cent. Pl. 1. n. 55. Sinapi Orient. maximum, rapi folio. I. R. H. Cor. 17. HALL. G. 236.*

3. *Sinapis filiquis hispidis, rostro obliquo longissimo. LINN. Sp. 2. HALL. H. 2. Sinapi album. BLACKW. T. 12. Sin. filiqua hirsuta, semine albo & rufo. I. R. H. Sin. api folio. C. B. In agris & hortis. A.*

4. *Sinapis filiquis glabris tetragonis. nigra. LINN. Sp. 3. Mat. med. 316. HALL. H. 1. BLACKW. 466. Sinapi rapi folio. C. B. I. R. H. In ruderatione. Haec species ob usum medicum & diaeteticum celebris. A.*

5. *H. Sinapis ramis fasciculatis, foliis inferioribus ovatis lacertis, summis lanceolatis integerrimis. LINN. Sp. 4. H. Vps. 191. *. Sinapi Ind. maximum, lactucae*

laetiferae folio. HERM. PAR. T. 230. *Sinapi Chinesse,*
Acanthi folio. GMELIN. A.

RAPHANVS. LINN. 736.

Calyx erectus, gibbus. Siliqua acuminata, articulata.

H. Raphanus siliquis teretibus torosis bilocularibus. *latus.* LINN. Sp. 1. *Raphanus bortenensis,*
 BLACKW. T. 81. *oblongus rotundus.* C. B. A.

Raphanus siliquis teretibus articulatis laevibus
 unilocularibus. LINN. Sp. 2. *Raphanistrum arvense,*
flore albo, & luteo. I. R. H. *Rapistrum flore albo,*
luteo. C. B. A.

B. Variet. *flore albo venoso.* *Lampfana apula* Plinii & Dioscoridis. COLUMN. Euphr. L p. 261. In
 agris auf dem Hagen, und hinter Grobnde. A.

DENTARIA. LINN. 726.

Dentaria foliis inferioribus pinnatis, superioribus simplicibus. *bulbifera.* LINN. Sp. 2. HALL. H.
 3. *Dentaria heptaphyllus baccifera,* CLVS. CXXI.
In alis foliorum profert bulbos nigros congestos, qui delapsi speciem propagant; siliquae raro maturescunt.
In sylvis circa Gottingam & beresynicis.

BRASSICA. LINN. 734.

Brassica radice cauleque tenui, foliis caulinis unilobatis cordatis sessilibus. *campestris.* LINN. Sp.
 2. *Brassica arvensis perfoliata, flore luteo.* Inter
 getes. A.

Brassica radice caulescente fusiformi. LINN. Sp. 3.
Napus sylvestris, sativa. C. B. In agris seritur. A.

H. Bras-

H. Brassica radice caulescente, terete, carnosa
 LINN. Sp. 5. Var. *bortenses variae*, *a. capitata*;
β. Sabauda, *γ. crispa*, *δ. laciniata*, *ε. cauliflora*,
h. Napo-brassica, *θ. Canlo-rapa* s. *gongylodes*. A.

H. Brassica radice caulescente orbiculari depressa
 carnosa. LINN. Sp. 4. Rapum. BLACKW. T. 231.
Rapa sylvestris, sativa, rotunda aut oblonga. A.

CARDAMINE. LINN. 727.

Vngues petalorum calyce duplo longiores. Stylus brevissimus, stigma integrum.

Cardamine foliis pinnatis, pinnis subrotundis, angulosis. HALL. H. 3. Card., *axillis stoloniferis*, (*amara*.) LINN. Sp. 12. Nasturtium aquae. maius & amarum. B. Prodr. 45. In valle ultra Hessel-dreisch. A.

Cardamine foliis pinnatis, imorum pinnis subrotundis, caulinis oblongis & lanceolatis. LINN. Sp. 11. HALL. H. 4. Cardamine *pratensis*, *flore magno purpurascens* & *albo*. I. R. H. Ubique in pratis vdis. A.

Cardamine caule foliisque hirsutis. HALL. H. 1. LINN. Sp. 10. Cardamine *pratensis*, *parvo flore*. I. R. H. In sylva Hercynica.

H. Cardamine foliis pinnatis, foliolis aequalibus lobatis, petiolatis. HALL. H. 252. *graeca*. LINN. Sp. 9. Cardamine *Sicula*, *fumariae folio*. I. R. H. Sio minimo affinis, *siliquis latis*. BOCC. Sic. 84. T. 44. f. 2. In Sicilia. A.

Carda-

Cardamine foliis pinnatis, pinnis laciniatis, HALL. H. 1. *impatiens*. LINN. Sp. 8. *Cardamine anthus*, *exiguo flore*. I. R. H. In sylvis circa *Göttingam* & *Hercynicis*.

H. *Cardamine* foliis pinnatis, foliolis lanceolatis basi unidentatis, *Virginica*. LINN. Sp. 14. Nasturtium *bursae pastoris folio*, *Virginianum*, *flore albo*, *siliqua compressa*, PLUKN. T. 101. fig. 4. A.

TURRITIS. LINN. 933.

Calyx erectus. Siliquae longissimae, compressae, cauli parallelae.

Turritis foliis radicalibus dentatis hispidis, caulinis integerrimis, amplexicaulibus glabris. LINN. Sp. 1. HALL. H. *Turritis foliis inferioribus Gichoreacis*, *superioribus Perfoliatae*. I. R. H. In sylvis. A.

Turritis foliis omnibus asperis, dentatis, caulinis amplexicaulibus. LINN. Sp. 2. HALL. H. 3. *Erythimo similis hirsuta, non lucinata, alba*. SAVH. Prodr. T. 42. In lapidosis. A.

ARABIS. LINN. 732.

Calycis folia deorsum gibba, in quibus glandulae. Siliquae laxae dispositae.

H. *Arabis* foliis amplexicaulibus dentatis, caule diffuso ramoso. LINN. Sp. 1. B. HALL. H. 1. *Draba* II. GLV. CXXIV. *Draba alba siliquosa*. G. B. In alpidibus *belveticis*. P.

Arabis foliis ovatis, vix dentatis, scabris, ad terram confertis. HALL. H. p. 562. *Arabis foliis petio-*

petiolatis lanceolatis integerrimis. (Thaliano.)
 LINN. Sp. 3. *Pilofella filiquata minor.* THAL.
 HARC. 84. T. 7. f. D. In agris sterilibus M. Hagen &
 in vallécula inter Waaken & Besigebansen. A.

H. Arabis (*Sisymbrium*) foliis lyrato-ferratis sca-
 bris radicalibus, caule subnudo ramoso. LINN. Sp. 8.
Sisymbrium HALL. 14. *Eruca coerulea in arenosis*
proveniens. C. B. I. R. H. In *Helvetia.* A.

H. Arabis foliis amplexicaulibus, filiquis ancipi-
 tibus linearibus, calycibus subpilosis. *pendula.*
 LINN. Sp. 6. H. Vps. 191. *. *Turritis latifolia hir-*
suta, filiquis pendulis. AMMANN. Rusb. 58. HALL
 G. 253. In *Sibiria.* A.

CHEIRANTHVS. LINN. 730.

Petala calyce gibbo longiora. Siliqua compressa,
 stigmate bifido.

Cheiranthus foliis lanceolatis, acutis, glabris. *Cheir.*
 LINN. Sp. 2. HALL. H. 563. *Leucoium luteum*
vulgare. C. B. I. R. H. *Leucoium aureum.* MATH.
 877. Sponte interdum in muris. v. g. bey dem Ob-
 servatorio. A.

β. Var. flore pleno. C. B.

H. *Cheiranthus* caule suffruticoso, foliis dense
 congestis, obverse ovatis, lanceolatis, glabris, inte-
 gerrimis. Rami sub floribus dense, foliis tecti, infe-
 rius nudi. Calyx ruber. Flores grandes, speciosi,
 kermesini coloris, fere unciales. Siliqua viridis,
 compressa, saepe undulata. An *Cheiranthus* simili
 fere nomine HALL. G. 255? Videtur ex descriptio-
 ne

nae florum, sed synonyma aliena esse videntur, Optime nostram exprimit *Leucoium hyemale diu durans*, folio viridi & livida glabre. I. B. III, 877. etsi florem album dicat, nec ablunderet *Leucoium marinum maius*. CLVS. 298. & *Keiri marin. flore candido*. LOB. ic. 332. nisi CLVSIVS annumam illam diceret. Fr.

H. *Cheiranthus* foliis lanceolatis integerrimis ob-
tusis incanis, filiquis apice truncatis, compressis,
caule suffruticoso. LINN. Sp. 4. *Leucoium incanq*
folio, hortense. C. B. flore albo, purpureo, violaceo,
pleno. *Viola alba & purpurea*. LOB. ic. 329. Fr.

H. *Cheiranthus* caule ramoso diffuso, foliis lan-
ceolato ovatis retusis, raris, integris, filiquis apice
subulatis. An *Cheiranthus Ebius*. LINN. Sp. 3.
Leucoium tblaspiot. fucie, flore albo. HERM. *Parad.*
T. 193. optime nostram exprimit. Minus convenit
Hesperis filiquis hirsutis, flore parvo rubello. DILL.
Elieb. T. 147. f. 178. A.

H. *Cheiranthus* filiquarum apicibus tridentatis.
LINN. Sp. 9. *Leucoium marinum*. CAMER. *hort.*
87. T. 24. MORIS. II. p. 242. S. III. T. 8. f. 13. In
maritimis. A.

HESPERIS. LINN. 731.

H. *Hesperis* caule ramoso, foliis lanceolatis, den-
tatis scabris, corollis minimis, filiquis longissimis,
teretibus, sessilibus. An *Hesperis Africana*. LINN.
Sp. 4. *Hesperis Afric. hieracii folio hirsuto*, flore
minimo purpurascens. NISSOL. *Act. Par.* An *He-*
speris sylvestris, parvo flore. u. *Prodr. 193.* Folia

dentata, incisuris raris, uno alterove denticulo notatis. Siliquae longissimae, teretes, fere sessiles, hispidae, stylo vix ullo, apice acuto. A.

H. Hesperis caule hispido ramoso patente. *triflis*. LINN. Sp. 1. Hesperis *peregrina*, siliquis articulatis. I. R. H. Hesp. Syriaca. CAMER. Hort. ic. 19. Hesperis *montana pallida odoratissima*. C. B. Hesperis. I. CLVS. 296.

Hesperis caule simplici erecto, foliis ovato-lanceolatis denticulatis, petalis mucrone emarginatis: *matronalis*. LINN. Sp. 2. Hesp. *styl.* III. CLVS. 297. Hesp. *styl. inodora*. I. R. H. HALL. Rupp. Fl. Jen. p. 78. n. 2. T. 1. In ruderosis ad arcem Pless & Grubenhagen.

β. Hesperis *bortensis*. C. B. Viola *matronalis*. DODON. Pempt. 161. & *bortulanorum*, flore albo, purpureo, utroque pleno,

γ. Hesperis caule erecto ramoso, foliis cordatis amplexicaulibus serratis villosis. LINN. Sp. 7. Leucoium *marinum alterum latifolium purpureo-violaceum*. LOB. ic. 333. Leucoium *marit. latifolium*. C. B. MORIS. Hist. II. p. 241. S. III. T. 8. f. 5.

Hesperis caule recto, foliis scabris, lanceolatis, rariter dentatis, siliquis tetragonis. Cheiranthus *erysimoides*. LINN. Sp. 1. Leucoium *luteum sylvestre bibracifolium*. G. B. HALL. Rupp. Fl. Jen. p. 77. n. 2. Leucoium *sylvestre*. CLVS. Hist. 299. HALL. G. 257. In M. Altenstollberg.

δ. Hesperis caule ramosissimo, folioso, foliis linearibus, rariter dentatis, glabris, siliquis teretibus, corollis vix calyce maioribus. *Inter folia longe linearia,*

neuria, obtusior sine, patula, quae caulem undique
 desistant, spicae florum minorum provenit. Siliquae
 longissimae ultra flosculos superiores elatae.

Hesperis exigua, flore luteo, folio angusto, den-
 tato. BOERH. HALL. G. 256.

CLEOME. LINN. 240.

Siliqua longe petiolata. Petala sursum flexa.

H. Cleome floribus gynandris, foliis digitatis.
 LINN. Sp. 2. Fl. Zeyl. 239. * Sinapis trum Indi-
 cum pentaphyllum, flore carneo non spinosum.
 HERM. Lugdb. 564. In regionibus calidissimis. A.

H. Cleome floribus dodecandris, foliis ternatis.
 LINN. Sp. 5. Sinapis trum Lusitanicum, triphyllum,
 siliqua maxima, flore albo. I. R. H. HALL. G.
 257. A.

H. Cleome floribus hexandris, genitalibus de-
 clinatis, siliquis subulatis, foliis ternatis lineari-lan-
 ceolatis. LINN. Sp. 7. HALL. G. 258. Sinapis trum
 Lusit. triphyllum, flore rubro, siliquis corniculatis.
 I. R. H. A.

PAPILIONACEAE DIADELPHIAE

Calyx monophyllus quinquesidus. Corolla ire-
 gularis, tetrapetala. Petalum supremum reliquis im-
 positum sine reflexo, Vexillum dictum. Lateralla
 aequalia, alae dicuntur. Infimum bipes, carinae
 nomine, compressum, stamina & fructum conti-
 ner. Stamina novem plerumque oriuntur ex vagina
 communi, fructum involvente, decimum solitarium

ex fundo calycis. Saepe omnium filamenta connata. Fructus unilocularis, siliquosus, bivalvis. Semina farinosa, vires blandae emollientes.

I. Fructus bilocularis.

GLYCINE. LINN. 797. Apios.

H. *Glycine* foliis pinnatis ovato-lanceolatis. LINN. *Sp.* 1. *H. Kpf.* 327. *Apios Americana*. CORNYT. *Canad.* T. 201. RIV. T. 165. In *Virginia*. P.

BISERRULA. LINN. 800.

Biserrula. LINN. *Sp.* 762. *Pelecinus*. I. R. H. *Secundaca peregrina*. CLVS. CCXXVIII. In *Europa australi*.

ASTRAGALVS. LINN. 799. HALL. 565.

H. *Astragalus caulescens*, spicis cylindricis subsessilibus, calycibus leguminibusque lanatis. *alopecuroides*. LINN. *Sp.* 1. *Amoen. Acad.* 2. p. 359. *. *Astragalus alpinus procerior alopecuroides*. I. R. H. 416.

H. *Astragalus caulescens erectus pilosus*, floribus spicatis, leguminibus subulatis pilosis. LINN. *Sp.* 4. *H. Vpf.* p. 226. n. 4. *Astragalus caule erecto, ex alis spicifero, siliquis teretibus bisfutis*. HALL. *Comm. Soc. Gott. Tom. II. T. XII. Astrag. villosus, erectus, floribus flavescentibus*. AMMANN. *Ruth.* 166. P.

H. *Astragalus caulescens strictus*, glaber, floribus racemosis pendulis, leguminibus triquetris utrinque mucronatis. *galegiformis*. LINN. *Sp.* 7. *. *Astrag.* AMMANN. 162.

H. *Astra-*

H. *Astragalus caulescens, erectiusculus, floribus spicatis, leguminibus erectiusculis nudis tumidis tereti-depressis; mucrone reflexo. uliginosus. LINN. Sp. 8. H. Vpf. 226. n. 3. ** Astrag. *pedunculis, foliis supparibus, floribus pendulis, siliquis erectis. Siliquis parum compressis.* Descripsi hanc plantam in *Comm. Gott. T. V. P.*

H. *Astragalus caulescens erectus laevis, pedunculis spicatis, leguminibus ovato-cylindricis stylo acuminatis. LINN. Sp. 9. Astrag. procerior non repens; flore viridi flavescente. DILL. Elth. T. 39. f. 45. In Virginia. P.*

H. *Astragalus caulescens prostratus, leguminibus subglobosis inflatis mucronatis pilosis. LINN. Sp. 11. H. Vpf. 226. Astrag. HALL. H. 11. Astrag. luteus, perennis, siliqua gemella, rotunda, vesicam referente. I. R. H. Cicer sylvestre, foliis oblongis, hispidis, maius. c. B. Circa Jenam & in Siberia. P.*

Astragalus caulescens prostratus, leguminibus subtriquetris arcuatis, foliis ovatis pedunculo longioribus. LINN. Sp. 13. Astragal. HALL. H. 1. Astrag. luteus perennis procumbens vulgaris. I. R. H. Glaux vulgaris, Glycyrrhiza sylv. Gesneri. CLVS. CCXXXIII. In pratis & sylvis. P.

H. *Astragalus caulescens procumbens, leguminibus subulatis recurvatis glabris. LINN. Sp. 14. H. Vpf. 5. Astrag. luteus annuus Monspeliacus, procumbens. I. R. H. Astragalus Monspellatus, CLVS. CCXXXIV. Monspelii. A.*

H. *Astragalus caulescens procumbens*, leguminibus contorduplicatis canaliculatis, villosis. LINN. *Sp.* 15. *H. Vss.* 227. n. 8. *Astrag. siliquis recurvis depressis hirsutis*. HALL. *Comm. Gott.* T. II. p. 339. *Tab. XI. In Sibiria.*

H. *Astragalus caulescens procumbens*, spicis pedunculatis, leguminibus prismaticis rectis triquetris apice uncinatis. LINN. *Sp.* 16. *Astrag. triangularis*. MONTENGU *id.* p. 110. *Astrag. Baeticus*. RIV. T. 105. A.

H. *Astragalus caulescens diffusus*, capitulis subsessilibus lateralibus, leguminibus erectis subulatis acutissime reflexis. LINN. *Sp.* 17. *Vicia Sesamacea apula*. COLUMN. *Ecphr.* I. p. 303. T. 301. *In Gallia Narbonensi.* A.

H. *Astragalus caulescens procumbens*, leguminibus capitatis cordatis acutis reflexis complicatis. LINN. *Sp.* 18. *Astragalus Hisp. siliqua epiglottidis simili, flore purpureo, maior*. HERM. *Lugd.* 74. T. 75. *Glaux Hispanica*. RIV. T. 109. A.

PHACA. LINN. 798.

H. *Phaca caulescens erecta* tota glabra sulcata, pedunculis folio longioribus, leguminibus erectis subtriquetris. In *Comm. Gott.* T. V. descripsi, addita icone. *Astragalus caulescens erectus, caule glabro sulcato*. LINN. *Sp.* 6. *Foliola* 8. *f. 9. parum linearis lanceolata glabra, Pedunculi, floribus delapsis, foliis triplo longiores. Spica laxa pauciflora flores parvi, violacei. Legumina unilocularia, subtri-*

triquetra, inflata percurrit fasciâ, tanquam fœpæ rudimentum. P.

H. Phaca pedunculis radicatis, hirsutissimis, foliolis ex ovato acutis saepe imbricatis. *Comm. Got.* T. V. c. 16. descripsi. *Astragalus acaules, scapo erecto foliis longiore, leguminibus subulatis inflatis, villosis erectis.* LINN. Sp. 27. H. Vpf. *Astragalus non ramifusus villosus & incavus spicatus, floribus purpureo-violaceis.* AMMANN. Ruth. 167.

PSORALEA. LINN. 801.

H. Psoralea foliis pinnatis, foliolis linearibus acuminatis, floribus axillaribus. *pinnata.* LINN. Sp. I. *Genistae affinis arbor Afric. monosperma, flore coeruleo, foliis pinnatis.* HERM. Lugdb. T. 273. In *Aethiopia.* Fr.

H. Psoralea foliis omnibus ternatis, pedunculis spicatis folio longioribus. LINN. Sp. 4. *Vulneraria foliis ternatis, floribus umbellatis.* HALL. G. 262. *Trifolium bituminosum.* DOD. Pempt. 566. RIV. T. 14. In *Italia, Gallia australi.* Fr.

H. Psoralea foliis simplicibus ovatis. *corylifolia.* LINN. Sp. 6. H. Vpf. 225. *Pfor. folio integro subrotundo.* JUSSIEV. *Ag. Par.* 1744. p. 382. *Loto affinis coryli folio.* PLUKN. Alm. 227. T. 96. f. 5. In *India.* A.

VULNERARIA. HALL. 569. TOURN. 211.

Anthyllis LINN. 773.

H. *Vulneraria (Anthyllis)* herbacea, foliis quaternopinnatis, floribus lateralibus. LINN. Sp. I. H.

Vps. 221. n. 1. *Vulneraria pentaphyllos*. HALL G. 261. *Anthyllis Hispanica*. RIV. T. 18. *Trifolium balicaphum*. CAMER. hort. 171. T. 47. In *Italia*. A.

Vulneraria foliis ad terram simplicibus, ovatis, ad caulem pinnatis. HALL H. 2. p. 570. *Anthyllis herbacea*, foliis pinnatis inaequalibus, capitulo duplicato. LINN. Sp. 2. *Vulneraria rufica*. 1. R.H.

β. *Vulneraria supina*, flore coccineo. DILL. Elth. 431. T. 320. f. 413. In *Helvetia*. P.

η. *Vulneraria* foliis pinnatis sericeis nitentibus. HALL. G. 261. *Anthyllis fruticosa*, foliis pinnatis aequalibus, floribus capitatis. LINN. Sp. 6. H. *Vps.* n. 3. *Barba Iovis*. C.B. RIV. T. 4.

DORYCNIVM. LINN. ed. pr. 602.

TOURN. 211.

η. *Dorycnium* foliis quinatis digitatis sessilibus. LINN. Cliff. 371. HALL. H. p. 570. *Lotus* LINN. Sp. 15. *Dorycnium Hispanicum*. CLVS. 100. *Monspeliensium*. LOBEL. ic. 51. In *Europa australi*. A.

LOTUS. LINN. 803. HALL. H. 571.

η. *Lotus* leguminibus solitariis membranaceo-quadrangulatis, bracteis ovatis, (caule erecto.) LINN. Sp. 2. *Lotus pulcherrima tetragonolobos*. CAMER. Hort. 91. T. 16. In *Sicilia*. A.

η. *Lotus* leguminibus solitariis membranaceo-quadrangulatis, bracteis ovato-lanceolatis, caule pro-

procumbente. HALL. H. 1. Lotus *siliquosus luteus minor & mollior.* 1. R. H. Lotus *tetragonolobus.* RIV. T. 39. In *Thuringia.* A.

H. Lotus leguminibus subbinatis linearibus compressis nutantibus. LINN. Sp. 6. Lotus *siliquis geminis, peregrinis.* BOERH. Lugdb. II. p. 98. HALL. G. 262. In *Europa australi.* A.

H. Lotus leguminibus subquinatis arcuatis compressis, caulibus diffusis. LINN. Sp. 8. Lotus *siliquis Ornithopodii.* C. B. CAMER. Hort. 91. T. 25. In *Sicilia.*

H. Lotus leguminibus subternatis, caule herbaeo erecto, foliis linearibus. LINN. Sp. 9. Lotus *angustifolia, flore luteo purpurascente, insulae S. Jacobi.* COMMEL. H. II. p. 165. T. 83.

H. Lotus capitulis dimidiatis, caule fruticoso, foliis nitidis. LINN. Sp. 10. Lotus *argentea Cretica.* PLVKN. Alm. 226. T. 43. f. 1. In *Syria, Creta.*

H. Lotus capitulis hirsutis, caule erecto hirtio, leguminibus ovatis. LINN. Sp. 11. Lotus *pentaphyllor, siliquosus, hirsutus.* 1. R. H. HALL. G. 262. In *Gallia Narbonensi.*

Lotus capitulis depressis, caulibus decumbentibus, leguminibus cylindricis. LINN. Sp. 13. Lotus *floribus umbellatis, siliquis pendulis, caule erecto.* HALL. Helv. 1. In *dumetis bey der Papiermühle.*

β. Lotus *floribus umbellatis, siliquis pendulis, caule procumbente.* HALL. H. 2.

H. *Lotus capitulis subglobosis, caule erecto, leguminibus rectis glabris.* LINN. *Sp.* 12. *Lotus villosus, flore altissimo, glomerato.* I. R. H. HALL. G. 262.

H. *Lotus oligoteratos, Cretica, lutea, filiquis longis erectis,* MORIS. HALL. G. 263.

ORNITHOPUS. LINN. 790. Ornithopodium. TOURN.

H. *Ornithopus foliis pinnatis, leguminibus articulatis levibus subarcuatis.* HALL. H. 572. LINN. *Sp.* 1. *Ornithopodium maius & minus.* C. B. I. R. H. In *Helvetia.* A.

H. *Ornithopus foliis pinnatis, leguminibus compressis rugosis.* LINN. *Sp.* 2. *Ornithopodium scorpioides, filiqua compressa.* I. R. H. HALL. G. 264. In *Italia.* A.

H. *Ornithopus foliis pinnatis, leguminibus semicircularibus levissimis.* HALL. G. 264. c. *deser.*

H. *Ornithopus foliis ternatis subsessilibus appendiculatis: impari maximo.* LINN. *Sp.* 3. *Ornithopodium portulacae folio.* I. R. H.

HIPPOCREPIS. LINN. 791.

H. *Hippocrepis leguminibus sessilibus solitariis utrinque exscissis.* LINN. *Sp.* 1. HALL. *Gott.* 269. *Ferrum equinum filiqua singulari.* C. B. GARIDEL. T. 114. *Ferrum equinum vulgare.* COLUMN. *Epbr.* L. F. 300. In *Italia.* A.

H. Hip-

Hippocrepis leguminibus pedunculatis confertis, margine exteriori repandis. LINN. *Sp.* 3. Ferrum equinum *siliquis umbellatis, utrinque emarginatis.* HALL. *Helv.* 572. Ferrum equinum *capitatum.* COLUMN. *Ecphr.* I. T. 301. In apricis M. *Plesi.* P.

SCORPIURVS. LINN. 792.

H. *Scorpiurus* pedunculis unifloris, leguminibus testis undique squamis obtusis. *vermiculata.* LINN. *Sp.* 1. *Scorpioides siliqua crassa boeltii.* 1. R. H. 402. A.

H. *Scorpiurus* pedunculis bifloris, leguminibus extrorsum obtuse aculeatis. *muricata.* LINN. *Sp.* 2. Hanc refert MORIS. II. p. 127. S. II. T. 11. f. 4.

H. *Scorpiurus* pedunculis subtrifloris, leguminibus extrorsum spinis distinctis acutis. *fulcata.* LINN. *Sp.* 3.

H. *Scorpiurus* pedunculis subquadrifloris, leguminibus extrorsum spinis confertis acutis. LINN. *Sp.* 4. *Scorpioides corniculis magis asperis in se contortis & convolutis.* MORIS. II. p. 127. S. II. T. 11. f. 2.

AESCHYNOMENE. LINN. 769.

H. *Aeschynomene* caule hispido, foliis acuminatis, leguminum articulis suborbiculatis. LINN. *Sp.* 2. ROY. *Lagdb.* 384. HALL. G. 265. c. *deser.* In *Jamaica.*

HEDY.

HEDYSARVM. LINN. 793.

H. Hedysarum foliis simplicibus lanceolatis obtusis, caule fruticoso spinoso. *Albogi*. LINN. Sp. 1. In *Oriente*.

H. Hedysarum foliis simplicibus ternatisque, floribus racemosis. *Canadense*. LINN. Sp. 15. Hedysarum *triphyllum Canadense*. CORNYT. *Canad.* T. 44. P.

H. Hedysarum foliis pinnatis, leguminibus articulatis aculeatis nudis rectis, caule diffuso, LINN. Sp. 28. Hedys. *clypeatum flore suaviter rubente*. I. R. H. 401.

fl. flore albo. I. R. H. Fr.

H. Hedysarum foliis pinnatis, leguminibus articulatis aculeatis flexuosis, caule diffuso. LINN. Sp. 30. Hedysarum *annuum, siliqua aspera undulata, intorta*. I. R. H. Hedys. *siliquis undulatis*. RIV. T. 99. In *Italia*.

H. Hedysarum foliis pinnatis, leguminibus monospermis aculeatis, corollarum alis calyce brevioribus. LINN. Sp. 32. Onobrychis *spicis densioribus, siliquis asperis*. HALL. *Helv.* 577. Onobrychis *foliis viciae, fructu echinato maior*. C. B. *Esparsetti* vulgo. Copiose seritur. Sponte in apricis Germaniae, Helvetiae. P.

H. Hedysarum foliis pinnatis, leguminibus monospermis aculeatis, petalis aequalibus. LINN. Sp. 33. Onobrychis *spicis rarioribus, siliquis aculeatis*. HALL. G. 267. Onobrychis *Cret. foliis Viciae, fructu*

fructu magno, cristato & aculeato. I. R. H. In *Gallia calidiori.*

CORONILLA. LINN. 789.

H. *Coronilla frumcosa*, corollarum vnguibus calyce duplo longioribus. *Emerus*. LINN. *Sp.* 1. *Coronilla frutescens*, *siliquis longissimis obscure articulat.* HALL. *H.* 4. *Emerus* I. R. H. *Coronilla scorpioides maior.* C. B. In *Helvetia & Europa australi.* Fr.

H. *Coronilla frumcosa*, stipulis subrotundis. LINN. *Sp.* 3. *Polygala Valentina.* CLVS. 98. *Coronilla* RIV. T. 92. In *Europa australi.*

H. *Coronilla foliolis denis*, leguminibus numerosis articulatis pendulis. HALL. *G.* 268. *Helv. n.* 4. *Coronilla scorpioides altera.* CLVS. 96. *Coronilla montana.* RIV. T. 93. In *Helvetia. Austria.*

H. *Coronilla herbacea*, leguminibus falcato-glabratis. LINN. *Sp.* 5. *Securidaca.* HALL. *H.* 576. *Securidaca lutea maior.* C. B. In *Italia. Hispania. A.*

Coronilla herbacea, leguminibus erectis teretibus torosis numerosis, foliis glabris. *varia.* LINN. *Sp.* 6. *Coronilla herbacea*, *flore vario.* I. R. H. *Securidacae II. prior species.* CLVS. 237. Inter segetes. P.

H. *Coronilla herbacea* leguminibus quinis erectis teretibus articulatis. LINN. *Sp.* 7. *Coronilla Cretica herbacea*, *flore parvo purpurascente.* I. R. H.

COLUTEA. LINN. 776.

H. Colutea arborea foliolis obverse ovatis oblongis, apice emarginatis. Colutea arborea foliolis obverse cordatis. LINN. Sp. 1. HALL. H. 575. Colutea vesicaria. C. B. 1. R. H. In Helvetia. Italia. Foliola duplo longiora, ac lata.

H. Colutea fruticosa, foliolis subrotundis, apice retusis. Colutea Orientalis, flore sanguinei coloris, lutea macula notato. I. R. H. Cor. 44. Diversa a priori videtur. Foliola subrotunda, aequae lata ac longa. Vexillum duabus maculis flavis notatum. Fr.

H. Colutea fruticosa, foliolis ovato-oblongis. LINN. Sp. 2. Colutea Aethiopica, barbae Jovis folio, flore phoeniceo. I. R. H. Colutea Aeth, flore purpureo. BREYN. Cent. T. 29. P.

H. Colutea herbacea, foliis linearibus. LINN. Sp. 3. Colutea Afr. annua, vesiculis compressis, foliis parvis mucronatis & cordatis. COMM. H. II. T. 44. VOLKAM. Norib. p. 118.

ROBINIA. LINN. 775.

H. Robinia pedunculis racemosis, foliis imparipinnatis. LINN. Sp. 1. Pseudo-Acacia. RIV. T. 63. Acacia AMER. Robini. CORNUT. Canad. 171. In Virginia. Fr.

GALEGA. LINN. 770.

Galega. LINN. Spec. 715. BLACKW. T. 91. vulgaris. CIVS. CCXXXIII. I. R. H. Ruta caprarum. MATH. 739. flore albo. G. flore subcaeruleo. In Helvetia. Italia. P.

INDI.

INDIGOFERA. LINN. 794.

H. Indigofera leguminibus arcuatis incanis, racematis folio brevioribus. *tinctoria*. LINN. Sp. 1. *Amoen. Acad.* l. p. 408. Anil 1. B. II. p. 945. In *India*. Fr.

GLYCIRRHIZA. LINN. 788.

H. Glycirrhiza leguminibus glabris. LINN. Sp. 2. *Riv. T.* 64. *filiquosa* & *Germanica*. B. Pin. 312. *vulgaris*. DODON. *Pempt.* 341. Colitur prope *Bambergam*. Fr.

OROBVS. LINN. 780.

Orobus caule simplicissimo, foliolis pluribus ovatis acutis, stipulis semisagittatis. LINN. *Cyff.* 366. HALL. H. 1. *vernus*. LINN. Sp. 4. Orobus *sylvaticus purpureus vernus*. C. B. Verno tempore vbique in sylvis. A.

Orobus caule simplici, foliis imis ovatis, superioribus ellipticis acutis. HALL. H. 3. *taberosus*. LINN. Sp. 5. Orobus *sylvaticus foliis oblongis glabris*. I. R. H. *Astragalus sylvaticus*. THAL. *Herc.* 7. T. 1. In sylva *Hercynica* circa *Elbingerode*.

Orobus caule ramoso, foliis sexiugis ovato-oblongis. LINN. Sp. 7. HALL. H. 4. Orobus *sylvaticus, viciae filii*. C. B. Orobus *Pannonicus*. II. CLVS. CCXXX. Circa *Rheinhausen*.

H. Orobus fruticosus, pedunculis unifloris, foliolis plurimis subtus tomentosis. *Robinia*. LINN. Sp. 3. *Aspalathus arborescens, pinnis foliorum crebrioribus oblongis*. AMMANN. *Ruth.* 285. *Caragana Sibiria*. ROYEN. 537. Fr.

H. Orobus pedunculis uni- & bifloris, foliis quaternatis, cuneiformibus, glabris. Robinia. LINN. Sp. 4. *Aspalathus frutescens maior latifolius, cortice aureo.* AMMANN. Ruth. 283. In Sibiria. Vtraque species a *Robiniae* definitione generica maxime aliena, fabrica calycis, corollae, & legumine proxime ad *Orobum* accedit. Fr.

H. Orobus foliis coniugatis subsessilibus, stipulis dentatis. *Lathyroides*. LINN. Sp. 1. *Lathyroides erecta, folio ovato acuminato, coerulescens viciae floris & siliquis.* AMMANN. Ruth. 151. T. 7. f. 2. P.

PISUM. LINN. 779.

H. Pisum stipulis crenatis, pedunculis multifloris. LINN. Cliff. 369. *fativum*. Sp. 1. BLACKW. T. 39. *Pisum hortense majus*. C. B. 3. *Pisum nigrum*. RIV. T. 39. A.

H. Pisum petioli decurrentibus membranaceis diphyllis, pedunculis unifloris. *Ochrus*. LINN. Sp. 4. *Ochrus folio integro capreolum emittente.* C. B. In Italia.

LATHYRVS. LINN. 781.

H. Lathyrus pedunculis unifloris, cirrhis aphyllis, stipulis sagittato cordatis. LINN. Sp. 1. *Aphaca*. HALL. H. 597. LOBEL. ic. II. p. 70. In Helvetia. Gallia. A.

H. Lathyrus pedunculis unifloris, foliis simplicibus, stipulis subulatis. *Nissolia*. LINN. Sp. 2. *Lathyrus foliis singularibus*. HALL. H. 1. *Nissolia vulgaris*. I. R. H. In Gallia. A.

H. La-

g. *Lathyrus* pedunculis unisloris, cirrhis diphyllis, leguminibus ovatis compressis dorso bimarginatis. LINN. *Sp.* 5. *Lathyrus* HALL, *H.* 3. *Lathyrus sativus flore fructuque albo.* L. R. H. Seritur in *Thuringia* & alibi. A.

h. flore coerulea. *Cicerula Aegyptia.* CLV. CCXXVI.

i. flore ex albo & coeruleo vario.

j. *Lathyrus* pedunculis unisloris, calyce brevioribus, cirrhis diphyllis, foliolis lanceolatis. LINN. *Sp.* 6. *Lathyrus orientalis, flore, vix conspicuo.* P. PERH. II. p. 42. HALL. *Gott.* 289. A.

k. *Lathyrus* flore miniato, siliquis & seminibus maculatis. HALL. *G.* 289.

l. *Lathyrus* pedunculis subunisloris, cirrhis polyphyllis, foliolis alternis. *articulatus.* LINN. *Sp.* 10. *Clymenum Hispanicum, folio vario, aliqua articulata.* L. R. H. In *Baetica.* A.

m. *Lathyrus* pedunculis bifloris, cirrhis diphyllis, foliolis ensiformibus, leguminibus glabris, stipulis bipartitis. LINN. *Amoen. Acad.* III. p. 417. *Lathyrus Hispanicus flore luteo.* RIV. *T.* 44.

n. *Lathyrus* pedunculis bifloris, cirrhis diphyllis, foliis ovato-oblongis, leguminibus hirsutis. *odoratus.* LINN. *Sp.* 12. *Lathyrus distuplatyphyllos hirsutus mollis, flore odore magna.* COMMEL. II. *T.* 80. In *Sicilia.* A.

o. flore ex albo & rubro vario.

p. *Lathyrus* pedunculis bifloris, cirrhis diphyllis, foliolis lineari-lanceolatis, leguminibus hirsutis,

feminibus scabris. LINN. *Sp.* 12. HALL. *H.* 9. *Lathyrus angustifolius*, *filiqua bisfuta*. C. B. In *Helvetia. Asia. A.*

H. *Lathyrus pedunculis bifloris*, cirrhis diphyllis, foliolis alternis lanceolatis. LINN. *Sp.* 13. *Lathyrus Tingitanus*, *filiqua Orobi*, *fore amplo ruberrimo*. MORIS. *H.* p. 5. *Lathyrus pedunculis bifloris*, *cirrhis polyphyllis*, *stipulis dentatis*. LINN. *Sp.* 14. *Clymenum Hispanicum*, *fore vario*, *filiqua plana*. L. R. H.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliolis ovalibus, internodiis nudis. LINN. *Sp.* 15. HALL. *H.* 9. *Lathyrus arvensis repens tuberosus*. C. B. Terrae glandes DOD. *Hist.* 550. In acclivi M. Heymberg. A.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliolis lanceolatis, cirrhis simplicissimis. *pratensis*. LINN. *Sp.* 16. HALL. *H.* 10. *Lathyrus sylv. lucus*, *folio Viciae*. C. B. In *prat.* P.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliis ensiformibus. LINN. *Sp.* 17. *Lathyrus foliis binatis ellipticis*, *floribus racemosis*. HALL. *H.* 6. *Gott.* p. 290. *Lathyrus sylvestris*. C. B. CLVS. CXXIX. In *sylva ultra Geismar*.

Lathyrus pedunculis multifloris, cirrhis diphyllis, foliolis lanceolatis, internodiis membranaceis. LINN. *Sp.* 18. HALL. *H.* 8. *Lathyrus latifolius*. C. B.

β. Varietas *angustifolia*. *Lathyrus Narbonensis maior angustifolius*. C. B. Circa *Blankenburg*.

H. Lathyrus pedunculis multifloris, cirrhis polyphyllis, stipulis lanceolatis. *palustris*. LINN. *Sp.* 20. Lathyrus *foliis senis linearibus*. HALL. *G.* 288. Clymenum *Parifense flore coeruleo*. I. R. H. In *Thuringia*. P.

H. Lathyrus paribus foliorum quinque, floribus racemosis, siliquis incurvis. HALL. *Comm. Gott.* T. II. 1792. p. 341. An Lathyrus *pedunculis multifloris, cirrhis polyphyllis, stipulis ovatis, basi acutis*. LINN. *Sp.* 21. H. *Vpf.* 217.?

VICIA. LINN. 782.

H. Vicia foliis quaternis, ovatis, floribus solitariis petiolatis. HALL. *Gott.* 291. Lathyrus. LINN. 9. Clymenum *Bithynicum, siliqua singulari, flore minore*. I. R. H. Habitus Lathyri, flos viciae.

Vicia pedunculis multifloris, foliolis octonis ovatis amplissimis. LINN. *Sp.* 1. HALL. *H.* 3. *Psidium sylvestre*. CLVS. CXXIX. Cracca *flore ochroleuco*. RIV. T. 52. An der *Längelerburg* versus *Makenrade*. A.

Vicia pedunculis multifloris, foliolis reflexis ovato mucronatis, stipulis subdentatis. LINN. *Sp.* 2. HALL. *H.* 5. Vicia *maxima dumetorum*. C. B. In dumetis ad prata prope *Herperhausen*, in *Eulenthal*.

Vicia foliolis ovalibus, stipulis denticulatis, siliquis racematim pendulis. HALL. *H.* 6. T. XIV. LINN. *Sp.* 3. Viciae *pulchrum genus, multifolium*. I. B. II. p. 316. *lisdem locis cum priori*.

H. *Vicia pedunculis multifloris*, foliolis ovatis, stipulis maximis folia aequantibus. *Folia octona, sena; ovato-elliptica, glabra. Stipulae maximae, crenulatae, acutae. Petioli duodecim flori & ultra. Calyx coerulescens. Vexillum venosum. Legumina longa, angusta, glabra.*

Vicia foliolis linearibus subtus sericeis, siliquis glabris, dense spicatis, pendulis, stipulis integris. HALL. H. 8. LINN. Sp. 5. *Vicia multiflora*. C. B. In pratis vdis.

H. *Vicia pedunculis subquinquefloris*, foliolis integerrimis, stipulis integris, leguminibus erectiusculis. LINN. Sp. 9. *Vicia Benghalensis hirsuta & incana; siliquis Pisi.* HERM. Lagdb. T. 615. A.

Vicia leguminibus sessilibus subbinatis erectis, foliis retusis, stipulis notatis. LINN. Sp. 10. HALL. H. 1. *Vicia vulgaris sativa*. C. B. 1. R. H. BLACKW. T. 423. Inter segetes; & copiose colitur.

Vicia leguminibus sessilibus erectis, foliis imis ovatis, superioribus linearibus. HALL. H. 3. *Vicia vicia angustiore folio*. C. B. 1. R. H. *angustifolia*. RIV. T. 55. Inter segetes circa Lenglern & M. Haven.

H. *Vicia* leguminibus sessilibus reflexis pilosis foliariis pentaspermis, corollae vexillis glabris. LINN. Sp. 12. *Vicia sylvestris lutea, siliqua hirsuta*. L. B. II. p. 313. HALL. G. 292.

H. *Vicia* leguminibus sessilibus reflexis pilosis polyspermis, corollae vexillis pilosis. An *Vicia*.

LINN.

LINN. Sp. 13. *Vicia fylv. lutea cum galea fusca*.
I. B. II. p. 314. HALL. G. l. c.

β. *Vicia orient. carina & alis albis, vexillo reflexo subfusco*. DILL. HALL. G. l. c.

h. *Vicia leguminibus subsessilibus solitariis pendulis glabris tetraspermis, foliolis linearibus emarginatis*. LINN. Sp. 14. *Vicia angustifolia purpureo-violacea, siliqua lata glabra*. MAGNOL. Bot. Monsp. 276. HALL. G. 293. *Semina sphaerica*.

h. *Vicia leguminibus sessilibus solitariis glabra nodosis, seminibus lentiformibus*. HALL. G. 293.

Vicia leguminibus subquaternis erectis, foliis ex ovatis obtuse decrescentibus. HALL. H. 4. LINN. Sp. 15. *Vicia sepium*. RIV. T. 56. *folia rotundiora acuta*. C. B. I. R. H. Ad sepes & in sylvis.

h. *Vicia leguminibus subsessilibus solitariis erectis, foliolis senis subovatis, stipulis denticulatis*. LINN. Sp. 16. *Vicia Narbonensis*. RIV. T. 57. HALL. G. 291. *Bona sylvestris*. DOD. Peimpt. 516. *Folia primo bina, deinde quaterna, suprema sena*. In Europa australi.

Vicia foliis linearibus obrusis, siliquis gemellis polyspermis. HALL. H. 9. *Ervum* LINN. 2. *Vicia minima cum siliquis glabris*. I. R. H. In agris. A. I

h. *Vicia floribus solitariis, seminibus compressis*. HALL. G. 296. *Ervum*. LINN. 4. *Lens monanthor*. HERM. Lugdb. 360. A.

FABA. TOVRN. 212.

h. *Faba*. C. B. I. R. H. *Faba maior*. BLACKW. T. 19. *Vicia caule erecto, petiolis absque cirrbis*. LINN.

Sp. 17. Varietates ex diversitate floris & feminis. Copiose feritur.

CICER. LINN. 783.

Cicer foliis serratis. LINN. Sp. 1. Cicer *arietinum*. DOD. *Pempt.* 525. *sativum*. C. B. Inter segetes *Hispaniae*.

ERVUM. LINN. 784. Lens TOURN.

H. Ervum feminibus compressis convexis. LINN. Sp. 1. Lens. HALL. H. 601. RIV. T. 35. Lens *vulgaris*. C. B. I. R. H.

β. Lens *maior*. RIV. T. 35. I. R. H. diversa non videtur. A.

Ervum pedunculis multifloris, feminibus globosis. *hirsutum*. LINN. Sp. 3. Vicia *siliquis hirsutis, pendulis, dispermis*. HALL. H. 10. Vicia *segetum cum siliquis plarimis hirsutis*. C. B. In agris. A.

H. Ervum foliis linearibus retusis octonis, pedunculis bifloris. Vicia *multiflora cum siliquis lotis*. HALL. G. 296. In *Pensylvania*. A.

H. Ervum leguminibus undato-plicatis. LINN. Sp. 5. Ervum *leguminibus pendulis*. HALL. H. 601. Ervum *verum*. I. R. H. In hortis rarius feritur. A.

H. Ervum caule, & foliolis obverse ovatis, hirsutis, petiolis mucronatis, leguminibus solitariis polyspermis erectis sessilibus. Circa *Berolinum*. A Vicia *minima*. RIV. RUPP. I. R. H. diversa videtur, quod pedunculi neque foliis longiores, neque cirrhosi sint, quales huic tribuit LINNAEVS *Cent. Pl. II. n. 185*.

ONONIS. LINN. 772. Anonis. HALL.

Ononis floribus racemosis geminatis, foliis ternatis, ramis spinosis. LINN. Sp. 1. Anonis floribus pene sessilibus, vaginis foliorum amplissimis HALL. H. 4. Anonis, resta bovis. TABERN. p. 528. Anonis spinosa, flore purpurea. C. B. L. R. H. BLACKW. T. 391. Ad vias. Fr.

β. flore albo. In colle prope Bovenden.

γ. spinis destituta iunior.

H. Ononis spicis foliosis, foliis simplicibus obtusis. LINN. Sp. 5. Anonis alopecuroides Sicula. I. R. H. HALL. G. 284. In Europa australiori. A.

H. Ononis pedunculis unisporis, filo terminatis, foliis ternatis. LINN. Sp. 7. Anonis pedunculis longis unisporis, leguminibus pendulis, teretibus hirsutis. HALL. H. 6. Anonis spinis carens, lutea maior. I. R. H. NATRIX RIV. T. 68.

H. Ononis pedunculis bifloris filo terminatis. LINN. Sp. 9. Anonis pedunculis longis, leguminibus pendulis articulatis compressis, HALL. G. 284. Anonis filiquis Ornithopodii. I. R. H. In Sicilia.

SPARTIVM. LINN. 765.

H. Spartium inerme, ramis oppositis teretibus, rariter foliosis, apice floriferis. iuncum. LINN. Sp. 1. Spartium Dioscoridis. CLVS. 101. Genista transalpina. CAMER. H. 12. XIV. Genista Hispanica. RIV. T. 66. In Provincia Hispania. Fr.

H. Spartium foliis ternatis solitariisque, ramis inermibus angulatis. Spartium scoparium. LINN. Sp.

7. Genista, TABERN. 1509. RIVIN. T. 65. Genista *angulosa* & *separia*. C. B. In arenosis Marchiae & circa Noribergam. Fr.

H. Spartium foliis solitariis ternatisque, ramis sexangularibus apice floriferis. LINN. Sp. 5. Non dum floruit, sed ex angulis tribus, per ramos a singulo folio decurrentibus, facile cognoscitur. Fr.

GENISTA. LINN. 766. HALL. 592.

H. Genista ramulis striatis & foliis ternatis villosis. *Canariensis*. LINN. Sp. 1. Cytisus minoribus foliis, ramulis tenellis villosis. B. Pin. 390. Cytisus L. CLVS. 94. Cytisus *Canariensis* *sempervirens* & *incanus*. COMMEL. Hall. T. 52. Fr.

H. Genista foliorum parte prima cauli adnata, elliptica, altera patente lanceolata. HALL. H. 4. Genista ramis ancipitibus articulatis, foliis ovato-lanceolatis. LINN. Sp. 2. Chamaegenista altera. CLVS. 104. Genista *sagittalis* *Pannonica*. CAMER. Hist. XIII. In austriaci Germanica. Fr.

Genista foliis confertis lanceolatis, ramis striatis erectis, floribus sessilibus spicatis. HALL. H. 1. LINN. Sp. 4. Genista *tinctoria* *Germanica*. C. B. 395. vulgaris. CLVS. 101. In sylvis & ad oras agrorum. Fr.

Genista foliis lanceolato-obtusis villosis, caule tuberculato decumbente. *pilosa*. LINN. Sp. 5. HALL. H. 2. Genista *ramosa* *foliis Hyperici*. C. B. 1. R. H. Chamaegenista *prima*. CLVS. 103. In sylva *bercynica* circa M. Bructerum. Fr.

H. Ge-

H. Genista spinis simplicibus, in ramis florigeris, foliis levibus lanceolatis solitariis. HALL. G. 186. LINN. Sp. 6. Genista minor asphaltoides. C. B. Prodr. 157. In arenosis Cellensibus.

Genista foliis ovato lanceolatis, hirsutis, spinis ramosis in ramis senescentibus. HALL. H. 3. LINN. Sp. 7. Genista spinosa minor Germanica. C. B. In Sylva ultra Rhoenhausen. Fr.

ARACHIS. LINN. 787.

Arachis LINN. Sp. 1. Arachidnoides Americana. NISSEOLE Aff. Par. 1727. HALL. G. 287.

CYTISVS. LINN. 785.

Cytisus racemis simplicibus pendulis, foliolis ovato-oblongis. Laburnum. LINN. Sp. 1. HALL. H. 1. p. 591. Laburnum. RIV. T. 62. Circa arcem Grubenhagen. Fr.

CROTALARIA. LINN. 771.

H. Crotalaria foliis ternatis ovatis, stipulis nullis, leguminibus hirsutis. incana. LINN. Sp. 11. Crotalaria foliis ternatis, petiolis nudis. CLIFF. 358. c. descr.

ERYTHRINA. LINN. 762.

H. Erythrina foliis ternatis, caule arboreo aculeato. Corallodendron. LINN. Sp. 2. Cliff. 354. Coral arbor Americana. COMMEL. H. I. T. 108. In America. Fr.

LUPINVS. LINN. 774.

H. *Lupinus calycibus alternis inappendiculatis*, labio superiore integro, inferiore tridentato. LINN. *Sp.* 2. *Lupinus flore albo.* CLVS. CCXXIX. RIV. T. 25. A.

H. *Lupinus calycibus semiverticillatis*, labio superiore bifido, inferiore subtridentato. *varius.* LINN. *Sp.* 3. *Lupinus sylv. flore purpureo, semine rotundo vario.* I. B. II. p. 291. *Lupinus maior, flore purpureo.* HALL. G. 270. A.

H. *Lupinus calycibus verticillatis appendiculatis*, labio superiore & inferiore integris. *hirsutus.* LINN. *Sp.* 4. *Lupinus villosus coeruleus maior.* B. *Prodr.* 148. *Lup. exoticus hirsutissimus.* I. B. *Hist.* II. p. 280. *flore coeruleo.* RIV. T. 27. A.

H. *Lupinus calycibus alternis appendiculatis*, labio superiore bipartito, inferiore integro. *angustifolius.* LINN. *Sp.* 6. *Lupinus flore coeruleo minore.* RIV. T. 28. *Indicus* CORNV. p. 207. A.

H. *Lupinus calycibus verticillatis appendiculatis*, labio superiore bipartito, inferiore tridentato. *luteus.* LINN. *Sp.* 7. *Lupinus sylv. flore luteo.* C. p. *Pin.* 348. RIVIN. T. 26. A.

PHASEOLVS. LINN. 777.

H. *Phaseolus caule volubili*, floribus racemosis geminis, bracteis calyce maioribus, leguminibus pendulis. LINN. *Sp.* 1. *Phaseolus vulgaris.* LOB. *sc.* 59. *Smilax bortenfis* f. *Phaseolus maior.* A. *Pin.* 339. *Mire ludit colore & mole seminis.*

H. Phaseolus caule volubili, pedunculis subquadrisfloris, bracteis calyce minoribus, leguminibus sublunatis superne granulatis. Phaseolus HALL. G. 271. n. 2. *Vexillum viride. Alae & carina albae, exsiccatae flavescences.* An Phaseolus caule volubili, leguminibus acinaciformibus sublunatis laevibus. LINN. Sp. 3. ? Videtur.

H. Phaseolus caule volubili, floribus racemosis geminis, bracteis calyce brevioribus, leguminibus pendulis (scabris). LINN. Sp. 2. Phaseolus puniceo flore. CORNVT. Canad. 184. *Flos priori triplo maior, semper coccineus. Legumina scabra, nodosa.*

H. Phaseolus caule volubili, petiolis hirsutis alatis, bracteis calyce maioribus, corolla concolore. *Caulis scaber. Pedunculus communis hirsutus, partiales laeves. Petala omnia flava, bracteae cum calyce ex viridi flavescences.* An Phaseolus caule volubili, florum vexillo revoluti, calycibus concolore. LINN. Sp. 4. *Cliff. 359. **?

H. Phaseolus caule volubili, vexillis carinaque spirahiter convolutis. *Caracalla.* LINN. Sp. 9. Phaseolus Amer. flore cochleato, odorato, vulgo Caracalla. HALL. Gott. 271. Phaseolus Indicus, flore cochleato, TRIUMPH. Obs. T. 94.

DIAD. DOLICHOS. LINN. 778.

H. Dolichos siliquis latissimis, superne verrucosis. HALL. G. 272. Dolichos leguminibus ovato-acinaciformibus, seminibus ovatis hilo arcuato versus alteram extremitatem. LINN. Sp. 1. Phaseolus Aegyptius

Aegyptius, nigro semine. 1. R. H. *Phaseolus niger* Lablab. ALPIN. *Aeg. T.* 75. A.

H. *Dolichos* foliis glabris, leguminibus angustis, erectis rigentibus. HALL. G. 272. *Phaseolus orthocarpus.* RIVIN. T. 32. *Vexillum extus subluteum, intus violaceum. Alae violaceae.*

H. *Dolichos* foliis glabris, intermedio subhastato, floribus gemellis, leguminibus cylindraceis, falcatis, deorsum recurvis. *Folia amplissima, lateralia extrorsum maxime gibba, angulo interiori plane destituta: intermedium utrinque gibbum, sinem versus acuminatum. Stipulae lanceolatae. Vexillum amplum extus ex luteo album intus obsolete purpureum, in origine duabus maculis flavis notatum. Alae inflatae purpureae. Carina alba. Semina violacea.* A.

H. *Dolichos* caule volubili perenni, foliis tomentosis, mollissimis, intermedio rhomboideo. Nondum floruit; ex habitu huc retuli. Fr.

H. *Dolichos* caule perenni, pedunculis capitatis, leguminibus strictis linearibus. LINN. *Sp.* 9. *Dolichos caule perenni lignoso.* H. Cliff. 360. T. 10. HALL. G. 273. Fr.

H. *Dolichos* volubilis annuus. HALL. G. 273.

TRIGONELLA. LINN. 804. *Foenum graecum.* TOURN.

H. *Trigonella* leguminibus pedunculatis congestis pendulis ovalibus compressis, caule diffusio, foliis subrotundis. LINN. *Sp.* 2. *Melilotus supina latifolia,*

trifolia, filiqua lata membranacea compressa. HALL. *Comm. Gott. T. I. p. 213. T. XIII. Act. Petrop. 8. p. 209. T. 12. In Sibiria. A.*

H. Trigonella leguminibus sessilibus subfoliariis erectis ensiformibus. LINN. *Sp. 6. Foenum graecum filiquis ensiformibus.* HALL. *G. 274. Foenum graecum sativum. C. B.*

H. Trigonella leguminibus paucioribus congestis erectis, linearibus, obtuse flexis. LINN. *Spec. 35. Foenum graecum filiquis binis angustis, obtuse flexis.* HALL. *G. 1. 6. A.*

H. Trigonella (*Foenum graecum*) leguminibus pedunculatis quinis congestis declinatis subfalcatis compressis. HALL. *H. l. c. spinosa.* LINN. *Sp. 4. Foenum graecum sylv. polyceratum creticum maius.* BREYN. *Cent. T. 33. f. 1. A.*

H. Trigonella (*Foenum graecum*) leguminibus pedunculatis numerosis sessilibus declinatis subfalcatis glabris. HALL. *l. c.* LINN. *Sp. 5. Foenum graecum sylv. minus polyceratum Monspeliense.* BREYN. *Cent. T. 33 f. 2. A.*

H. Trigonella (*Foenum graecum*) leguminibus spicatis numerosis falcatis declinatis subhirsutis. HALL. *l. c.* Trifolium. LINN. *Sp. 7. Trifolium corniculatum.* IL. DOD. *Pempt. 573. In Italia. A.*

MEDICAGO. LINN. 895.

H. Medicago leguminibus lunatis margine integerrimis, caule arboreo. LINN. *Sp. 1. Medicago trifolia frutescens, incana.* I. R. H. HALL. *G. 275.*

Cyti-

Cytisus Marantae. LOB. II. p. 46. MARTYN. *Georg. Virgil.* p. 232. qui pro vero *Cytiso* veterum habet. In regno *Neapolitano*. Fr.

H. *Medicago* foliis pinnatis, pinna extrema maxima, leguminibus reniformibus margine dentatis. LINN. *Sp.* 4. *Medicago Vulnerariae facie Hispanica*. I. R. H. In *Hispania*. A.

H. *Medicago* leguminibus reniformibus margine dentatis, foliis ternatis. LINN. *Sp.* 3. *Medica filiqua reniformi, ciliata, amplissima*. I. R. H. *Lunaria radiata Itatorum*. LOB. II. p. 38. In *Italia*. A.

H. *Medicago* pedunculis racemosis, leguminibus lunatis, caule prostrato. LINN. *Sp.* 6. *Medica* HALL. H. 3. *Falcata* RIVIN. T. 84. *Medica flavo flore*. CLVS. II. CCXXXIII. In prat. A.

H. *Medicago* spicis ovalibus, leguminibus reniformibus monospermis, caule procumbente. *lapulina*. LINN. *Sp.* 7. *Medica* HALL. H. 4. *Trifolium prat. luteum, capitulo brevior*. C. B. *Pin.* 328. *Melilotus minima*. RIV. T. 8. In prat. A.

H. *Medicago* caule erecto, pedunculis racemosis, leguminibus contortis. LINN. *Sp.* 5. *Medica* HALL. H. 2. *Medica sativa vulgo Luzerne*. In *Germania australiori & Helvetia*. P.

H. *Medicago* pedunculis racemosis, leguminibus cochleatis spinosis, caule procumbente tomentoso. LINN. *Sp.* 8. *Medica marina*. CLVS. CCXXXIII.

H. *Medicago* leguminibus cochleatis, stipulis dentatis, foliis serratis, caule diffuso. LINN. *Sp.* 9. Ad hanc refert LINNAEVS tanquam varietates omnes re-

reliquas cochleatis, quae ad varias species reduci posse videntur.

1. Medicago leguminibus solitariis cochleatis compressis planis, stipulis ciliatis, caule diffuso. SAUVAG. *Medica fructu orbiculato, compresso, plano.* RIV. HALL. 275.

2. Medicago leguminibus solitariis cochleatis compressis umbilicatis. *Medica fructu scutellato.* HALL. H. 276.

3. Medicago fructu tornatili. RIV. HALL. l. c.

4. Medicago fructu turbinato. RIV. β. *Medica fructibus rugosis, compressis, floribus racemosis.* HALL. l. c. γ. *Medica dicarpos, fructu turbinato, rugoso.* MORIS II. p. 153.

5. Medicago fructu coronulam vtrinque dentatam referente. SAUVAG. *Medica coronata.* HALL. l. c.

6. Medicago fructu echinato. *Medica fructu turbinato, breviori, brevissime echinato.* HALL. l. c. 277.

β. *Medica fructu polygono.* HALL. 278.

γ. *Medica turbinata echinata.* RIV.

δ. *Medica fructu turbinato, echinato, rigidiori.* HALL. l. c.

ε. *Medica fructu longius echinato.* RIV.

7. Medicago leguminibus spinosis. *Medica spinosa, fructu conico truncato, spinis distinctis.* HALL. 279.

β. *Medica fructu ovato, angustiori, hirsuto, spinis se decussantibus.* HALL. l. c.

γ. *Medica*

7. *Medica cochlearia spinosa maior, capsulae spinis longioribus sursum & deorsum tendentibus.* MORIS. I. p. 153. S. II. T. 15. f. 8. 9.

8. *Medicago fructu spinoso, globoso, foliis oblongis, argute dentatis.* HALL. G. 278. *Trifolium cochlearium spinosum Syriacum, foliis laciniatis.* BREYN. Cent. T. 34.

MELILOTVS. HALL. 587. *Trifolii*
Sp. LINN.

H. *Melilotus (Trifolium)* spicis oblongis, leguminibus feminudis mucronatis, caule erecto. LINN. Sp. 1. *Melilotus maior Jacqua violacea.* HALL. G. 281. *Lotus bortenensis odorata.* B. Pln. 330. *Lotus urbana.* BLACKW. T. 284. In Bohemia. A.

H. *Melilotus* floribus minimis dense spicatis, leguminibus monospermis rugosis, caule erecto. *Melilotus lutea minor, floribus & siliculis minoribus, spicatis & dense dispositis.* MORIS. II. p. 167. A.

Melilotus (Trifolium) leguminibus racemosis nudis, rugosis, dispermis, caule erecto. LINN. Sp. 4. HALL. H. 1. *Melilotus officinarum.* C. B. 1. R. H. BLACKW. T. 80. flore flava & albo. Vbique inter segetes.

H. *Melilotus* foliis hirsutis rariter dentatis, scapo paucifloro. HALL. Comm. Gott. 1752. T. II. p. A.

H. *Melilotus (Trifolium)* leguminibus racemosis nudis dispermis rugosis obtusis, caule erecto, foliolis integris. LINN. Sp. 5. *Melilotus Italica folliculis rotundis.* C. B. *Melilotus Italica.* CAMER. Hort. ic. 29. In Italia.

H. Me-

H. Melilotus (*Trifolium*) leguminibus racemosis nudis dispermis membranaceis, caule erectiusculo. LINN. *Sp.* 6. Melilotus *Cretica*, fructu orbiculari plano, maximo. 1. R. H. *Trifolium peltatum Creticum*. C. B. *Prodr.* T. 142.

TRIFOLIUM. LINN. 802.

Trifolium caule procumbente, & florigeris & foliis petiolo longissimis. HALL. *Helv.* 5. *Trifolium capitulis umbellaribus, leguminibus tetraspermis, caule repente.* LINN. *Sp.* 12. *Trifol. prat. flore albo.* 1. R. H. In pratis.

H. *Trifolium capitulis villosis quinquefloris, involucri centrali reflexo rigido fructum involvente, subterraneum.* LINN. *Sp.* 15. *Trifolium album trilocum subterraneum gastonium reticulatum.* MORIS. II. p. 132. S. II. T. 14. f. 5. In Gallia. A.

H. *Trifolium spicis villosis longis, corollis monopetalis, caule erecto, foliis serrulatis.* LINN. *Sp.* 17. *Trifolium spicis villosis, calyce glabro, dentibus villosis, imo longissimo.* HALL. H. 11. *Trifol. mont. spica longissima rubente.* C. B. 328. In Thuringia.

Trifolium spicis subvillosis, cinctis stipulis oppositis membranaceis, corollis monopetalis. LINN. *Sp.* 18. *Trifol. spicis villosis foliis insidentibus, vaginarum caudis capillaceis.* HALL. H. 14. *Trifol. prat. purpureum.* C. B. In pratis, & ob pabulum feritur. P.

Trifolium spicis villosis, foliis insidentibus, vaginarum caudis laticioribus. HALL. H. 13. *Trifol.*

mont. purpureum. C. B. Hanc LINNAEVS cum priori iungit.

H. Trifolium spicis villosis oblongis obtusis aphyllis, calyce tomentoso, foliolis subrotundis. LINN. *Sp.* 21. Trifol. *spica f. brotunda rubra*. C. B. Lagopus *maximus*. CLVS. CCXLVI.

Trifolium spicis villosis conico-oblongis, denticulis calycinis setaceis subaequalibus, foliolis linearibus. LINN. *Sp.* 23. HALL. *H.* 9. Trifolium *arvense humile spicatum f. Lagopus*. C. B. *Pin.* 328. In agris.

H. Trifolium spicis ovatis pilosis. denticulis calycinis lanceolatis patulis. LINN. *Cliff.* 375. *. *stellatum*. LINN. *Sp.* 24. Trifolium *stellatum*. C. B. *Pin.* 329. *Prodr.* 143. In Europa calidiori. A.

H. Trifolium spicis ovatis, calycibus patulis: lacinia infima maxima lanceolata, foliis ovatis. LINN. *Sp.* 25. Trifol. *clypeatum argenteum*. ALPIN. *Exot. T.* 306. In Oriente. A.

Trifolium caule erecto, spicis albis brevibusque terminato, foliis nervosis, inferne subhirsutis lanceolatis ferratis. HALL. *H.* 8. LINN. *Sp.* 29. Trifolium *mont. album*. C. B. I. R. H. In pratis siccis. v. g. prope *Hessendreisck*.

H. Trifolium spicis ovatis, calycibus inflatis glabris quinquedentatis, involucris univcrsalibus pentaphyllis. LINN. *Sp.* 32. HALL. *G.* 282.

H. Trifolium spicis subovatis, calycibus inflatis dorso gibbis, caulibus prostratis. LINN. *Sp.* 33. Trifol. *pratense folliculatum*. I. R. H.

Tri-

Trifolium capitulis subrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus. LINN.

Sp. 35. Trifol. *fragiferum purpureum folio oblongo.*

I. R. H. *fragiferum Friscum.* C. B. CLVS. *Gur. post.*

T. 39. In pratis *gegen die Maschmühle.*

Trifolium spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus nudis, caule erecto.

LINN. *Spec.* 36. HALL. *H.* 1. Trifolium *lupulinum alterum minus* RAH. In pratis.

Trifolium spicis ovalibus imbricatis, vexillis deflexis persistentibus, calycibus pilosis, caule erecto.

LINN. *Sp.* 37. HALL. *H.* 3. Trifol. *montanum lupulinum.* C. B. *Prodr.* 140. In sylvis.

Trifolium spicis ovalibus imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. LINN.

Sp. 38. Trifol. *pratense luteum, capitulo lupuli s. agrarium.* C. B. I. R. H. In ficcis.

H. Trifolium foliis quinis ciliatis sessilibus, leguminibus polyspermis. LINN. *Sp.* 9. *Lupinaster foliis purpureis & albis, siliquis minoribus.* AMMANN. *Rutb.* 143. 144. In *Sibiria.*

PAPILIONACEIS ADFINES.

Cum *Papilionaceis* habitu, similitudine quadam floris, & fructu conveniunt, vario tamen modo corolla, staminibus, & situ partium ab illis differunt.

AMORPHA. LINN. 768.

Vexillum solum adest, alae & carina deficiunt.

H. *Amorpha*. LINN. *Sp.* p. 713. *H. Cliff.* 353.
 T. 19. *Barba Jovis Amer. pseudo-acaciae foliis,*
flosculis purpureis minimis. *H. Angl.* In *Carolina.*

SOPHORA. LINN. 456.

Stamina decem distincta.

H. *Sophora* foliis pinnatis; foliolis numerosis
 villosis oblongis. LINN. *Sp.* 1. *Ervum Orient. al-*
pecuroides perenne, fructu longissimo. DILL. *Elth.*
 136. T. 112.

CERCIS. LINN. 458.

Corolla pentapetala florem papilionaceum refe-
 rens. Stamina distincta.

H. *Cercis* foliis cordato-orbiculatis glabris. *Sili-*
quastrum. LINN. *Spec.* 1. *Siliqua sylv. rotundifo-*
lia. C.B. *Arbor Judae.* DODON. *Pempt.* 786. In
Europa australiori; Palaestina; Aegypto.

CASSIA. LINN. 461.

H. *Cassia* foliolis trijugatis quadrijugatisque sub-
 ovatis. LINN. *Sp.* 8. *Senna Italica foliis obtusi-*
MORIS. II. p. 200. S. II. T. 24. f. 2. In *Aegypto.* Fr.

H. *Cassia* foliolis quinquejugatis ovato-lanceola-
 tis margine scabris, exterioribus maioribus, glandula
 baseos petiolorum. LINN. *Sp.* 10. Fr.

H. *Cassia* foliolis quinquejugatis ovato-lanceola-
 tis glabris, glandula baseos petiolorum. LINN. *Sp.*
 11. ROYEN. *Lugd.* 468. HALL. *Gott.* 299. C
descr. Fr.

H. *Cassia*

H. *Callia* foliolis multiungatis, calycibus acutis, floribus pentandris. LINN. *Sp.* 25. *H. Cliff.* 497. *T.* 36. HALL. *G. l. c. c. descr.* *Senna spuria Virginiana*, *Mimosae foliis, floribus parvis nictitantibus*. PLVKN. *T.* 314. *f.* 3. Fr.

MIMOSA. LINN. 597.

H. *Mimosa* foliis subdigitatis pinnatis, caule aculeato hispido. LINN. *Sp.* 14. *Mimosa spinosa* III. *siliquis parvis echinatis*. BREYN. *Cent.* p. 40. *T.* 18. Fr.

H. *Mimosa* inermis, foliis bipinnatis, spicis cernuis, floribus decandris, inferioribus castratis apetalis, siliquis linearibus glabris. LINN. *Sp.* 17. *Mimosa spuria de Fernambuke, dicta Mimosa Italica*. ZANON. *T.* 123. Fr.

H. *Mimosa* floribus superioribus pentapetalis decandris, inferioribus decapetalis sterilibus. HALL. *Gott.* 300. *c. descr.* Fr.

H. *Mimosa* spinis geminis distinctis, foliis duplicato-pinnatis, partialibus vtrinque quinopluribus. LINN. *Cliff.* 209. *Mimosa*. LINN. *Sp.* 29. *Acacia ALP. Aegypt.* *T.* 9. Fr.

H. *Mimosa* spinis geminis distinctis, foliis bipinnatis, partialibus octoingatis. LINN. *Sp.* 30. *Acacia Indica Farnesiana*. ALDIN. *Farnes.* 2. Fr.

PARKINSONIA. LINN. 460.

H. *Parkinsonia* LINN. *Sp.* p. 375. *H. Cliff.* *T.* 13. *p.* 157. *Parkinsonia aculeata, foliis Mimosae uni costae affixis*. WALTH. 6. *T.* 13. *In America*. Fr.

GLEDITSIA. LINN. 1025.

H. *Gleditsia triacanthos*. LINN. *Sp.* p. 1016.
H. Vpf. 298. GRONOV. *Virg.* 183. DV. HAMEL
Tr. des arbres. T. I. p. 265. *sqq.* T. 105. Fr.

FUMARIA. LINN. 760.

Vexillum calcaratum. Filamenta duo lateralia,
 singula tribus antheris.

Fumaria caule simplici bifolio, radice cava, foliis inter flores integris. HALL. *H.* p. 604. *Fumaria bulbosa, radice cava, maior.* C.B. Vbique verno tempore in dumetis.

Fumaria caule simplici multifolio, radice bulbosa solida, foliis inter flores palmatis. HALL. *H.* 605. *Fumaria bulbosa, radice non cava, maior.* C.B. Rarius in hortis. Circa Norbergam. *Fumaria caule simplici, bracteis longitudine florum.* LINN. *Sp.* 3. diversissimas hasce species iungit.

H. *Fumaria* siliquis linearibus pauciculis polyspermis, caule erecto. LINN. *Sp.* 5. Capricoides. *Fumaria siliquosa Canad. flore carnea & mucosa* BARREL. *ic.* 108. A.

H. *Fumaria* siliquis linearibus tetragonis, caulibus diffusis acutangulis, radice perenni. LINN. *Sp.* 9. *Fumaria calyce donata, siliquis oblongis latiusculis* HALL. *G.* 302. c. *descr. floris.* *Fumaria lutea.* C.B. *Pseudofumaria flore luteo.* RIVIN. *Tetrap. irreg.* T. 74. In Italia. P.

Fumaria pericarpis monospermis racemosis, caule diffuso, foliis subrotundis. LINN. *Sp.* 7. HALL.

H. 3. Fumaria offic. & Dioscoridi. C. B. BLACKW.
T. 237. In agris & hortis.

H. *Fumaria pericarpis monospermis spicatis*,
caule erecto, foliolis capillaribus. LINN. *Sp.* HALL.
G. l. c. *Fumaria minor, tenuifolia.* C. B. Capnos
tenuifolia. CLVS. CCVIII. *Fumaria, tenuifolia ere-*
cta Hisp. purpurea. BARREL. ic. 41. β . flore albo,
roseo. In Europa australi.

H. *Fumaria pericarpis monospermis, caulibus se-*
convolventibus. HALL. *G.* 303. LINN. *Sp.* 9. *Fu-*
maria viticulis & capreolis plantis vicinis adhae-
rens. C. B. *Folia extrema subcirrhosa.*

H. *Fumaria siliquis ovatis inflatis polyspermis,*
foliis cirrhiferis. LINN. *Sp.* 11. *Cysticaphos Africa*
scandens. BOERH. *Lugdb. I. T.* 310. In *Acthū-*
pia. A.

H. *Fumaria floribus albis.* HALL. *G.* 301.

POLYGALA. LINN. 761.

Carina sine anteriori appendiculata. Stamina
octo connata.

Polygala floribus cristatis racemosis, caulibus her-
baceis simplicibus procumbentibus, foliis lineari-
lanceolatis. LINN. *Sp.* 2. α . *Polygala foliis omnibus*
acutis. HALL. *H.* 606. n. 1. *vulgaris maior.* C. B.
CLVS. 324. In pratis & sylvis.

β . *Polygala foliis imis subrotundis, superioribus*
angustis acutis. HALL. *Helv.* 2. *Polygala minor.*
VAILL. In locis aridis apricis. Varietas prioris.

H. *Polygala floribus imberbibus sparsis* : *carina* apice subrotundo, caule fruticoso, foliis lanceolatis. LINN. *Sp.* 11. *Polygaloides* HALL. *H.* 607. *Chamaebuxus flore Coluteae*. C. B. *Anonymos flore Coluteae*. CLVS. *L. p.* 105. Circa *Noribergam*.

AGGREGATAE. S. DIPSACEAE.

In plurimis flores in capitulum congesti, & calyce communi excepti, in nonnullis flores tantum paniculati, singuli squamis distincti, & in multis receptaculo communi innati. In omnibus, excepti Globularia, femini vnico innascitur flos in plurimis monopetalus, in paucioribus pentapetalus. Stamina quatuor aut quinque corollae affixa, antheris distinctis. Tuba vnica.

STATICE. LINN. 348. Limonium. T.

Statice caule nudo simplici capitato. *Armeria*, LINN. *Sp.* 1. Statice. RVPP. 96. *Limonium aphyllacaulon gramineum globosum*. MORIS. III. S. XV. T. 1. f. 29. *Caryophyllus mont. maior, flore globoso*. C. B. 211. *flore rubro & albo*. In sabulosis Germaniae. v. g. circa *Herzberg*. &c.

β. Statice *montana minor*. I. R. H. *Caulis humilis foliis aequalis. Flores maiores*.

γ. Statice caule nudo paniculato tereti, foliis laevibus, obverse ovatis, LINN. *Sp.* 2. *Limonium maritimum maius*. C. B. 408. 295. Ad littora maris,

δ. Statice caule ancipiti dichotomo nudo, foliis lanceolatis mucronatis radicalibus, floribus alternis di-

stantibus. LINN. Sp. 6. HALL. G. 349. GMEL. Sib. II. p. 223. T. 92. *Limonium elatius*, *Plantaginis foliis procumbentibus in aculeum terminatis, floribus albis spicatis*. AMMANN. Ruth. n. 130. In Sibiria. P.

H. Stace foliis radicalibus alternatim pinnato-sinuatis: caulinis ternis triquetris subulatis decurrentibus. LINN. Sp. 13. *Limonium peregrinum Asplenii folio*. C. B. *Limonium Rauwolfianum*. CLVS. Cur. post. p. 33. In Palaestina. A.

H. Stace caule fruticoso folioso, floribus solitariis, foliis lanceolatis vaginantibus, *monopetala*. LINN. Sp. 11. *Limonium lignosum*. BOCC. Sic. T. 17. Fr.

LAGOECIA. LINN. 251.

H. Lagoecia. LINN. Sp. 203. Cuminoides. I. R. H. *Cuminum sylvestre*. C. B. 186. CAMER. Epis. 519. In Oriente. A.

VALERIANA. LINN. 43.

Semine pappo plumoso. *Valerianae*. T.

H. Valeriana floribus monandris calcare donatis, foliis lanceolatis integerrimis. LINN. Sp. 1. HALL. H. 6. *Valeriana rubra*. DOD. Pompt. 351. I. R. H. In Italiae, Helvetiae rupibus. A.

H. Valeriana floribus monandris, foliis imis integris, superioribus pinnatifidis. LINN. Sp. 2. *Valeriana folio Calcitrapae*. C. B. I. R. H. A.

Valeriana floribus triandris dioicis, foliis pinnatis. LINN. Sp. 4. *Valeriana foliis imis ovalibus*,
Aa 5
supe-

superioribus pinnatis, sexu distincta. HALL. H. 2. a. *Mar.* *Valeriana palustris minor.* C. B. 1. R. H. *flos maior.* *Femina* β. *Valeriana palustris inodora, florum laciniata.* C. B. 1. R. H. Flores minores. In paludosis.

α. *Valeriana floribus triandris, foliis omnibus pinnatis. officinalis.* LINN. Sp. 5. HALL. H. 1. *Valeriana sylvestris.* L. CUVS. LX. *maior.* C. B. 1. R. H. BLACKW. 371. In vdis sepius & sylvis.

H. *Valeriana floribus triandris, foliis radicalibus indivisis, cauliculis pinnatis.* LINN. Sp. 6. *Valer. Dortensis.* C. B. Caulis non sulcatus. In Gallia.

H. *Valeriana floribus triandris, foliis omnibus integris ovato-oblongis, subdentatis, caule simplici.* LINN. Sp. 9. HALL. H. 4. *Valer. alpina, fol. Scrophulariae.* C. B. Prodr. 87. In Helvetiae montibus.

Semine non papposo. *Valerianellae.* TOURN. HALL.

*Valeriana caule dichotomo, foliis oblongis rari-
ter lacinis, seminis mucrone simplici. Valerianella.*
HALL. H. 1. *Valeriana.* LINN. Sp. 14. α. *Valerianella mucronis praecox humilis, semine compressa.*
I. R. H. *Lactuca agnina.* TABERN. Ubique vero
tempore in gramineis. A.

*Valeriana (Valerianella) caule dichotomo, fo-
liis plerumque ferratis, feminis corona tridentata.*
HALL. H. 1. LINN. Sp. cit. β. *Valerianella arven-
sis, serotina, alba, seminis turgidiore.* I. R. H.
Caulis

Caulis magis ramosus. Squamulae flores excipientes glabrae. Serius floret.

H. Valeriana caule dichotomo, foliis pinnatifidis, seminis corona radiatim dentata. Valerianella *femine stellato*. I. R. H. LINN. Sp. cit. var. *γ. coronata*. Squamulae floribus subiectae hirsutae, fructus hirsutus.

H. Valeriana caule dichotomo, foliis lanceolatis serratis, seminis corona globosa inflata. Valerianella *Cretica fructu vesicario*. I. R. H. Cor. 6. BOERH. I. T. 75. Semen subhirsutum in apice coronatur membranula in vesiculam globosam inflata, orificio contracto dentato, in cuius fundum apert. seminis prominet.

H. Valeriana floribus diandris tingentibus, foliis ovatis sessilibus. LINN. Sp. 2. Valerianella *Indica rubra*. C. B. Valerianella *Cornucupioidea echinata*. COLUMN. Ecphr. 306. Semen calyce quinque-dentato coronatum. In calidiori Europa.

H. Valeriana floribus tetrandris, foliis pinnatifidis, seminibus palcae ovali adnatis. LINN. Spec. 16. Valerianella *tetraseemon*, femine foliolo infidente. HALL. Commenta. Gott. 1751. p. 297. T. IX. In *Sibiria*.

BOERHAAVIA. LINN. 9.

H. Boerhaavia caule erecto, pedunculis folio glabro longioribus. An Boerhaavia *erecta*. LINN. Sp. 10. HALL. G. 352.

SCABIOSA. LINN. 108. HALL. Succisa,
HALL.

Corollae quadrifidae; involucrum seminis den-
tatum ciliatum.

Scabiosa (*Succisa*) foliis ad terram ovatis, ad
caulem oblongis parum dentatis. HALL. H. 671.
n. 1. Scabiosa corollulis quadrifidis aequalibus, cau-
le simplici, ramis approximatis, foliis lanceolato-
ovatis. LINN. Sp. 4. Succisa glabra. C. B. Succisa
f. Morfus Diaboli. MATH. 623. BLACKW. T. 142. P.
In pratis vdis, circa Hurste, Bovenden & ber-
cynicis.

H. Scabiosa foliis subhirsutis, pinnatifidis, lanceo-
latis, squamis calycinis hirsutis acutis, flore bre-
vioribus. Calycis folia exteriora lineari-lanceolata,
pilosa, squamae interiores apice fuso rigido arista-
tae. Involucrum seminis brevissime ciliatum. Se-
gmenta floris albi omnia obtusa, inferius minus,
apice viridi. A.

H. Scabiosa (*Succisa*) capitulis nudis globosis,
foliis glabris pinnatis solidis. HALL. Comm. Gott. T.
I. p. 225. En. Gott. 354. Secundum descriptionem
a priori diversa videtur.

H. Scabiosa foliis pinnatis, foliolis lanceolatis
ferratis, capitulis globosis hispidis congestis. HALL.
H. 672. n. 2. LINN. Sp. 1. Scabiosa alpina, foliis
Centaurii maioris. C. B. In Helvetia. P.

H. Scabiosa corollulis radiantibus, caule hispido,
foliis sessilibus ferratis, squamis calycinis acutis.
Omnino eadem esse videtur Scabiosa flosculis qua-
drifidis,

divifidis, foliis pinnatifidis, laciniis lateralibus erectiusculis. LINN. *Art. Vpf.* 1744. p. 11. T. I. *Tatarica* LINN. *Sp.* 7. Nam etiam nostrae folia radicalia saepe sunt pinnatifida, lobis imbricatis. In *Tataria.* *Bienn.*

Scabiosa corollulis radiantibus, caule hispido, foliis subhirsutis, imis ovatis, superioribus semipinnatis. HALL. *H.* 670. n. 1. LINN. *Sp.* 8. Scabiosa *pratensis hirsuta officinarum.* C. B. I. R. H. *Sylvestris.* BLACKW. T. 185. In pratis & ad vias.

ASTEROCEPHALVS. HALL. H. 668.

Corollulae quinquefidae; seminis involucrum annulo fimbriato plicatili coronatum; ex apice seminis setae quinque circa florem.

Asterocephalus (*Scabiosa*), foliis radicalibus ovatis crenatis, caulinis pinnatis setaceis, capitulis fructiferis globosis. HALL. *H.* 670. LINN. *Sp.* 10. Scabiosa *capitulo globoso minor.* C. B. In collibus ficcis apricis. P.

H. Asterocephalus foliis pinnato-compositis, foliolis cuneiformibus incisis, superioribus linearibus, capitulis fructiferis subrotundis. Egregie hanc exprimit Scabiosa *glabra, carnosus foliis virentibus.* HERM. *Parad. Bat. ic.* p. 221. Nec alia esse videtur Scabiosa *mont. foliis scabiosae vulgaris.* C. B. I. R. H. Priori similis, diversa tamen *caule ramossissimo altitudinis humanae, foliis omnibus pinnatis, & setis circa florem fere semper paucioribus ac quinque.* P.

H. Astero-

H. *Astrocephalus* foliis bipinnatis linearibus, ca-
pitolis radiantibus, ochroleucis. Scabiosa LINN.
Sp. 17. Scabiosa folio multifido, flore flavescente.
VIL GLVS. III. MORIS. III. S. VI. T. 13. f. 23.
III. HALLERVS dubitat, num a *vulgari* satis sit di-
versus.

H. *Astrocephalus* foliis lyratis, lobis profunde
serratis, capitulis globosis, annulo seminis amplissi-
mo, patulo. Scabiosa LINN. 11. Sc. maior Hispanica.
GLVS. I. stellata, folio laciniato, maior. C. B. 1. R. H.
MALL. G. 353. A.

H. *Astrocephalus* foliis inferioribus lyratis, ser-
ratis, superioribus linearibus integerrimis, capitu-
lis cylindricis hispidis. Scabiosa atropurpurea.
LINN. Sp. 12. Scab. peregrina rubra, capite oblon-
go. C. B. Scabiosa VI. Indica. GLVS. III. Caulis
herbaceus, annuus, diffusus. A.

H. *Astrocephalus* foliis inferioribus petiolatis
ovatis serratis, superioribus pinnatis linearibus, ca-
pitulis cylindricis hispidis. Radix perennis, caulis
firmior erectus ramosus. Folia glabra, nitida. Pe-
dunculi hirti. An Scabiosa Afr. frutescens maxi-
ma, foliis tenuissime incis. BOERH. Lugdb. I. p.
128. In Africa. P.

H. *Astrocephalus* caule fruticoso, foliis petio-
latis obovatis, crenatis, hirsutis, pedunculis hirtis
ramos terminantibus. Scabiosa LINN. Sp. 14. Sca-
biofa Afr. frutescens, foliis rugosis & crenatis.
HERM. Par. ic. p. 219. 220. Fr.

H. *Astrocephalus* foliis planis, carnosiss, infe-
rioribus pinnatis, ramorum integerrimis, calyce
flore

florem superante. GAIL. Sib. II. p. 213. T. 87.
HALL. G. 353. A.

H. *Asterocephalus* foliis carnosis hirsutis lanceolatis, calyce folioso florem excedente, corollulis aequalibus. *Planta humilis, caulis dichotome ramosus, folia sessilia, crassiuscula; folia calycis uncialia, hirsutissima, inaequalia; flos exiguus, corollulae omnes aequales, nec radiantes, limbo quinquesido regulari, carneae. Setae ex apice seminis rigidae, corollulis longiores. Fufius illam descripsi in Comment. Gott. T. V. A.*

DIPSACVS. LINN. 107.

Dipsacus foliis connatis ferratis, capitulis ovalibus, aristis calycinis rectis. HALL. H. 1. LINN. Sp. 1. α *Dipsacus sylv. aut virga pastoris, maior.* C. B. BLACKW. T. 50. Ad oras agrorum, v. g. circa *Herperhausen*. A.

H. *Dipsacus* foliis sessilibus ferratis, capitulis ovalibus; aristis calycinis recurvis. HALL. G. 355. *Dipsacus sativus, fullonum.* C. B. In usum fullonum feritur.

H. *Dipsacus* foliis connatis sinuatis. LINN. Sp. 3. *Dips. folio laciniato.* C. B.

KNAUTIA. LINN. 109.

H. *Knautia*. LINN. Sp. p. 101. *Lychni-Scabiosa, flore rubro, annua.* BOERH. Lugdb. I. p. 131. In Oriente.

SYNGENESIA. LINN. f. Flores compositi, antheris quinque coalitis.

Plantae *flore composito* dicuntur, in quibus plures flosculi calyce communi comprehensi sunt. In omnibus hisce receptaculo communi innascuntur semina, quae singula ex apice proferunt florem monopetalum tubulosum, in aliis in limbum breviter quinquefidum, in alijs in ligulam planam extrorsum versam, terminatum. Ex pariete interno tubi floralis nascuntur quinque filamenta, antheris totidem linearibus in cylindrum connatis, per quem ascendit stylus, stigmate bifido ex tubo eminente.

Flosculis omnibus tubulosis, limbo quinquefido.

I. CAPITATAE. Tubus floris longissimus, limbi segmenta longiora; capitulum oblongum ventricosum.

EGHINOPS. LINN. 829.

Singulis flosculis in globum collectis calyx peculiaris.

H. Echinops calyculis unisporis, caule multifloro, foliis spinosis, supra nudis. *Sphaerocephalus*. LINN. *Sp.* 1. *maior*. I. R. H. Carduus *sphaerocephalus latifolius vulgaris*. C. B. In Italia. P.

H. Echinops caule subunisporo, foliis duplicatopinnatifidis, foliolis latiusculis vicinis. GMEL. *Sib.* II. p. 100. LINN. *Sp.* 2. H. *Vff.* 248. *. In *Sibiria*.

H. Echinops calycibus fasciculatis: lateralibus sterilibus, foliis pinnatifidis supra strigosis. LINN. *Sp.*

Sp. 3. *Onopordus minor*, annuus, capite magno.
 1. R. H. Scabiosa *Cardusifolia annua*. HERM. Par.
 T. 224. *Sphaerocephalus annuus*. CAMER. Hort.
 ic. 45. In *Hispania*.

Placenta alveolata.

ONOPORDON. LINN. 834.

Onopordon calycibus squarrosis, foliis decurren-
tibus ovaro-oblongis sinuatis, caule brachiato. LINN.

Sp. 1. HALL. H. p. 674. *Spina alba tomentosa lili-*
zifolia sylv. D. B. *Carduus alatus tomentosus, acan-*
thi folio, vulgaris. 1. R. H. Ad vias.

β. *Carduus luteus tomentosus, Acanthi folio*
flave magno. HALL. G. 357. A.

h. *Onopordon calycibus imbricatis, caule parum*
ramoso. LINN. Sp. 3. H. Vps. 249. *Carduus to-*
mentosus, Acanthi folio, altissimus Lusitanicus.
 1. R. H. HALL G. 357.

h. *Onopordon calycibus squarrosis, foliis linea-*
tibus pinnatifidis, vix caule latioribus. LINN. Sp.
 2. *Acanthium Illyricum.* LOBEL. II. ic. 1. In *Eu-*
ropa australi. A.

h. *Onopordon acaulon, capitulis squarrosis ner-*
vo foliorum insidentibus. HALL. G. 357. c. descr. B.

Placenta paleis aut pilis rectis.

CYNARA. LINN. 835.

h. *Cynara foliis subspinosis pinnatis indivisisque,*
calycinis squamis ovatis. LINN. Sp. 1. *Cynara*

B b

hort.

hort. foliis non aculeatis. C. B. Scolymus Dioscoridis. CLVS. CLIII.

H. Cynara foliis spinosis: omnibus pinnatifidis, calycinis squamis ovatis. LINN. Sp. 2. Cynara spinosa, cuius cauliculi cstantur. C. B. Scolymus aculeatus. TABERN. 1075.

LAPPA. HALL. 674. TOURN. Artium.

LINN 830.

Lappa capitulis tomentosis. HALL Helv. 1. Artium. LINN. Sp. 1. β. Lappa minor montana, capitulis tomentosis. C. B. Vbique ad vias.

Lappa capitulis glabris. HALL. H. 2. Artium. LINN. Sp. 1. α. Lappa maior, f. Artium Dioscoridis. C. B. 1. R. H. Bardana officina. lisdem locis cum priori.

Lappa capitulis glabris minimis. HALL. H. 3.

SERRATVLA. LINN. 831.

Serratula foliis pinnatifidis, petiolis multifloris, calycibus glabris spinulosis. Serratula tinctoria. LINN. Sp. 1. α. C. B. Carduus HALL. H. 5. Serratula Matthioli. CLVS. VIII. In M. arcis Pless. P.

H. Serratula foliis pinnatifidis, petiolis unifloris, calycibus squalidis. Serratula praecoxa. LINN. Sp. 1. β. Carduus inermis, foliis glabris pinnatis, lacinia extrema maxima, capitulis squelidis. GMEL. Sib. II. T. 20. In Sibiria. Planta priori omnibus partibus multo maior. Squamae calyciniae dense conglutinatae & pulvere sordido fere tectae, nec ora alba, uti prior species, ornatae. P.

H. *Serratula* (*Carduus*) caule ramoso, foliis pinnatifidis, foliolis inferiorum dentatis, squamis calycinis ex lanceolato-spinosis, appressis. Gmel. Sib. II. n. 38. T. 17.

Serratula foliis dentatis spinosis. LINN. Sp. 141. *Carduus foliis semipinnatis, spinosis, capitulis oblongis mitioribus*. HALL. H. 5. *Carduus arvensis*. TABERN. 300. In segetibus humidioribus.

H. *Serratula* caule ramosissimo, ramis parulis, foliis petiolatis, ovato-acutis, nervosis, lunate dentatis, flosculis calycem aequantibus. Sic interim definire liceat plantulam, quae licet *figura foliorum & capituli* ad *Cirsium* atq. *boni Herinci folia*. HALL. H. T. 22. n. 16. accedere videatur, caule tamen ramosissimo, aliisque notis longe ab illo secedit. Folia inferne incana, nec lanugine albida, Calycis parvi cylindrici squamae lanceolatae, ciliatae. Flosculi carnei coloris calycem non superant. Pappus simplex, sub floris longitudine. Placenta villosa.

CARDUVS, LINN. 832. *Cirsium*.

HALL.

H. *Carduus* foliis lanceolatis dentatis ciliatis decurrentibus: spinis marginalibus duplicibus. LINN. Sp. 1. *Acarna maior, caule folioso*. C. B. *Chamaeleon Salmatensis*. BLVS. CLV. *Squamae calycinae in aetate validissimum flavescens* abest.

Carduus (*Cirsium*) foliis decurrentibus spinosis, pinnatis, laciniis divaricatis, capitulis ovatis tomentosis spinosis, caule piloso. HALL. H. 7. LINN. Sp.

50. *Carduus lanceolatus laevisfolius*. C. B. Ad. vias & in sylvis.

Carduus foliis spinosis semidecurrentibus, capitulo spinoso nutante. HALL. H. 1. LINN. Sp. 4.

Carduus nutans. P. R. H. Ad. vias.

51. *Carduus* foliis decurrentibus laciniatis, subtus tomentosis ex margine spinosis, capitulis sessilibus congestis aculeatis cylindricis. HALL. H. 677. N. 3.

Carduus LINN. Sp. 5. A. RUM.

52. *Carduus* foliis decurrentibus sinuatis ciliatis subtus tomentosis, capitulis sessilibus terminibus congestis molliter spinosis. HALL. H. 10. 2. LINN. Sp. 6. *Carduus caule crispo*. T. B. III. P. I. p. 59. P. R. H. Ad. sepes.

53. *Carduus* (*Cirsium*) caule pene nudo, alis crispis obducto, capitulis mitibus, in vertice congestis. HALL. H. 8. GMEL. Sib. II. T. 23. f. 2. *Carduus palustris*. LINN. Sp. 8. C. B. Prodr. 156. *Cirsium prat. polycephalon vulgare*. I. R. H. In p. & sylvis vdis.

β. flore albo. In sylva supra Gersmar.

H. *Carduus* (*Cirsium*) foliis ciliatis, imis ovatis, superioribus semipinnatis. HALL. Comm. Gott. T. I. p. 306. En. Gott. 362.

54. *Carduus inermis* foliis rigidis ciliatis, scapo longo paucifloro. HALL. H. 677. T. 19. *Cirsium* Bot. S. 29. Circa Jenam.

55. *Carduus* foliis amplexicaulibus angulato-spinosis, floribus solitariis, subsessilibus, obvallatis. HALL. H. 677. T. 19. *Cirsium* Bot. S. 29. Circa Jenam.

As Tuboquinis. LINN. Sp. 14. *Cirsium spinosum, var. foliorum lacteis.* HALL. G. 361. *Carduus lacteus Syriacus.* CAMER. Hort. 35. T. 10. In Syria. Hispania.

H. *Carduus foliis amplexicaulibus hastato-pinnatifidis spinosis, squamis calycinis margine apiceque spinosis.* LINN. Sp. 13. *Silybum nervis foliorum lacteis.* HALL. H. 684. *Carduus albis maculis notatus vulgaris.* C. B. I. R. H. *Carduus Mariae.* BLACKW. T. 79.

H. *Carduus foliis ex cordato-lanceolatis margine ferratis & spinosis, squamis calycum membranaceis lacris spinosis, capitulis nutantibus.* GMEL. Sib. II. p. 47. T. 19. *Cnicus foliis cordatis, petiolis crispis spinosis amplexicaulibus, floribus cernuis.* LINN. Sp. 5. H. Vpf. 251. *. In Sibiria.

H. *Carduus (Cirsium) foliis spinosis hirsutis, pinnis semibilobis, capitulis tomentosis sphaericis pungentibus.* HALL. H. 6. *Carduus eriophorus.* LINN. Sp. 15. *Card. Ericephalus.* CLVS. CLIII. *Carduus capite rotundo tomentoso.* I. R. H. In montosis Sueviae & Helvetiae. P.

Carduus (Cirsium) inerme, radicibus teretibus congestis, pinnis foliorum semibilobis, ramis unifloris. HALL. G. 360. *Carduus tuberosus.* LINN. Sp. 18. *Carduus prat. Asphodeli radice, foliis tenuiter incis.* C. B. flore purpureo. In sylvis. v. g. *Gottingenf.* & *Eulenloch.* β. flore albo *bey Harfse & Herperhausen.*

H. *Carduus foliis lanceolatis amplexicaulibus, serraturis spinoso-setaceis, capitulis triphyllis.*

Bb 3

LINN.

LINN. Sp. 22. Omnia conveniunt, sed color floris in nostra planta *purpureus* est, quem *album* dicit.

Carduus acaulis, calyce glabro. LINN. Sp. p. 1199. in *app.* *Cirsium caule floris longitudine.* HALL. 11. *Cirsium acaulos flore purpureo.* I. R. H. In ficcis acclivibus. In horto caulis elongatur. P.

CNICVS. LINN. 833.

Cnicus caule erecto, foliis inferioribus laciniatis, superioribus integris contaxis. LINN. Sp. 2. *Cirsium inerme*, foliis inter flavescentia folia congestis. HALL. H. 4. *Carduus prat. latifolius.* C. B. LOR. II. p. 11. In pratis vdis.

H. *Cnicus caule diffuso*, foliis dentato-finuatis. LINN. Sp. 1. *Cnicus sylv. hirsutior f. Carduus benedictus.* C. B. offic.

CARTHAMVS. LINN. 838.

H. *Carthamus foliis ovatis integris ferrato-aculeatis. tinctorius.* LINN. Sp. 1. *Cnicus sativus f. Carthamus officinarum.* C. B. *Cnicus vulgaris.* CLVS. CLII. In Aegypto. A.

H. *Carthamus caule piloso*, superne lanato, foliis interioribus pinnatifidis, summis amplexicaulibus dentatis. LINN. Sp. 2. *Atractylis succo sanguineo, flore luteo.* HALL. H. 685. *Atractylis lutea.* C. B. I. R. H. COLUMN. Ephr. p. 19. T. 23.

CARLINA

CARLINA. LINN. 836.

Carlina caule uniflora. HALL. H. 686. n. 2.
 LINN. Sp. 1. Carlina *acaulis*, *magno flore*. C. B.
 I. R. H. In collibus versus *Allendorf*.

β. Carlina *caulescens*, *magno flore albo*. C. B.
 I. R. H. Varietas *hortensis* eiusdem plantae, quae in
 loco culto caulem cubitalem profert.

Carlina caule multiflora, floribus terminalibus.
 LINN. Sp. 4. Carlina *sylvestris vulgaris*. C. B. In
 apricis aridis.

Floribus in radio sterilibus.

CENTAVREA. LINN. 880.

I. *Rhapontica*. Calycibus haevibus inermibus.

H. Centaurea calycibus inermibus: squamis li-
 nearibus subulatis, foliis pinnatis serratis. LINN. Sp.
 1. *Rhaponticum foliis pinnatis ciliatis, calycibus*
acutis. HALL. G. 364. *Serratula annua semine cilia-*
ri elegantissimo. I. R. H. *Senecio-Carduus apulus*.
 COLUMN. *Ecpbr.* p. 34. *ic.* In *Italia australi*.

H. Centaurea calycibus inermibus subrotundis
 glabris, squamis ovatis, foliis sinuatis. LINN. Sp.
 2. *Rhaponticum flosculis sterilibus quindecimfidis,*
squamis calycinis rotundis. HALL. l. c. *Cyanus ori-*
entalis maior moschatus. MORIS. III. S. VII. T. 25.
 f. 5. ZANON. T. 62. *Amberboi flore purpureo*
odorato. VAILL.

β. flore albo.

γ. flore luteo.

H. *Centaurea squamis calycinis ovatis glabris, foliis pinnatis, foliolis decurrentibus, serratis & integris.* GMEL. Sib. II. p. 89. T. 41. f. 2. LINN. Sp. 6. *Rhaponticum foliis pinnatis, pinnis serratis, squamis calycinis rotundis.* HALL. Comm. Gött. 1751. p. 206. *Centaureum minus* L. vulg. CLVS. X. In *Sibiria*, & alpebus *Italiae*.

Centaurea calycibus squamosis, foliis lanceolatis radicalibus sinuato-dentatis, ramis angulatis LINN. Sp. 26. *Rhaponticum foliis squalidis tomentosis, inferioribus ad b. sin semipinnatis, superioribus integris.* HALL. H. 3. *Jacea nigra prat. latifolia.* C.B. In pratis.

H. *Centaurea calycibus squamosis, foliis indivisis integerrimis decurrentibus.* LINN. Sp. 30. *Rhaponticum foliis linguiformibus decurrentibus, capitulis argenteis.* HALL. G. 364. *Centaureum minus Orient. rectum glasti folio, flore luteo.* COM-MEL. rar. T. 39. P.

Cyani. Squamis calycinis serrato-ciliatis.

Centaurea calycibus serratis: foliis linearibus integerrimis: infimis dentatis. LINN. Sp. 12. *Cyanus flosculis radialibus, latisimis, semiseptifidis.* HALL. H. 4. *Cyanus segetum.* C. B. Inter segetes. β. flore purpureo. A.

H. *Centaurea calycibus serratis, foliis lanceolatis decurrentibus, caule simplicissimo.* LINN. Sp. 11. *Cyanus caule alato, foliis integris subtus tomentosis.* HALL. H. 5. *Cyanus maior.* BLACKW. T. 266. *mont. latifolius.* C. B. I. R. H. In montosis *Helvetiae*.

• *Centaurea* (*Cyanus*). foliis ellipticis dentatis rigidis, squamis calycinis ferratis. HALL. G. 370. *Enicurus perennis coeruleus Tingitanus*, HERM. Lugdb. T. 163. *Carthamus foliis lanceolatis acuminato-seratis*. LINN. Sp. 3.

H. *Centaurea calycibus ciliatis, foliis pinnatifidis linearibus, caule paniculato*. LINN. Sp. 13. *Cyanus pinnis foliorum pinnatis, pinnis acutis gracilibus dentatis*. HALL. H. 1. *Jacea foliis candicantibus, laciniatis, calyculis non splendentibus*. I. R. H. *Stoebe Austriaca elatior*. CLVS. X. In *Thuringia*.

H. *Centaurea calycibus ciliatis, foliis tomentosis pinnatifidis, foliolis obtusis ovatis integerrimis, exterioribus maioribus*. LINN. Sp. 16. *Cyanus foliis pinnatis tomentosissimis*. HALL. G. 369. *Jacea arborea, argentea, Ragusina*. ALP. exot. 116. ZANNON. Hist. T. 92.

H. *Centaurea calycibus ciliatis subrotundis, foliis tomentosis indivisis pinnatifidisque integerrimis, caule declinato*. LINN. Sp. 19. GMEL. Sib. II. p. 96. T. 42. f. 2. *Cyanus foliis pinnatis, pinnis rotundis, superioribus ovatis, caule prostrato*. HALL. G. 368. *Comm. Gott. T. I. p. 204. In Sibiria*.

Centaurea calycibus margine ciliatis, foliis pinnatifidis, pinnis lanceolatis dentatis. LINN. Sp. 21. *Cyanus*. HALL. H. 2. *Jacea vulgaris lactuata*. I. R. H.

β. flore albo.

γ. flore ochroleuco.

Bb 5

δ. flore

g. flore flavo. Ex Sibiria. HALL. Comm. Gott. T. I. Tab. VII.

h. flore helvolo.

i. foliis latioribus glaberrimis firmulis. Ex Sibiria.

h. Centaurea (*Cyanus*) calycibus ciliatis, foliis pinnatis tomentosis, pinnis subrotundis, flore helvolo. HALL. Comm. Gott. T. I. p. 203. En. Gott. 367. In Sibiria.

Centaurea calycibus recurvato-plumosis, foliis indivisis. LINN. Sp. 7. *Cyanus foliis ovato-lanceolatis subasperis, glumis calycis in reflexam plumulam definentibus*. HALL. H. 6. Jacea IV. Austr. vill. so capite. CLVS. VII. Jacea cum squamis pinnatis f. villoso capite. I. R. H. In sylvis bercynicis.

h. Centaurea calycibus recurvato-plumosis, foliis imis pinnatis dentatis: summis lanceolatis. LINN. Sp. 8. GMEL. Sib. II. p. 98. T. 43. *Cyanus foliis calycinis plumosis reflexis, foliis pinnatis*. HALL. Comm. Gott. T. I. p. 202. Tab. 6.

h. Centaurea (*Cyanus*) calycis plumulis reflexis, foliis linguiformibus subhirsutis. HALL. G. 369. Comm. Gott. T. I. p. 201. Centaurea calycibus ciliatis, vilis setaceis reflexis, foliis integerrimis. GMEL. Sib. II. p. 100. Tab. 45. f. 1. 2.

Calcitrapae. Squamis calycinis spinosis.

h. Centaurea (*Rhaponticum*) calycibus glabris, simplicissime & breviter spinosis, foliis dentato pinnatifidis. HALL. Gott. 364. LINN. Sp. 46. Jacea foliis

foliis trichonactis, altissima, flore purpurea. I. R. H. Nostrae plantae folia omnia glabra, & lacina sunt. A.

H. *Centaurea (Rhaponticum) calycibus hirsutis, simplicissime spinosis. HALL. G. 365.*

H. *Centaurea calycibus palmato-spinosis subaequalibus, foliis decurrentibus sinuatis spinulosis. LINN. Sp. 37. H. Vps. 272. Calcitrapa caulibus amplissime alatis, spinis calycinis palmatis. HALL. G. 367. Jacea Cyanoides, caule alato. HERM. Par. 189. ic. In Creta. A.*

H. *Centaurea calycibus duplicato-spinosis, foliis lanceolatis decurrentibus subdentatis, capitulis subsessilibus lateralibus. LINN. H. Vps. 10. Solstitialis. LINN. Sp. 7. Calcitrapa lutea vulgaris, caule alato. VAILL. HALL. G. 366. Flores saepe conferti, foliis linearibus succincti. A.*

H. *Centaurea calycibus palmato-spinosis, foliis decurrentibus spinulosis repando-dentatis. LINN. Sp. 32. Jacea laciniata fionchi folio. B. Prodr. 128. HERM. Par. T. 675. Carduus maritimus canescens, alato caule. I. R. H.*

H. *Centaurea calycibus duplicato-spinosis lanatis, foliis semidecurrentibus integris sinuatisque. LINN. Sp. 39. Calcitrapa foliis linguliformibus, capitalis tomentosis. HALL. G. 366. Calcitrapa lutea, alato caule, capite eriophoro. VAILL. Act. Par. 1718. P. 212.*

Centaurea (Calcitrapa) calycibus sessilibus duplicato-spinosis, aculeis rigidissimis, foliis pinnatifidis

dis linearibus dentatis. LINN. Sp. 40. HALL. H. 689.
Carduus stellatus. C. B. *Carduus maricatus* vulgo
Calcitrapa dictus. CLVS. VII. Circa Wernigerode.

Centaurea (*Calcitrapa*) foliis semipinnatis, spinis
 calycinis ptaelongis ciliatis. HALL. G. 365.

II. CORYMBIFERAE.

Flosculi tubulosi breves, aut omnes androgyni
 limbo quinquesido, aut in ambitu feminini fistulosi,
 in capitulum planum, latum, non ventricosum
 collecti.

α. Semine nudo.

TANACETVM. LINN. 848.

Tanacetum foliis pinnatis, pinnis semipinnatis
 serratis, floribus umbellatis. HALL. H. 694. LINN.
 Sp. 7. *Tanacetum vulgare luteum*. C. B. I. R. H.
 BLACKW. T. 464. Ad ripas fossarum.

β. Var. foliis crispis. C. B.

γ. *Tanacetum* foliis ovatis integris serratis.
 LINN. Sp. 8. *Balsamita mas*. BLACKW. T. 98. Men-
 tha hort. corymbifera. C. B. In Narbona. P.

δ. *Tanacetum* foliis rigidis integris dentatis, sca-
 po unifloro. HALL. Gott. 370. *Chrysanthemum*
flosculis omnibus uniformibus hermaphroditis. LINN.
 Sp. 14. *Bellis spinosa*. ALPIN. Exot. T. 326. In
 Africa.

ARTEMI-

ARTEMISIA. LINN. 849.

H. *Artemisia* caule fruticoso ramosissimo, foliis
caulinis pinnato-multifidis linearibus, floralibus in-
divisis, spicis secundis reflexis. LINN. Sp. 3. Gmel.
Sib. II. p. 115. T. 51. *Planta fruticosa, foliis*
multifidis, pinnulis remotis, linearibus longis, ca-
pitulis subcylindricis sessilibus dense spicatis, squa-
mis calycinis ora membranaceis, flosculis omnibus
androgynis, paucioribus, vix ultra quinque. Fr.

H. *Artemisia* foliis ramosissimis setaceis, caule
erecto suffruticoso. LINN. Sp. 2. *Abrotanum mas-*
angustifolium Matis. C. B.

H. *Artemisia* foliis planis longe petiolatis dupli-
cato-pinnatis, caule procumbente laxè spicato.
HALL. H. 6. LINN. Sp. 4. *Abrotanum campestre.*
Gmel. Ad viam Thuringia. Marchia.

H. *Artemisia* foliis pinnatis, caulibus ascenden-
tibus hirsutis, floribus globosis cernuis subsolitariis,
receptaculo villoso. LINN. Sp. 8. Fl. Suec. ed. alt.
n. 733. *Abſinthium paniculatum palustre minus, argen-*
teo sericeoque folio. BOCC. Mus. II. T. 71. In Sili-
riae & Succiae rupibus. Fr.

H. *Artemisia* foliis incanis pinnatis, pinnis bilobis
floribus subglobosis pendulis secundis: receptaculo
villoso. LINN. Sp. 12. HALL. H. 4. *Abſinthium.*
BLACKW. T. 7. *Abſinthium Ponticum f. Romanum*
offinarum, Dioscoridis. C. B. 1. B. H. Passim in
ruderosis spontanea quasi. P.

H. *Artemisia* foliis bipinnatis, foliis tomentosis
nitidis, pinnis transversis, racemis simplicibus, re-
cepta-

receptaculo villos. LINN. Sp. 11. *Artemisia radice perenni, foliis pinnatis pinnatifidis incisiss serratis, calycibus subrotundis viridibus nutantibus.* GMEL. Sib. II. p. 123. T. 58.

h. *Artemisia* foliis simplicibus tomentosis obtusis lanceolatis; inferioribus cuneiformibus trilobis, receptaculo villos. LINN. Sp. 17. *Artemisia foliis radicalibus a caulinis diversis.* GMEL. Sib. II. p. 61. f. 1. 2. Fr.

Artemisia foliis pinnatifidis planis incisiss, subtus tomentosis, racemis simplicibus, floribus ovatis erectis, radio quinquefloro. LINN. Sp. 13. HALL. H. 5. *Artemisia vulgaris.* C. B. I. R. H. BLACKW. T. 431. Tabern. p. 7.

h. *Artemisia* foliis multipartitis tomentosis, floribus subrotundis nutantibus: receptaculo nudo. LINN. Sp. 9. *Abiesinthium Ponticum tenuifolium incanum.* C. B. In *Hungaria, Sibiria.* P.

h. *Artemisia inodora*, foliis incanissimè pinnatis, pinnis longis linearibus, floribus dense spicatis erectis HALL. Gott 373. *Racemi plurimi, erecti, equuli fere paralleli, floribus congestis cæcis, calyce laxo, ex viridi fusco.*

h. *Artemisia* foliis linguiformibus tomentosis, integris, lateralibus sæpe palmatis, floribus cylindricis spicatis cernuis. LINN. Sp. 15. *Abiesinthium maritimum, lavendulae folio.* C. B. *Abrotanum latifolium* Erh. *Artemisia folio.* COLUMN. Ephr. II. T. 76. Ad litorea maris.

h. Ar-

H. *Artemisia* foliis lineari-lanceolatis integerrimis glabris. LINN. *Sp.* 16. GMEL. *Sib.* II. p. 126. T. 59. 60. *Abrotanum line. folio acriori & odorato.* 1. R. H. *Draco herba.* DOD. *Pempt.* 799. In *Sibiria.* P.

H. *Artemisia* lini folio hirsuto, inodorum. AMMAN. *Ruth.* 147. Num satis a priori diversa?

COTULA. LINN. 868.

H. *Cotula* foliis lanceolato-linearibus amplexicaulibus inferne dentatis. LINN. *Sp.* 2. *Ananthocyclus Coronpi folio.* DILL. *Elth.* 27. T. 23. f. 26. In *Frisiae* inundatis. A.

H. *Cotula* receptaculis subtus inflatis turbinate. LINN. *Sp.* 4. *Chamaerictum Aethiop. lanuginosum.* BREYN. *Cent.* 148. T. 73. A.

CARPESIVM. LINN. 851.

H. *Carpesium* floribus terminalibus. LINN. *Sp.* 1. p. 859. *Tanacetum flore singulari nutante, foliis longioribus cincto.* HALL. *H.* 693. GOTT. 371. *Alter cornuus.* COLV. *Ecph.* 251. 66. In *Helvetia Italia.* A.

SANTOLINA. LINN. 847.

H. *Santolina* pedunculis unifloris, foliis quadrisariam dentatis. *Chamaecyparissus.* LINN. *Sp.* 1. *Abrotanum femina vulgare, foliis teretibus.* G. B. CLYS. 341. In *Europa australi.*

H. *Santolina* pedunculis unifloris, foliis linearibus. LINN. *Sp.* 1. *Abrotanum femina foliis Re-risina-*

risinarini. C. B. *Abrotanum femina Nardoni*. CLVS.
342. Fr

H. *Santolina corymbis simplicibus coarctatis, foliis pinnatifidis dentatis*. LINN. *Sp.* 4. *Elichrysum inodorum glabrum Coronopi folio annuum*. HERM. H. *Lugd.* T. 227. *Chrysanthemum corymbiferum*. TRIVMP. *Obs.* T. 86. *Caulis ramosissimus. Folia pinnata, pinnis distantibus lineari-lanceolatis, apice saepe trifidis. Capitula oblonga; calycis squamae flavescentes.*

ANACYCLVS. LINN. 869.

H. *Anacyclus foliis decompositis linearibus, laciniis divisis teretiusculis acutis, floribus flosculosis*. LINN. *Sp.* 3. *Chrysanthemum Valentium*. CLVS. 332.

β. Semine papposo.

GNAPHALIVM. LINN. 850. Filago.

HALL.

Gnaphalium sexubus tota planta distinctis. HALL. n. 1. *Gnaphalium caule simplicissimo, corymbo simplici terminali, sarmentis procumbentibus*. LINN. *Sp.* 1.

α. *Androgyna, flore breviori. Elichrysum mont. flore & folio rotundiore*. I. R. H.

β. *Femina, flore longiori. Elichrysum mont. folio & flore longiore*. I. R. H. In collibus siccis apricis.

H. *Gnaphalium foliis lingulatis subtus tomentosis, caule umbella lactea terminato*. HALL. H. 3. *Gnaphalium foliis lineari-lanceolatis, acuminatis, alternis*

ternis, caule superne ramoso, corymbis fastigiatis.
 LINN. Sp. 3. Gnaph. *Americanum*. CLVS. 327.

H. Gnaphalium foliis amplexicaulibus integerri-
 mis acutis subtus tomentosis, caule ramoso. LINN.
 Sp. 5. Gnaph. *Afr. latifoliam foetidum, capitulo*
argenteo. COMMEL. II. T. 56. VOLKAM. *Fl. Neri*
194. dc. β. flore aurea. P.

H. Gnaphalium foliis lanceolatis, caule tomento-
 so ramoso, floribus terminalibus glomeratis con-
 cis. LINN. Sp. 8. Elichrysum *obtusifolium, capitulis*
argenteis conglobatis. DILL. *Elth.* T. 108. P.

H. Gnaphalium foliis lineari-lanceolatis, caule infer-
 ne ramoso, corymbo decomposito terminali.
 LINN. Sp. 20. Elichrysum *Afr. folio oblongo angu-*
sto, flore rubello, postea aureo. DILL. *Elth.* T. 167.
 f. 127. P.

Gnaphalium foliis lingulatis, caule herbaceo
 simplicissimo, floribus umbellatis flavis, flosculis o-
 mnibus androgynis. HALL. G. 376. Gnaphalium,
 LINN. 19. *Stoechas citrina latifolia offic.* C. B.
Circa Rheinhausen & über der Papiermühle.

H. Gnaphalium foliis confertis angusto-lanceola-
 tis, caule fruticoso: corymbo composito. LINN.
 Sp. 13. Elichrysum *Afr. frutescens, angustis &*
longioribus foliis incanis. COMMEL. II. T. 55. Fr.

H. Gnaphalium foliis linearibus, caule fruticoso-
 ramoso, floribus corymbosis cylindricis. LINN. Sp.
 14. *Stoechas citrina tenuifolia altera.* I. B. III. 155.
In Europa calidiori.

Gnaphalium caule ramosissimo diffuso, floribus confertis terminalibus fuscis. LINN. *Sp.* 30. HALL. *H.* 9. *Elichrysum aquat. minus, capitulis foliatis.* I. R. H. Ad fossas, in agris vdis.

Gnaphalium caule simplicissimo, floribus spicatis sparsis axillaribus. *sylvaticum.* LINN. *Sp.* 31. HALL. *H.* 5. *Elichrysum spicatum.* I. R. H. In sylvis.

Gnaphalium caule erecto ramosissimo, glomerulis ad alas foliorum longe sessilibus. HALL. *H.* 11. *Gnaph. paniculatum hirsutum.* LINN. *Sp.* 32. *Filago incana tomentosa erecta.* VAILL. Locis aridis. e. g. in colle prope *Bovenden.*

Gnaphalium caule erecto ramoso, foliis perangustis, glomerulis distitis in summo caule sessilibus. HALL. *H.* 12. *montanum.* LINN. *Sp.* 12. *Filago minor.* I. R. H. In sabulosis. v. g. *M. Hagen.*

Gnaphalium caule dichotomo erecto, capitulis globosis axillaribus terminalibusque. LINN. *Sp.* 35. HALL. *H.* 10. *Gnaph. vulg. maius.* C. B. *Filago f. herba impta.* I. R. H. Locis siccis & inter stipulas.

H. *Gnaphalium* foliis alternis linearibus acutis planis, pedunculis longissimis unisporis. *saxatile.* LINN. *Sp.* 36. *Elichryso sylvestri similis, flore oblongo.* C. B. *Prodr. T.* 123.

H. *Gnaphalium (Filago)* foliis ovatis undulatis, pedunculis unisporis. HALL. *G.* 378. *Elichrysum sylv. latifolium, flore magno singulari.* I. R. H.

BACCHARIS. LINN. 853.

H. Baccharis foliis lanceolatis serratis, floribus corymbosis, caule fruticoso. LINN. *Sp.* 1. Erigeron. HALL. *G.* 379. n. 1. Conyza *frut. foliis angustioribus nervosis*. FEVILL. *Peruv. T.* 37. Senecio *Afr. arbor. folio serrato*. BOERH. *In America*. Fr.

H. Baccharis foliis obovatis, superne emarginato-crenatis glaucis, floribus in alas congestis, caule fruticoso. LINN. *Sp.* 4. Erigeron HALL. *G.* 379. Senecio *Virginianus arbor. atriplicis folio*. HERM. *Par. Bat. T.* 225. *In Virginia*. Fr.

ERIGERON. LINN. 855.

Erigeron floribus ex alis foliorum paniculatis, flosculis femininis omnibus lingulatis. HALL. *H. n.* 4. Erigeron *caule floribusque paniculatis*. LINN. *Sp.* 3. Conyza *annua acris alba elatior, Linariae foliis*. BOCC. *Sic. T.* 46. Exoticae quidem originis nunc in omnibus sylvis frequentissima. A.

H. Erigeron foliis inferioribus dentato-laciniatis, superioribus integris. LINN. *Cliff.* 407. LINN. *Sp.* 4. Senecio *Bonariensis purpurascens, foliis imis Coronopi*. DILL. *Elth. T.* 257. f. 334. *In America australi*. A.

H. Erigeron foliis lingulatis, floribus solitariis per ramos sessilibus. HALL. *G.* 380. Erigeron *ramis lateralibus multifloris, calycibus squarrosis*. LINN. *Cent. Pl. l. n.* 81. Virga aurea minor, foliis glutinosis & graveolentibus. I. R. H.

Erigeron pedunculis alternis unifloris alaribus. LINN. *Sp.* 6. HALL. *H.* 3. Conyza *coerulea*. TA-

BERN. 861. After *arv. cocculeus acris*. 1. R. H. In
siccis apricis. A.

CONYZA. LINN. 854.

Conyza foliis ovato-lanceolatis, floribus laxo
corymbosis. HALL. H. 704. *squarrosa*. LINN. Sp. 1.
Conyza maior vulgaris. C. B. In M. *Plais* aliisque
collibus siccis. P.

CHRYSOCOMA. LINN. 845.

H. *Chrysocoma* herbacea, foliis linearibus gla-
bris, calycibus laxis. LINN. Sp. 4. *Oxyris Austr.*
CLVS. 325. *Conyza Linariae folio*. 1. R. H. In
Helvetia. P.

H. *Chrysocoma* fruticosa, foliis linearibus dorso
decurrentibus. LINN. Sp. 1. *Coma aurea Afr. fru-*
ticans, foliis Linariae angustis, maior. COMMEL.
H. II. T. 45. In *Africa*.

TVSSILAGO. LINN. 856. *Petafites*. HALL. 706.

TOURN

Tussilago scapo imbricato unifloro, foliis sub-
cordatis angulatis denticulatis. LINN. Sp. 3. *Peta-*
sites scapo unifloro, flosculis in ambitu lingulatis.
HALL. H. 6. *Tussilago f. Farfara*. 1. R. H. BLACKW.
T. 204. In argillofis vdis. P.

Tussilago (*Petafites*) thyrso ovato, flosculis omni-
bus androgynis. LINN. Sp. 6. HALL. H. 7. *Petafi-*
tes maior & vulgaris. C. B. BLACKW. T. 222. Ad
Leinam & fossas. P.

Tussi-

Tussilago (Petasites) thyrsis oblongo, flosculis femineis nudis plurimis, paucissimis androgynis. LINN. *Sp.* 6. HALL. *H.* 3. *Petasites maior, flosculis pedunculis longis insidentibus.* DILL. *Elth.* T. 230. f. 297. *Bey der Weender Papiermühle; in der Contrescarpe.*

CACALIA. LINN. 841.

H. *Cacalia foliis reniformibus acutis denticulatis.* LINN. *Sp.* 10. *Cacalia.* HALL. *H.* 705. *Cacalia glabro folio.* CLVS. CXV. *In Helvetiae montosis.*

H. *Cacalia (Kleinia) caule fruticoso, foliis carnosis ovato-oblongis, petiolis basi linea triplici deductis.* LINN. *Sp.* 2. *H. Cliff.* 395. *Anteuphorbium.* DOD. *Pempt.* 378. DILL. *Elth.* T. 55. f. 2. 3. Fr.

H. *Cacalia (Kleinia) caule fruticoso, levi ventricoso, foliis lanceolatis planis, petiolorum cicatricibus obsolete.* LINN. *Sp.* 3. *Cliff.* 395. *Cacalianthemum folio Nerii glauco.* DILL. *Elth.* T. 54. f. 2. Fr.

H. *Cacalia (Kleinia) caule fruticoso terete, foliis carnosis compressis lanceolatis.* LINN. *Sp.* 2. *Senecio Afr. arboreus, Ficoidis folio & facie.* COMMEL. *rar.* T. 40.

EUPATORIUM. LINN. 842.

Eupatorium foliis plerumque ternatis. HALL. *H.* 704. *Eupat. foliis digitatis.* LINN. *Sp.* 10. *Eupat. cannabinum.* C. B. I. R. H. *Eup. Avicennae.* BLACKW. T. 110. *Ad rivas. P.*

H. Eupatorium foliis ovatis serratis petiolatis, caule glabro, calycibus simplicibus. Eupatorium *Scrophulariae folio, floribus albis*. HALL. G. 381. Ageratum LINN. Sp. 3. Valeriana *Vrticae folio, flore albo*. CORNYT. Canad. T. 21. P.

Semine aristis coronato.

AGERATVM. LINN. 843.

H. Ageratum caule piloso, foliis petiolatis hirsutis ovatis crenatis, stylis flore longioribus reflexis. Ageratum. LINN. I. *Flosculi albi*. An Eupatorium *Betonicae folio*. HERM. Par. ic. p. 162. ? Sed folia sessilia ibi pinguntur.

H. Ageratum caule subhirsuto, foliis petiolatis oblongis crenatis glabris ora ciliatis, stylis tubum floris vix superantibus. An Eupatorium *Amer. berbaceum, Melissa folio, villosum, flore coerulesco*. HALL. G. 323 ?

XERANTHEMVM. LINN. 851.

H. Xeranthemum foliis lanceolatis patentibus, receptaculo paleaceo, feminum pappo quinqueseto. LINN. Sp. 1. Ptarmica *Austriaca*, CLVS. XI.

BIDENS. LINN. 840.

Bidens foliis tripartito divisis, calycibus subliosis, seminibus erectis. LINN. Sp. 1. HALL. H. 709. n. 1. Cannabina *aquat. folio tripartitum diviso, c. b.* Ad fossas. β . Var. flore radiato.

Bidens

Bidens foliis longe ellipticis indivisis, seminibus erectis, floribus cernuis. HALL. H. 2. LINN. Sp. 2.
Bidens folio non dissecto. 1. R. H.

β. Varietas *flosculis in radio lingulatis, semina abortivo.* *Coreopsis foliis lanceolatis serratis oppositis amplexicaulibus.* LINN. Sp. 7.

H. *Bidens* foliis pinnatis subpilosis, corollis flosculosis, calycibus simplicibus, seminibus divergentibus. LINN. Sp. 5. *Bidens latifolia hirsutior, semine angustiore radiato.* DILL. Elth. T. 43. f. 51.

H. *Bidens* foliis ovatis & tripteris, caulibus hirtis brachiatis. HALL. G. 383. *Bidens foliis ovatis serratis: inferioribus oppositis, superioribus ternatis; intermedio maiore.* LINN. Sp. 11. c. descr. A.

RADIATAE.

Flores in disco tubulosi, quinquefidi, androgyni; in radio lingulati, feminini aut abortivi.

I. Placenta paleacea, semine paleis coronato.

VERBESINA. LINN. 874.

H. *Verbesina* foliis alternis decurrentibus undulatis obtusis. LINN. Sp. 1. *Chrysanthemum Curassavicum, alato caule, floribus aurantiis.* HERM. Par. T. 125. In *Surinamo.* A.

H. *Verbesina* foliis oppositis lanceolatis serratis, floribus alternis geminis subsessilibus. *prostrata.* LINN. Sp. 8. *Eupatorio-phalacron menthae arvensis folio.* DILL. Elth. T. 113. f. 138. Eandem iconem addita descripsi in *Comm. Soc. Gott. T. III. Tab. XI.*

COREOPSIS. LINN. 879.

H. Coreopsis foliis subternatis integerrimis. LINN. *Spec.* 2. H. *Vpf.* 269. *. HALL. G. 385. Chrysanthemum *Virgin. folio acutiore lacvi trifoliato, f. anagyridis folio.* MORIS. III. S. VI. T. 3. f. 44. P.

H Coreopsis foliis lanceolatis ferratis alternis petiolatis decurrentibus. LINN. *Sp.* 8. HALL. G. 385. Chrysanthemum *caule alato Canadense.* ZANON. T. 58.

HELIANTHVS. LINN. 877.

H. Helianthus radice annua, flore amplissimo nutante. LINN. *Cliff.* 419. *foliis omnibus cordatis, nervis pone basin unitis: extorsum denudatis.* LINN. *Sp.* 1. Corona solis, I. R. H. A.

H. Helianthus foliis inferioribus cordatis: nervis prope basin unitis denudatis: superioribus ovatis, multiflorus. LINN. *Sp.* 2. Hel. *radice tereti inflexa perenni.* *Cliff.* 419. Corona Solis minor femina. TABERN. ic. 764. P.

H. Helianthus foliis ovato-cordatis: nervis infra folium unitis, radice tuberosa. LINN. *Sp.* 3. Corona Solis, *flore parvo, radice tuberosa.* I. R. H. In *Brassilia.*

H. Helianthus foliis lanceolatis scabris, caule stricto: inferne glabro, radice fusiformi. LINN. *Sp.* 6. Helianthus *maior Virginianus.* ZANON. *Hist.* T. 58. P.

RVDBECKIA. LINN. 878.

H. Rudbeckia foliis pinnatis, extrema pinna amplissima. HALL. G. 386. *foliis compositis laciniatis.* LINN. Sp. 1. Obeliscotheca *Hydrophylli folio, foliorum lobis latioribus.* VAILL. β . Variet. *lobis angustioribus.* P.

H. Rudbeckia foliis asperis ovatis dentatis, radii petalis emarginatis. LINN. Sp. 3. Obeliscotheca *integrifolia, radio aureo, umbone atrorubente.* DILL. Elth. T. 218. f. 285.

H. Rudbeckia foliis oppositis hirsutis ovato-acutis, calyce imbricato cylindrico; radii petalis pistillatis. Sic interim definire liceat plantam, quae, si vlla, noui generis nomen meretur. *Calyx cylindricus uncialis, squamis rotundatis dense imbricatus. Flores radii apice emarginato, immarcescibiles. Discus conicus. Semina omnia longa triquetra, quae in radio, nuda, cum petalo cartilagineo unita; quae sub disci floribus, simplici aut duplici longiori arista coronata; squamae calycis longitudine, semina distinguunt.* A Rudbeckia, cuius habitum refert, petalis radii femininis & calyce differt. Vide iconem sub fine libri.

SIEGESBECKIA. LINN. 873.

H. Siegesbeckia LINN. Cliff. 412. T. 23. *. H. Vps. 267. HALL. G. 385. *petiolis sessilibus, calycibus incrassatis.* LINN. Sp. 1. Bidenti similis, *foliis latissimis serratis.* BVXB. Cent. III. T. 52. In Media. A.

BUPHTHALMUM. LINN. 876.

H. Bupthalmum foliis oppositis lanceolatis, petiolis bidentatis. LINN. *Sp.* 1. *Asteriscus frutescens, Leucoii foliis sericeis & incanis.* DILL. *Elth.* T. 38. f. 44. In *Virginia.* Fr.

H. Bupthalmum calycibus acute foliosis, ramis alternis, foliis lanceolatis amplexicaulibus integerrimis. *spinosum.* LINN. *Sp.* 2. After. I. CLVS. XIII. After *luteus, foliis ad florem rigidis.* C. B.

H. Bupthalmum calycibus obtuse foliosis sessilibus axillaribus, foliis oblongis obtusis. LINN. *Sp.* 3. After. *annuus odoratus, Lusitanicus.* I. R. H. ZANON. *Hist.* T. 24. In *Creta. Lusitania.* A.

H. Bupthalmum foliis lineari-lanceolatis, serratis, villosis, calycibus nudis. HALL. *H.* 711. n. 2. *salicifolium.* LINN. *Spec.* 5. After III. *Austr.* I. CLVS. XIII. After *luteus maior, foliis succisae.* C. B. In *Helvetia.*

H. Bupthalmum calycibus foliosis, foliis cordatis serratis trinerviis basi hinc brevioribus. LINN. *Sp.* 7. HALL. *G.* 388. *Chrysanthemum Scrophulariae folio Americanum.* PLVKN. T. 99. f. 1. P.

Bupthalmum foliis incanis pinnatis, pinnis argute serratis. HALL. *H.* 3. *Anthemis* LINN. *Sp.* 16. *Chamaemelum Chrysanthemum.* I. B. III. p. 122. Bupthalmum *Tunaceti minoris foliis.* C. B. I. R. H. In sylvis & collibus.

H. Bupthalmum (*Anthemis*) caulibus simplicibus unifloris decumbentibus, foliis pinnato multifidis.

LINN.

LINN. *Sp.* 12. *Pyrethrum officinarum*. BLACKW. *T.* 390. DOD. *Pempt.* 327.

ACHILLEA. LINN. 871.

H. Achillea foliis ellipticis obtusis acute serratis fasciculatis, floribus umbellatis. HALL. *G.* 389. LINN. *Sp.* 2. *Ageratum foliis serratis*. C. B. *Parmica lutea suaveolens*. I. R. H. *Balsamita minor*. DOD. *Pempt.* 295. In Gallia Narbonensi.

Achillea foliis integris lanceolatis, minutissime serratis. LINN. *Cliff.* 412. HALL. *H.* 1. LINN. *Sp.* 10. *Parmica vulgaris*. CLVS. XII. BLACKW. *T.* 276. *foliis longis serratis, flore albo*. I. R. H. In pratis. v. g. gegen die Mäschmühle. Var. hort. *flore pleno*.

Achillea foliis punctatis bipinnatis nudis, lacinii linearibus dentatis. LINN. *Sp.* 14. HALL. *Holv.* 8. *Millefolium*. BLACKW. *T.* 13. *vulgare album*. C. B. I. R. H. β . *vulgare purpureum minus*. I. R. H. Prior var. vbique in gramineis; altera potius *bortensis*. P.

Achillea foliis bipinnatis, pinnis primis remotis, ultimis confertis, pinnulis semipinnatis. HALL. *H.* 9. LINN. *Sp.* 15. *Millefolium nobile*. I. R. H. Achillea *Millefolia odorata*. I. B. III. p. 145. In M. Altenstollberg; circa Stollberg & Kulbra in muris. P.

H. Achillea pinnulis foliorum aequalibus confertis tomentosis, pinnulis imorum trifidis, summorum simplicibus. HALL. *H.* 11. LINN. *Sp.* 4. *Millefolium tomentosum luteum*. C. B. *Stratiotes millefolia flavo flore*. CLVS. 330. In Valleſia. P.

H. *Achillea* foliis pinnatis, pinnis semipinnatis, foliolis lanceolatis incis fissatis serratis subtus lanigeris. LINN. *Sp.* 5. *Millefolium Orient. foliis Tanacetii incanitis, radiis pallide luteis.* BOERH. *Eugdb.* L. p. 112. I. R. H. *Cor.* 37.

ANTHEMIS. LINN. 870. *Chamaemelum.* T.

H. *Anthemis* caule ramoso, foliis pinnato-multifidis setaceis, calycibus villosis pedunculatis. LINN. *Spec.* 13. *Santolina foliis repetito-divisis planis,* HALL. *G.* 375. *Cotula flore luteo radiato.* I. R. H.

H. *Anthemis* foliis pinnato-multifidis, foliorum apicibus subspinosis. LINN. *Sp.* 2. *Chamaemelum Leucanthemum Hisp. magno flore.* C. B. *Prodr.* 70. In *Gallia Narbonensi.*

H. *Anthemis* foliis planis pulposis pinnatis, pinnis dentatis, caule ramoso. HALL. *G.* 391. LINN. *Sp.* 3. *Chamaemelum maritimum.* I. B. III. p. 122.

H. *Anthemis* foliis pinnato-multifidis planis; laciniis linearibus acutis trifidis, pedunculis longissimis. LINN. *Sp.* 10. *Chamaemelum Orient. absinthii folio.* I. R. H. *Cor.* 37.

H. *Anthemis* foliis pinnato-compositis linearibus acutis subvillosis. LINN. *Sp.* 7. *Chamaemelum foliis lanuginosis pinnatis, pinnis capillariter multifidis brevioribus.* HALL. *H.* 2. *Chamaemelum odoratum.* DOD. *Pempt.* 270. *Chamaem. nobile f. Leucanthemum odoratius.* C. B. I. R. H. In *Helvetia.*

β. Var. *flore pleno & semipieno.*

H. *Chamaemelum* Chium vernum, folio crassiore, flore magno. 1. 4. H. Cor. 37. *Flos maior. Folia lata & robusta*; LINNAEVS tamen pro varietate prioris habet.

Anthemis (*Chamaemelum*) foliis pene glabris, bipinnatis, pinnis latiusculis linearibus. HALL. H. 3. Anth. receptaculis conicis, paleis setaceis, seminibus nudis. LINN. Sp. 9. *Cotula foetida*. BLACKW. T. 67.

MATRICARIA. HALL. 718. LINN. 867. *Chrysanthemum*. LINN. 866.

H. *Matricaria* (*Chrysanthemum*) foliis lanceolatis superne ferratis vtrinque acuminatis. LINN. Sp. 2. *Bellis Amer. procerior serotina ramosa, flore amplissimo*. PLVKN. T. 17. f. 2.

Matricaria foliis imis petiolatis; caulinis amplexicaulibus ferratis. HALL. H. 3. *Chrysanthemum*. LINN. Sp. 3. *Bellis sylv. caule folioso, maior*. C. B. *Bellis maior*. CAMER. In gramineis.

Matricaria foliis glaucis amplexicaulibus, profunde dentatis. HALL. G. 393. *Chrysanthemum segetum*. LINN. Sp. 10. CLVS. 334. *Chrysanth. folio minus secto glaucum*. I. B. III. 103. Vbiq̃ue in arenosis circa *Hesfeld*, versus *Seeburg*: in ducatu *Cellenfi*. A.

H. *Matricaria* (*Chrysanthemum*) foliis pinnatifidis incisis extorsum latioribus. LINN. Sp. 11. *Chrysanth. maius folio valde laciniato, flore croceo*. I. B. III. p. 104. *Chrysanth. Creticum*. CLVS. 335. β. Var. *semiflosculis fistulosis*. A.

H. *Matricaria foliis pinnatis, pinnis distantibus parallelis semipinnatis.* HALL. G. 392. *Matricaria Orient. folio Tanaceti, flore suaverubente.* I. R. H.

Matricaria foliis compositis planis, foliolis ovatis incis, pedunculis ramosis. HALL. H. 2. LINN. Sp. 1. *Matricaria.* BLACKW. T. 192. *vulgaris.* C. B. I. R. H. In cultis & ruderatis. In horto radios saepe deponit.

Matricaria foliis pinnatis, pinnis laxe & capilliter multifidis, receptaculis haemisphaericis. HALL. H. 1. *Matricaria Chamomilla.* LINN. Sp. 3. *Chamaemelum vulgare f. Leucanthemum Dioscoridis.* C. B. I. R. H. BLACKW. T. 296. In agris ante portam Albanam ad radicem M. Heymberg. Circa Noribergam abunde.

PYRETHVM. HALL. 720. *Chrysanthemi*
Sp. LINN.

Pyrethrum foliis pinnatis, pinnis semipinnatis, serratis. HALL. H. 1. *Chrysanthemum.* LINN. Sp. 12. *Matricaria folio Tanaceti, flore maiore, semine umbilicato.* I. R. H. *Tanacetum inodorum.* L. CLVS. 338. In sylvis Montanis. P.

H. *Pyrethrum foliis incanis triplicato-pinnatis tomentosis.* HALL. G. 393. *Ptarmica Orient. foliis Tanaceti incanis, semisflosculis brevioribus.* I. R. H.

H. *Pyrethrum foliis pulposis, rariter pinnatis, pinnulis bidentatis & tridentatis.* HALL. Gott. 393. *Chrysanthemum* LINN. I. *Leucanthemum Canariense, sapore Pyrethri.* WALTH. Hort. T. 24.

CALEN-

CALENDULA. LINN. 885.

H. Calendula foliis ovatis integerrimis, floribus parvis pallentibus. HALL. H. 722. *Caltha arvensis*. C. B. I. R. H. In *Helvetia & Thuringia* ad lacum falsum. A.

H. Calendula foliis ovatis acutis, flore magno aurantio. Calendula *feminibus radii cymbiformibus echinatis: disci bicornibus*. LINN. Sp. 1. utramque iungit.

H. Calendula foliis ovato-lanceolatis dentatis hirsutis, pedunculis superne incrassatis. LINN. Sp. 3. Calendula *humilis Afr. flore intus albo, extus violaceo*. HERM. Lugdb. T. 105. Radii *extus dilute violacei*. A.

H. Calendula foliis linearibus subdenticulatis & integris, radiis floris distantibus. An *Bellis Afr. florum pedunculis foliosis, foliis angustis & integris*, COMMEL. H. II. T. 34. A.

OSTEOSPERMUM. LINN. 887.

H. Osteospermum foliis oppositis palmatis, petiolis foliosis. HALL. Gott. 395. c. *charact.* LINN. Sp. 1. *Chrysanthemum perenne Virgin. maius, Plutani Orient. folio*. MORIS. III. S. VI. T. 7. f. 55. P.

H. Osteospermum foliis ovalibus rigidis dentatis. LINN. Sp. 3. *Chrysanthemoides Afrum, Populi albae foliis*. DILL. Eltb. T. 68. f. 79. In *Aethiopia*.

ERIOCEPHALVS. LINN. 890.

H. Eriocephalus *Africanus*. LINN. *Sp.* 926. Eriocephalus *sempervirens*, foliis fasciculatis & digitatis. DILL. *Elth.* T. 110. f. 134. Fr.

Semine paleis coronato.

TAGETES. LINN. 865.

H. Tagetes caule subdiviso diffuso. *petula*. LINN. *Sp.* 1. Flos Africanus. DODON. *Pempt.* 255. Tagetes *minor flore fulvo maculato*. DILL. *Elth.* T. 279. f. 361. In Mexico. A.

H. Tagetes caule simplici erecto, pedunculis nudis vni-floris. LINN. *Spec.* 2. Tagetes *maior, flore luteo pallescente*. VAILL.

β. flore aurantio, simplici.

γ. flore pleno.

δ. radiis fistulosis. Tagetes *Mexicanus, flore fistuloso, simplex*. COLUMN. *Ecpbr.* II. T. 46. f. 1. A.

ARCTOTIS. LINN. 836.

H. Arctotis foliis pinnatis, pinnis undulatis bipinnatis, subtus tomentosis. HALL. *G.* 406. LINN. *Sp.* 3. Anemonospermus *Afra, folio Jacobae tenuiter laciniato, flore aurantio pulcherrimo*. BOERH. *Lugdb.* T. 100. COMMEL. *H.* II. T. 22. Fr.

H. Arctotis foliis tomentosis pinnatis, pinna extrema maxima, flosculis radiantibus vicinis. HALL. *G.* 407. LINN. *Sp.* 1. Anemonospermus *Afra, foliis & facie Taraxaci incanis*. BREYN. *ic.* 27. T. 15. A.

HELE-

HELENIA. LINN. 863.

H. *Helenia foliis decurrentibus*. LINN. *Sp.* 1.
After *Floridanus aureus*, *caule alato*. PLVKN. *T.*
372. *f.* 5. After *luteus alatus*. CORNV. *Canad.*
T. 63. P.

Semine papposo, placenta nuda.

BELLIS. LINN. 864.

Calyx haemisphaericus imbricatus ex duplici serie squamarum.

Bellis scapo nudo unifloro, semine villoso. LINN.,
Sp. 1. HALL. *H.* 722. *Bellis minor sylvestris*. C. B.
I. R. H. BLACKW. *T.* 200. In omnis pratis. *Semina*
brevi pappo semper coronata. β . Var. *hort. flore*
pleno.

H. *Bellis scapo nudo unifloro*, semine papposo.
Bellidastrium. HALL. *Helv.* 723. *Doronicum*. LINN.
Sp. 4. In montibus *Helveticis*.

H. *Bellis caule diviso folioso*, foliis lanceolatis.
After LINN. *Sp.* 19. *Bellis ramosa umbellifera*. COR-
NVT. *Canad.* *T.* 194. In *Canada*. A.

INVLA. LINN. 860.

Calyx latus, laxis. Flosculi plurimi. Antherae
inferne setas demittunt.

H. *Inula foliis ovatis rugosis*, subtus tomentosis
ferratis, calycum squamis ovatis. LINN. *Sp.* 1.
After HALL. *H.* 6. After *omnium maximus*, *Hele-*
nium dictus. I. R. H. *Helenium vulgare*. C. B. In
Anglia. *Helvetia*. P.

• *Inula* foliis radicalibus petiolatis lingulatis, caulinis subamplexicaulibus serratis, floribus corymbosis. After *luteus folio succisae*. C. B. HALL. G. 397. *Conyza* III. *Austr.* ci. vs. XX. An *Inula*. LINN. Sp. 3. Fufius hanc & fequentem descripsi in *Comment. Gott. T. V. p. 13. fqq.* Circa *Berolinum, Jenam, A.*

• H. *Inula* foliis lanceolato-ovatis serratis, asperis, subtus incanis, caule subunifloro. *Comm. Gott. T. V. l. c.* After HALL. H. 5. After *luteus, hirsuto salicis folio*. C. B. 266. Circa *Berolinum, A.*

• *Inula* foliis lanceolatis recurvis serrato-ciliatis glabris, floribus folitariis, ramis angulatis. LINN. Sp. 7. After HALL. H. 8. After *mont. luteus, salicis folio glabro*. C. B. In M. *Altenstollberg. P.*

• *Inula* foliis oblongis amplexicaulibus, caule hirsuto paniculato, squamis calycinis setaceis. *dy-senierica*. LINN. Sp. 5. After. HALL. H. 3. After *prat. autumnalis, Conyzae folio*. I. R. H. *Conyza media*. C. B. LOB. ic. 345. Ad fossas. A.

• *Inula* caule ramosissimo prostrato, foliis undulatis amplexicaulibus, radiis brevissimis. LINN. Sp. 6. After HALL. H. 4. *Conyza pulicaria*. BLACKW. T. 103. After *palustris parvus, globoso flore*. I. R. H. In agris vdis & ad fossas.

H. *Inula* foliis linearibus carnosis tricuspidatis. LINN. Sp. 11. After *marit. folio tereti crasso tridentato*. I. R. H. *Crithmum Chrysanthemum*. DOD. *Pempt.* 706.

ASTER. LINN. 858.

Calyx oblongus. Flosculi pauciores. Semiflosculi plurimi calyce longiores.

After

Aster foliis lanceolatis integerrimis carnosis glabris, ramis inaequatis, floribus corymbosis. LINN. *Sp.* 4. GMEL. *Sib.* II. p. 187. *T.* 80. f. 2. *Tripolium maius coerulæum.* C. B. DOD. *Pempt.* 379. In pratis salis circa *Harste.*

H. **Aster foliis lanceolatis scabris trinerviis integerrimis, supremis ovatis, pedunculis nudiusculis corymbosis, squamis calycinis obtusis.** LINN. *Sp.* 5. **Aster** HALL. *H.* 2. **Aster** *atticus coeruleus, vulgaris.* C. B. I. R. H. BLACKW. *T.* 109. DOD. *Pempt.* 266. In M. *Pless.* P.

H. **Aster foliis omnibus ovato-lanceolatis scabris integerrimis, pedunculis corymbosis, squamis calycinis acutis ciliatis.** HALL. *G.* 396. **Aster** *latifolius, Tripolii flore.* I. R. H. P.

H. **Aster ramis divaricatis, foliis ovatis serratis, floralibus integerrimis, pedunculis unifloris nudis, divaricatus.** LINN. *Sp.* 6. GRON. *Virg.* 99. In *Virginia.* P.

H. **Aster caulibus asperis, foliis lanceolatis integerrimis semiamplexicaulibus, floribus terminalibus, pedunculis foliosis.** LINN. *Sp.* 15. HALL. *G.* 396. **Aster** *novae Angliae altissimus hirsutus, floribus amplis purpureo-violaceis.* HERM. *Par.* T. 98. VOLKAM. p. 50. ic. P.

H. **Aster foliis amplexicaulibus integris linguatis glabris, pedunculis foliosis unifloris, calycibus squarrosis.** HALL. *G.* 396. **Aster** *novi Belgii latifolius, paniculatus, floribus dilute violaceis.* HERM. *Lugdb.* T. 67. An **Aster** LINN. 25. ? P.

H. After (*Solidago*) foliis lanceolatis veniosis amplexicaulibus glabris, pedunculis foliosis, calycis squamis lanceolatis appressis. HALL. G. 400. After *novi Belgii latifolius*, *paniculatus*, *floribus saturate violaceis*. HERM. Lugdb. p. 67. ic. P.

H. After (*Solidago*) foliis ovato-lanceolatis serratis petiolatis, caule paniculato, ramis multifloris. HALL. G. 400. LINN. Sp. 17. After *latifolius autumnalis*. CORNVT. Canad. T. 65. P.

H. After (*Solidago*) caule præalto ramosissimo, foliis linearibus integerrimis, floribus terminalibus. HALL. G. l. c. LINN. Sp. 7. After *novae Angliae*, *Linariae foliis*, *Chamaemeli floribus*. HERM. Par. T. 95. Flores albi. P.

H. After (*Solidago*) foliis linearibus trinerviis, floribus corymbosis. HALL. G. 400. After *Tripolii flore*. I. R. H. P.

H. After caule paniculato, pedunculis racemosis, pedicellis foliosis, foliolis linearibus integerrimis. *ericoides*. LINN. Sp. 16. After *ericoides dumosus*. DILL. Elth. T. 36. f. 40. P.

H. After foliis ovatis angulatis dentatis petiolatis, calycibus terminalibus patentibus foliosis. LINN. Sp. 29. After *Chenopodii folio annuus*, *flore ingenti specioso*. DILL. Elth. T. 34. f. 38. A.

SOLIDAGO. LINN. 859.

Calyx oblongus, imbricatus. Semiflosculi pauciores, calyci subaequales. Flores plerumque racemosi.

H. *Solidago* foliis lanceolatis subcarnosis glaberrimis marginè scabriusculis, panicula corymbosa.

LINN.

LINN. Sp. 1. *Solidago maxima*. CORNYT. *Canad.* 168. *Virga aurea Novboracensis glabra, caulibus rubentibus, foliis angustis glabris*. HERM. Fl. 26. A.

H. *Solidago paniculato-corymbosa, racemis recurvis, floribus adscendentibus, foliis trinerviis subseratis scabris*. LINN. Sp. 2. *Virga aurea angustifolia, panicula speciosa, Canadensis*. PLVKN. T. 236. f. 1. P.

H. *Solidago paniculato-corymbosa, racemis recurvis, floribus adscendentibus, foliis enerviis subintegerrimis, altissima*. LINN. Sp. 3. *Virga aurea altissima serotina, panicula speciosa patula*. MARTYN. Cent. T. 14. P.

H. *Solidago caule obliquo, spicis secundis, pedunculis erectis foliolatis ramosis, foliis lanceolatis integerrimis*. LINN. Sp. 6. HALL. G. 399. *Virga aurea Limonii folio, panicula uno versu disposita*. I. R. H. PLVKN. T. 235. f. 2. In Mexico. P.

H. *Solidago caule flexuoso, foliis ovatis acuminatis serratis, racemis lateralibus simplicibus*. LINN. Sp. 7. *Virga aurea mont. Scrophulariae folio*. PLVKN. T. 235. f. 3. *Virga aurea Can. Asterisci folio*. HERM. Par. T. 244.

Solidago caule subflexuoso angulato, racemis paniculatis erectis confertis. LINN. Sp. 9. HALL. H. 729. *Virga aurea vulgaris latifolia serrata*. C. B. I. R. H. BLACKW. T. 169. In sylvis.

SENECIO. HALL. 730. LINN. 857.

Othonna. LINN. 888.

Calyx conicus ex squamis linearibus parallelis, simplici serie in cylindrum conglutinatis, accedentibus ad basin squamulis vagis.

H. Senecio corollis nudis, foliis amplexicaulibus laevibus laceris, caule herbaceo erecto. LINN. *Sp.* 1. Senecio *Amer. altissimus, blattariae f. hieracii foliq.* HERM. *Par. T.* 226. A.

• Senecio corollis nudis, foliis pinnato-sinuatis amplexicaulibus, floribus sparsis. LINN. *Sp.* 4. HALL. *H.* 7. Senecio *minor vulgaris.* C. B. 1. R. H. Erigeron *Senecio.* BLACKW. *T.* 132. In agris & hortis. A.

H. Senecio corollis revolutis, foliis sinuato-dentatis, inferioribus petiolatis, pedunculis trifloris folio longioribus. Senecio *triflorus.* LINN. *Sp.* 5. ROYEN. *n.* 13. Senecio *Aegyptius, foliis Matriariae.* 1. R. H. A.

H. Senecio corollis revolutis, foliis amplexicaulibus lanceolatis dentatis, squamis calycinis apice purpureis. Senecio *lividus.* LINN. *Sp.* 7. H. *Vpf.* 267. *n.* 7. Jacobaea *Hisp. minus laciniata, petalis brevissimis.* 1. R. H. *Folia saepe subtus rubent.* A.

H. Senecio corollis revolutis, foliis amplexicaulibus apice trilobis serratis, squamis calycinis sphaecelatis laceris. LINN. *Sp.* 8. H. *Vpf.* 4. *c. descr.* In *Hispania.* A.

• Senecio corollis revolutis, foliis viscidis pinnatis, pinnis angulatis, squamis calycinis laxis longitudine perianthii. HALL. *H.* 5. LINN. *Sp.* 9. Senecio *hirsutus viscidus graveolens.* DILL. *Elth. T.* 258. *f.* 336. Senecio *incanus pinguis.* In sylvis caeduis, in primis areis carbonariorum. A.

β. *Jacobaea Senecionis folio incano*. I. R. H. HALL. *Rupp. Fl. Jen. T. III.* Senecio. LINN. *Sp.* 10. Flores minores, folia minime viscida.

Senecio corollis radiantibus, foliis pinnatis, pinnis multiformibus dentatis, floribus corymbosis. HALL. *H.* 3. LINN. *Sp.* 19. *Jacobaea vulgaris laciniata*. C. B. I. R. H. *Ad fossas*. P.

Senecio corollis radiantibus, foliis pinnatifidis, subtus tomentosis, pinnis linearibus patentissimis, rariter incis. MOEHRING. *Commerc. Nor.* 1743. p. 117. HALL. *H.* 4. Senecio LINN. *Sp.* 15. *Jacobaea altissima, foliis Erucæ Artemisiaeve similibus*. RVPP. *Fl. Jen.* 176. In sylvis. v. g. *Eulenloch*. P.

H. Senecio corollis radiantibus foliis semiamplexicaulibus, semipinnatis, pinnis aequalibus angulatis, HALL. *G.* 402. LINN. *Sp.* 13. *Jacobaea Afr. annua, floribus suaverubentibus*. VDLK. *Noric.* p. 225. *ic.* A.

H. Senecio foliis pinnatifidis tomentosis, laciniiis sinuatis, caule fruticoso. Othonna LINN. *Sp.* 9. *Jacobaea maritima*. C. B. *Cineraria*. DOD. *Pempt.* 642.

Senecio corollis radiantibus, floribus corymbosis, foliis vtrinque lanceolatis, serratis, glabriusculis. LINN. *Sp.* 23. HALL. *H.* 9. *Solidago Sarvaetica*. DOD. 141. *Consolida aurea*. TABERN. 556. In sylvis. P.

Senecio corollis radiantibus, foliis ovato-lanceolatis subtus hirsutis, floribus corymbosis. HALL. *H.* 10. LINN. *Sp.* 22. *Jacobaea nemorensis, latiore*
Dd 4 rigi-

rigidiore & hirsuto folio. RVPP. *Jen.* 142. In sylvis *hercynicis*. P.

1. Senecio foliis ensiformibus acute serratis, subtus villosis, floribus corymbosis. LINN. *Sp.* 21. HALL. *H.* 11. *Consolida palustris*. TABERN. 555. *Jacobaea palustris altissima, foliis serratis*. I. R. H. In lacu ad *Denkerhausen*. P.

2. Senecio (*Solidago*) palustris, altissima, incana & tomentosa, foliis latioribus serratis. AMMANN. 220.

H. Senecio foliis cordato-oblongis amplexicaulis scabris acuminatis serratis, caule fruticoso. LINN. *Sp.* 26. *Jacobaea Afr. frutescens, foliis rigidis & hirsutis*. COMMEL. II. p. 149. T. 75. Fr.

H. Senecio foliis imis cordatis serratis, superioribus lingulatis integris. HALL. G. 403. *Solidago*. GMEL. *Sib.* II. T. 71. Senecio. LINN. *Sp.* 6. *Jacobaea Pannonica*. II. CLVS. XXII. In sylvis *Thuringiae*.

H. Senecio foliis suborbiculatis denticulatis petiolatis. Othonna. LINN. *Sp.* 1. *Jacobaea Afr. benderae terrestris folio, repens*. COMMEL. II. p. 147. T. 73.

H. Senecio foliis subcordatis crenatis, caule subnudo simplicissimo. Othonna. LINN. *Sp.* 4. *Jacobaeastrum Cacaliae folio*. AMMANN. *Ruth.* 221. T. 24. In *Sibiria*.

H. Senecio foliis crenatis, inferioribus petiolatis ovato-acutis, caulinis sessilibus lanceolatis, floribus

bus corymbosis. Dorla Orient. Limonii folio amplissimo. DILL. Elzb. T. 105.

ARNICA. LINN. 861.

Arnica foliis ovatis integris, caulinis geminis oppositis. LINN. Sp. 1. HALL. H. 737. Doronicum Plantaginis folio, alterum. C. B. 1. R. H. Doronicum VI. Pannonicum. CLVS. XVIII. Arnica offic. In omnibus pratis Hercynicis. P.

DORONICVM. LINN. 862.

H. Doronicum foliis cordatis obtusis: radicalibus petiolatis, caulinis amplexicaulibus. LINN. Sp. 1. Doronicum Romanum. BLACKW. 239. Aconicum Pardalianches. DOD. Purg. 305. In montibus Helvetiae. P.

PLANIPETALAE S. CICHORACEAE.

Flosculi omnes aequales lingulati androgyni, apice quinquedentato. Plantae huius classes omnes fauciatae lac stillant, amarae, saponaceae.

I. Semine papposo, placenta nuda.

TARAXACVM. HALL. 739. Leontodon.
LINN. 817.

Taraxacum pinnis foliorum triangularibus, calyce inferne reflexo. HALL. Helv. 1. Leontodon. LINN. Sp. 1. Dens Leonis, Taraxacum. BLACKW. T. 1. C. B. 1. R. H. Vbique in gramineis. A.

Taraxacum foliis hispidis triangulariter dentatis, caule erecto hispido, pappo plumoso. HALL. H. 6.

D d 5

Leon-

lium. I. R. H. Hier. *pilosum*, foliis variis dentatis. DILL. *Elth.* T. 149.

Hieracium foliis glabris amplexicaulibus dentatis, caule paniculato, calycibus hispidis. LINN. *Sp.* 16. HALL. *H.* 20. Hierac. *mont. latifolium glabrum minus*. C. B. In pratis vdis. v. g. *um die Papiermühle*.

H. Hieracium caule ramoso, foliis firmis, infimis petiolatis, reliquis ex ovato-lanceolatis: omnibus sinuosis, petiolorum instar dentatis. GMEL. *Sib. II.* p. 16. T. 24. Crepis LINN. *Sp.* 9. In *Sibiria*. P.

CREPIS. LINN. 819.

• Crepis foliis ad terram serratis, ad caulem pinnatis, pinnis alterne maioribus; caule nudo pene nudo multifloro. Hieracioides. HALL. *Helv.* 2. Crepis. LINN. *Sp.* 10. Hieracium *maius erectum angustifolium*, caule laevi. C. B. I. R. H. In agris.

Crepis (*Hieracioides*) foliis semipinnatis, subasperis amplexicaulibus, caule hispido folioso & multifloro. HALL. *H.* 3. Crepis *foliis bastato-lyratis*, caule piloso. LINN. *Sp.* 12. Hier. *Chondrillae folio birsutum*. C. B. In agris & hortis.

• Crepis foliis glabris integris & varie incisis, caule ramosissimo, floribus minimis. HALL. *H.* 4. Crepis. LINN. *Sp.* 11. Hieracium *chondrillae folio glabrum*. C. B. In hortis.

• Crepis foliis ovatis, caule brachiato, floribus perexiguis. HALL. *G.* 411. Hieracioides *annua* gl.

glutinosa, floribus parvis. VAILL. *Act. Par.* 1721.

Crepis foliis hirsutis pinnatis, pinnis angulatis alterne maioribus. HALL. *H.* 1. LINN. *Sp.* 13.
Hieracium amygdalas amaras olens, f. odore Hier. Apuli suave rubentis. I. R. H. *Über der Papiermühle*, in *M. Pless* sub arce.

INTYBVS. HALL. *Comment. Gott.* 1751, p. 219.

Habitus Hieracii, sed placenta alveolata, & flores umbellati.

Intybus caule folioso, foliis ovato-lanceolatis, dentatis hirsutis. Hieracium. HALL. *H.* 16. LINN. *Sp.* 25. Hier. *frutic. folio subrotundo.* C. B. Hieracii *Sabaudi varietas altera.* I. B. II. p. 1030. In sylvis.

Intybus caule folioso, foliis amplexicaulibus ellipticis acutis dentatis. HALL. *G.* 419. *Helv.* p. 747. Hierac. *frutic. latifolium hirsutum.* C. B.

Intybus caule folioso, foliis linearibus lingulatis dentatis firmis. HALL. *Comm. Gott.* 1751. p. 219. Hieracium. *Helv.* 22. LINN. *Sp.* 26. Hier. *fruticosum angustifolium maius.* I. R. H. In dumetis.

β. Var. *foliis angustis vix dentatis.*

PICRIS. LINN. 812.

Picris foliis amplexicaulibus lingulatis dentatis, perianthiis imbricatis. HALL. *H.* 751. LINN. *Sp.* 2. Hieracium *asperum maiori flore in agrorum limitibus.* I. R. H.

HEL-

HELMINTHOTHECA. VAILL. HALL.

H. Helminthotheca. HALL. G. 413. *Picris perianthiis involacro maximo cinctis*. LINN. Sp. 1. *Hieracium echioides*, *capitulis Cardui benedicti*. C. B. A.

SONCHVS. LINN. 813.

• *Sonchus pedunculis tomentosis, calycibus glabris*. LINN. Sp. 4. *Sonchus foliis levibus teneris, calycibus glabris*. HALL. H. 3. *larvis laciniatus latifolius*. C. B. In hortis. A.

• *β. Sonchus foliis rigidiusculis ferratis, calycibus laevibus*. HALL. H. 2. *Sonchus asper laciniatus & non laciniatus*. C. B. In cultis. A.

• *Sonchus pedunculis calycibusque hispidis subumbellatis, foliis basi cordatis semipinnatis*. LINN. Sp. 3. HALL. H. 1. *Sonchus repens, multis Hieracium maius*. I. R. H. In vdis segetibus. A.

H. *Sonchus pedunculis calycibusque hispidis subumbellatis, foliis basi sagittatis: palustris*. LINN. Sp. 2. *Sonchus asper arborescens*. C. B.

• *Sonchus pedunculis squamosis, floribus racemosis*. LINN. Sp. 6. HALL. H. 4. *Sonchus laevis laciniatus coeruleus alpinus*. C. B. In sylvis Hercyniæ circa M. Bruclerum. P.

H. *Sonchus pedunculis subsquamosis, floribus racemosis, foliis lyrato-hastatis*. LINN. Sp. 7. *Sonchus Canad. altissimus laciniatus, flore coerulescente*. VAILL. Act. Par. 1721. p. 198.

H. *Sonchus pedunculis squamatis, foliis lanceolatis indivisis sessilibus*. LINN. Sp. 8. GMEL. Sib. II.

p. 11. T. 3. *Lactuca Salicis folio, flore coeruleo.*
AMMANN. Ruth. 211. In Sibiria.

PRENANTHES. LINN. 816.

Prenanthes flosculis quinis, foliis lyrato-hastatis.
LINN. Sp. 4. HALL. H. 755. n. 1. *Lactuca sylv.*
murorum, flore luteo. I. B. II. p. 1004. *Sonchus*
laevior vulgaris. II. CLVS. CXLVI. In sylvis.

CHONDRILLA. LINN. 815.

Chondrilla. LINN. Sp. 796. HALL. H. 755. a
Chondrilla *iunceae viminea arvensis.* TABERN. 487.
C. B. In apricis. v. g. über der Maschbrücke.

LACTVCA. LINN. 814.

H. *Lactuca foliis rotundatis, caule corymboso.*
sativa. LINN. Sp. 2. BLACKW. T. 88. *capitata.* B.
non capitata. γ. *crispa.* In hortis cultura varie
mutatur. A.

H. *Lactuca foliis ovatis dentatis, costa spinosa.*
HALL. Gott. 415. *Lactuca sylv. Italica, sanguineis*
maculis adspersa. HERM. Par. p. 194. ic. *Lactuca*
sylv. odore viroso. C. B. I. R. H.

Lactuca foliis semipinnatis, costa spinosa. HALL. •
H. 3. *Lactuca sylv. costa spinosa.* C. B. I. R. H. In
vallo vrbis. *Lactuca foliis carina aculeatis dentatis.*
LINN. Sp. 3. hanc cum priori iungit.

H. *Lactuca foliis lingulatis & semipinnatis, costae*
principio spinoso. HALL. G. l. c. *saligna* LINN. Sp.
5. *Lactuca sylv. altera, angusto saligno folio, costa*
albi-

albicante. HALL. *Rapp. Jen.* 207. T. 4. In *Thuringia*. A.

• H. *Lactuca* foliis glabris pinnatis, caule brachiato. HALL. *H.* 2. LINN. *Sp.* 6. *Chondrilla coerulescens*. TABERN. p. 176. In *Thuringia*.

SCORZONERA. LINN. 811.

Scorzonera foliis angustissimis semipinnatis & integris, caule folioso. HALL. *H.* 2. *Scorz. foliis linearibus dentatis*. LINN. *Sp.* 6. *Scorz. foliis laciniatis*. I. R. H. Ad vias. v. g. circa *Gescheröder Feld*.

β. *Scorzonera* foliis semipinnatis, caulibus erectis ramosis & foliosis. HALL. *G.* 416.

Scorzonera foliis semipinnatis, caule pene nudo, pappo sessili. HALL. *H.* 4. *Leontodon*. LINN. *Sp.* 4. *Hieracium minus praemorsa radice*. LOB. *ic.* 237. *Hierac. Chondrillae & Coronopi folio*. C. B. I. R. H. In pratis & gramineis. v. g. gegen der *Maschmühle*.

h. *Scorzonera* foliis gramineis firmulis, caule brachiato multifloro. HALL. *G.* 417.

H. *Scorzonera* caule ramoso, foliis amplexicaulis integris serrulatis. LINN. *Sp.* 2. *Scorz. latifolia sinuata*. C. B. BLACKW. T. 406. *Scorz. maior* Hiff. L. CLVS. CXXXVII. In *Hispania*. *Sibiria*.

H. *Scorzonera* caule subnudo, foliis nervosis planis ovato-lanceolatis. LINN. *Sp.* 1. *Mat. med.* 371. *Scorz. humilis latifolia nervosa*. C. B. I. R. H. CLVS. CXXXVIII. In pratis & caeduis sylvarum. v. g. circa *Berolinum*. Spontanea caule uniflora, in hortis culta multiflora.

H. *Scorzonera* foliis omnibus antrorsum dentatis amplexicaulibus, squamis calycis ovatis alatis. LINN. Sp. 7. *Chondrilla Tingitana*, floribus luteis, *popaveris hortensis folio*. HERM. Lugdh. p. 659. ic.

H. *Scorzonera* foliis glaucis, interioribus semipinnatis, superioribus amplexicaulibus integerrimis, squamis calycis cordatis alatis. LINN. Sp. 8. *Crepis*. HALL. G. 413. Utraque modo dicta dubii generis.

TRAGOPOGON. LINN. 810.

Tragopogon calycibus corollae radium aequantibus, foliis carinatis amplexicaulibus graminibus. LINN. Sp. 1. HALL. H. 758. *Tragopogon prut. luteum majus*. C. B. Y. R. H. In vallo vrbis & pratis.

β. Var. flore viridi. Im Göttinger Wald.

H. Tragopogon calycibus corollae radio longioribus, foliis integris strictis, pedunculis superne incrassatis. LINN. Sp. 3. *Tragop. purpureo-coeruleum Perri folio*. C. B. *Tragop. crocifolium mont. flore nigro purpureo*. COLVMN. Ecphr. p. 230. ic.

IK. Tragopogon calycibus corollae radio longioribus, foliis integris, seminibus laevibus, disci plumosis, radii setaceis. LINN. Sp. 4. H. Vpf. 243. *. *Tragop. gramineo folio, flore suaverubente*. COLVMN. Ecphr. I. T. 23 IV. A.

III. Tragopogon calycibus corollae radium aequantibus, foliis lineari-ensiformibus integris carinatis. *Scorzonera*. LINN. Sp. 2. *Scorz. Laxitanica gramineo folio, flore pallide luteo*. BYXB. Cent. II. p. 26. T. 21.

H. *Tragopogon calycibus corolla brevioribus inermibus, foliis lyrato-sinuatis*. LINN. *Sp.* 6. *H. Vpsl.* 344. *Chondrilla prior*. CLVS. CXLIH. A.

H. *Tragopogon calycibus corolla brevioribus aculeatis, foliis pinnato-hastatis*. LINN. *Sp.* 7. HALL. G. 419. *Sonchus asper laciniatus Creticus*. C. B. *Prodr.* 60. A.

H. *Tragopogon calycibus corolla brevioribus hispida, foliis scabris caulinis oblongis, caule ramosissimo*. LINN. *Sp.* 8. *Sonchus asper subrotundo folio, maior & minor*. C. B. *Prodr.* 60. *Folia aspera, margine aculeis exiguis scabra. Folia calycis simplici serie, pilis rigidis hispida, accessoriis ad basin folioli vagis laxis. Semina omnia incurva ex brevissimo stylo papposa, exteriora foliis carinatis calycis involuta pappo breviori*. A.

Placenta pilosa, semine papposo.

ANDRYALA. LINN. 820.

H. *Andryala foliis integris*. LINN. *Sp.* 1. *Eriophoros foliis dentatis, flore extus melino, intus subrubente, & utrinque luteo*. VAILL. 1721. p. 277. *Sonchus lanatus*. DALECH.

Placenta paleacea, semine papposo.

HYPOCHAERIS. LINN. 822.

H. *Hypochaeris foliis dentato-fimbriatis scabris, caule ramato nudo, pedunculis squamosis*. LINN. *Sp.* 6. HALL. H. 1. *Hieracium dentis leonis folio obtusa maius*. C. B. *Hierac. intybaceum*. TABERN. 489. In sylvis & gramineis.

Hypo-

Hypochaeris glabra, calycibus oblongis imbricatis, caule nudo, foliis glabris dentato-sinuatis. LINN. Sp. 3. *Hypochaeris chondrillae folio, parvo flore.* VAILL. 1721. p. 14. *Hyoseris altera.* TABERN. 490. In graminis M. Hagen.

Hypochaeris caule subnudo, ramo solitario, foliis ovato-oblongis dentatis hirsutis. LINN. Sp. 4. *Hieracium alp. latifolium, flore magno.* C. B. HALL. H. n. 2. T. 24. *Hierac. latifolium.* CLVS. CXXXIX. In pratis bercynicis circa Elbingerode.

H. *Hypochaeris* calycibus aequalibus hispidis. *Achyrophorus.* LINN. Sp. 2. *Rhagadiolus.* HALL. G. 422. n. 3. In Creta. A.

Semine foliis coronato.

CATANANCHE. LINN. 824.

H. *Catananche* squamis calyceinis inferioribus lanceolatis. *lutea.* LINN. Sp. 2. *Chondrilla cyanoides lutea, coronopi folio non diviso.* BARR. ic. 1135. In Creta. A.

H. *Catananche* squamis calycinis inferioribus ovatis. *coerulea.* LINN. Sp. 1. *Chondrilla coerulea cyan capitulo.* C. B. *Chondrilla.* III. DOD. Pempt. 638. In Provincia. A.

HYOSERIS.

Semina exigua recta setis rigidis paucioribus coronata, in ambitu saepe nuda.

H. *Hyoseris* scapis unifloris nudis, foliis glabris lyrato-hastatis angulatis, seminibus omnibus setis coronatis. LINN. Sp. 2. ROYEN. 127. *Taraxacum*

Ee 2

nastrum

nastrium *dentis leonis folio ad summum radice*.
VALL. Paris. 131. p. 180. In Gallia Narbonensi.

H. *Hyoseris* caule folioso ramofo, calyce levis
 longissimis circumvallato, seminibus in ambitu
 dis. *Rhagadiolus* HALL. 421. *Crepis barbata*.
 LINN. *Sp. 4. Hieracium calyce barbato. COLUM.*
Epbr. II. p. 28. Ic. Hier. Baeticum medio nigra.
 HERM. *Par. p. 185. Ic. In Italiae maritimis.*

RHAGADIOLVS.

Semina longa cylindrica, in ambitu foliis caly-
 cis obvoluta, corona ciliata pecta, in medio levis
 coronata.

H. *Rhagadiolus* capitulo subgloboso. squamis ca-
 lycis seminigeris longioribus hispidis. HALL. G. 422.
An Hyoseris. LINN. Sp. 6? Squamae calycis dorso
undique scabrae. Setae seminum albae genicula an-
gustatae.

H. *Rhagadiolus* capitulo ovato, squamis calycis
 seminigeris longioribus, inferne glabris, apicem
 versus dorso scabris. *Hyoseris* LINN. *Sp. 7. Lapsana*
H. Ppf. 6. Squamae calycis glabrae, apicem ver-
sus serrato-denticulatae, setae seminum subsistae
capillares.

H. *Rhagadiolus* capitulo depresso angulato, squa-
 mis calycis seminigeris brevioribus, glabris, gibbis,
 nodosis. HALL. G. 422. *Lapsana* LINN. *Sp. 2. Ci-*
chorium verrucarium, Zacintha. CLVS. CXLIV.

LAPSA NA LINN. 823.

Semina omnia nuda.

Lapsana calycibus fructus angulatis, pedunculis tenuibus ramosissimis. LINN. Sp. 1. Lapsana foliis imis pinnatis, superioribus lanceolatis serratis.

HALL. Helv. 1761. Lapsana. I. R. H. DODON. Pempt. 675. In sylvis caeduis & locis ruderosis.

Lapsana calycibus fructus depressis, caule nudo diviso, foliis ovatis dentatis. HALL. G. 423. Hyoseris LINN. Sp. 5. Hyoseris angustifolia. TABERN. 180. ic. Hieracium minimum. CLVS. CXLII. In arenosis Germaniae.

H. A. Lapsana calycibus fructus vndique patentibus, radiis subulatis, foliis lanceolatis indivisis. stellata. LINN. Sp. 3. Hieracium Norbogensis falcatum. EOB. ic. 240.

H. Lapsana calycibus fructus vndique patentibus, radiis subulatis, foliis lyratim, lobis rotundatis. LINN. Sp. 4. H. Vps. 245. Rhadagioma Lapsanae folia. LINN. 1753. ic. 245.

CICHORIVM. LINN. 825.

Cichorium caule simplici, foliis dentato-grenatis, pinnis triangularibus. LINN. Sp. 1. Cichorium sylvestre f. Officinarium. C. B. I. R. H. BLACKW. T. 183. Vbique ad oras agrorum.

H. Cichorium caule simplici, foliis integris crenatis. LINN. Sp. 2. Cichorium latifolium f. Endivia vulgaris. C. B. Intybum sativum. DOD. Pempt. 634. A.

H. *Cichorium caule dichotomo spinoso*. LINN.
Sp. 3. Cichorium spinosum. C. B. *Prodr. p. 62. ic.*
Chondrillae genus elegans, coeruleo flore. CLYS.
 CXLV. *Ad litora maris Siciliae.* A.

SCOLYMUS. LINN. 816.

H. *Scolymus foliis margine attenuatis*. LINN.
Sp. 1. Scolymus Theophrasti, Narbonensis. CLYS.
 CLIII. *In Europa australi.*

A P P E N D I X.

OMISSA, ET ADDENDA.

Pag. 11. post. l. 19.

CERATONIA. LINN. 983.

H. *Ceratonia*. LINN. *Sp. p. 1026. Mat. med. 455.*
Silqua edulis, sicuti Carobe filigata. BLACKW. T.
 209. *In Europa australi. Planta diphyta, fr. legu-*
men magnum compressum.

Pag. 250. post. l. 26.

PETIVERIA. LINN. 417.

Semina sex, styli quatuor.

H. *Petiveria alliacea*. LINN. *Sp. p. 342. Aff.*
Stoeckh. 1744. p. 287. T. 7. Petiveria Solani foliis,
luculis spinosis. PLUM. *Gen. 50. In Jamaica,*

Pag. 26. post. l. 15.

anethodoh i: RIVINIA. LINN. 1900.

H. Rivinia. LINN. *Spec. p. 121. Rivinia humilis racemosa, baccis puniceis.* PLUM. *Gen. 48. Solanoides Amer. Circææ foliis.* TOURN. *Art. Par. 1706.*

Pag. 3. post. l. 10.

PHYTOLACCA. LINN. 321.

H. Phytolacca baccis petiolatis, decem striis notatis. HALL. *G. 211. vulgaris.* DILL. *H. Elth. T. 239. f. 309.* Virginiae indigena, nunc sponte in Italia provenit. P.

H. Phytolacca baccis sessilibus, octo striis notatis. HALL. *G. 212. DILL. Elth. T. 308. Flores subvirides. Phytolacca foliis integerrimis.* LINN. *Sp. p. 441. vtramque iungit. P.*

P. 22. post. l. 7.

2. CANNA. LINN. 1. Cannacorea. T.

Stamen unicum & stylus unicus figura laciniæ coronæ.

H. Canina foliis ovatis vtrinque acuminatis nervosa, flore luteo punctato. LINN. *Sp. 1. qui hanc & sequentem iungit. A. B. H. R. T. III. P.*

H. Canna foliis ovatis vtrinque acuminatis nervosis, flore coccineo splendente. HALL. *G. 66. R. K. H. R. V. T. 110. Commer. Litt. Nor. 1743. p. 290. Tab. II. f. 3. fgg. P.*

Pag. 91.

1211.1001.01.257

Genera *Bravelia* & *Haemadathus* poni debuissent
post *Narcissus*, p. 95.

Pag. 118: post l. 18.

H. *Thalictrum* seminibus pendulis triangularibus,
stipulis caulinis, caule tereti. *aquilegifolium*. LINN.
Sp. 12. HALL. H. 309. *Thalictrum Aquilegiae fe-*
lis, *florum staminibus purpureis aut albis*. I. R. H.
In *Helvetia*, P.

P. 125: post l. 15.

VERATRVM. LINN. 1013. HALL.

H. *Veratrum* racemo composito, corollis patentissimis. LINN. Sp. 2. GMEL. Sib. I. p. 76. *Veratrum flore atrorubente*. HALL. G. 87. P.

P. 149: post l. 23.

. *Cistus suffruticosus procumbens*, stipulis lanceolatis, foliis oblongis revolutis subpilosissimis. LINN. Sp. 26. *Helianthemum foliis subhirsutis longe ellipticis*. HALL. H. 1. *Helianth. vulgare, flore luteo*. C. B. I. R. H. In siccis apricis. P.

. *Cistus herbaceus erectus hirsutus stipulatus*, floribus foliariis subsessilibus, folio ternato opposito. *Caulis simplex vna cum foliis & calyce hirsutus. Folia petiolata lanceolata linguata integerrima, stipulis magnis lanceolatis foliis similibus, ut folia ternata dici possint; superiora foliaria singula opposita floribus secundis, brevi pedunculo innixis.*

An

An *Cistus salicifolius*, LINN. *Sp.* 19. ? *Cistus annuus* L. CLVS. 76. Icon sequenti magis convenit. A.

H. *Cistus herbaceus* patulus villosus stipulatus, floribus solitariis alaribus & terminalibus, stipulis minimis. *Priori ex habitu similis, sed in eodem licet solo multoties minor. Stipulae angustissimae. Flores alterni, plerique tamen cauliculi uniflori.* A.

Pag. 275. post. l. 17.

FRAXINVS. LINN. 1026.

Fraxinus foliolis ferratis, floribus apetalis. excellior. LINN. *Sp.* 1. HALL, *H.* 167. C. B. I. R. H. In sylvis & sepibus.

P. 312. post. l. 16.

GLECHOMA. LINN. 643. *Chamaeclema*.

T. HALL.

Glechoma (Chamaeclema) caulibus procumbentibus radicatis, foliis reniformibus crenatis. HALL, *H.* 1. LINN. *Sp.* 2. *Hedera terrestris vulgaris.* C. B. BLACKW. *T.* 225. Ad sepes & sub arboribus.

β. *Chamaeclema flore magno.* HALL. *H.* 2. *Hedera terrestris montana.* C. B. In vallo suburbano. Num Varietas.?



5

1. The first of these is the fact that the
2. second of these is the fact that the
3. third of these is the fact that the
4. fourth of these is the fact that the
5. fifth of these is the fact that the
6. sixth of these is the fact that the
7. seventh of these is the fact that the
8. eighth of these is the fact that the
9. ninth of these is the fact that the
10. tenth of these is the fact that the

Chrysomelidae

1. The first group of people who are not allowed to enter the country are those who are not citizens of the United States and who are not permanent residents of the United States. This group includes all foreign-born individuals who are not citizens or permanent residents of the United States.

1. The first step in the process is to identify the problem or issue that needs to be addressed. This involves gathering information and understanding the context of the problem.

- HOLLYWOOD, CALIF. (AP) - A major Hollywood strike is expected to begin today as the Screen Actors Guild and the Screen Writers Guild vote to join the 100,000-member Writers Guild of America in a walkout against the television industry.
- The strike would be the largest in the history of the entertainment industry, and could cost the industry billions of dollars in lost revenue.
- The strike would also affect the production of new television series, and could lead to the cancellation of existing series.
- The strike would also affect the production of new motion pictures, and could lead to the cancellation of existing films.
- The strike would also affect the production of new television series, and could lead to the cancellation of existing series.
- The strike would also affect the production of new motion pictures, and could lead to the cancellation of existing films.
- The strike would also affect the production of new television series, and could lead to the cancellation of existing series.
- The strike would also affect the production of new motion pictures, and could lead to the cancellation of existing films.


abricato, radii petalis







3 2044 102 804







3 2044 102 804 267

